

# Informative Inventory Report

about Belgium's annual submission of air emission  
data reported in February 2015 under the  
Convention on Long Range Transboundary Air  
Pollution CLRTAP

March 2015



**Prepared by:**

Flemish Environment Agency (VMM)

Alfons Van de Maelestraat 96

9320 Erembodegem

Belgium

Telephone +32 (0)53 72.67.19

Fax +32 (0)53 72.66.19

<http://www.vmm.be>

E-mail: [h.pien@vmm.be](mailto:h.pien@vmm.be), [i.vanvynckt@vmm.be](mailto:i.vanvynckt@vmm.be), [c.debosscher@vmm.be](mailto:c.debosscher@vmm.be), [e.debrabanter@vmm.be](mailto:e.debrabanter@vmm.be),  
[r.vercruysse@vmm.be](mailto:r.vercruysse@vmm.be), [v.cornelis@vmm.be](mailto:v.cornelis@vmm.be), [m.vandenhende@vmm.be](mailto:m.vandenhende@vmm.be)

Walloon Agency for Air and Climate (AWAC)

Walloon Public Service

Av. Princes de Liège 7

5100 Jambes

Belgium

Telephone +32 (0) 81 33.59.33

Fax +32 (0) 81 33.59.32

<http://www.airclimat.wallonie.be>

E-mail: [isabelle.higuet@spw.wallonie.be](mailto:isabelle.higuet@spw.wallonie.be), [andre.guns@spw.wallonie.be](mailto:andre.guns@spw.wallonie.be),  
[camille.vercruysse@spw.wallonie.be](mailto:camille.vercruysse@spw.wallonie.be), [julien.hoyaux@spw.wallonie.be](mailto:julien.hoyaux@spw.wallonie.be)

Brussels Environment (IBGE-BIM)

Gulledelle 100

1200 Brussels

Belgium

Telephone +32 (0)2 775.76.81

Fax +32 (0)2 775.76.21

<http://www.ibgebim.be>

E-mail: [fgoor@environnement.irisnet.be](mailto:fgoor@environnement.irisnet.be), [mcadena@environnement.irisnet.be](mailto:mcadena@environnement.irisnet.be)

Belgian Interregional Environment Agency (IRCEL-CELINE)

Avenue des Arts/Kunstlaan 10-11

1210 Brussels

Belgium

Telephone +32 (0)2 227.56.72

Fax +32 (0)2 227.56.99

<http://www.irceline.be>

E-mail: [vanpoucke@irceline.be](mailto:vanpoucke@irceline.be)

Environment, nature and energy department (LNE)

Graaf de Ferrarisgebouw

Koning Albert II-laan 20, bus 8

1000 Brussel

Belgium

<http://www.lne.be>

E-mail: [steven.lauwereins@lne.vlaanderen.be](mailto:steven.lauwereins@lne.vlaanderen.be)

## Table of contents

Executive Summary .....	vi
Chapter 1. Introduction .....	1-1
1.1. National inventory background .....	1-1
1.2. Institutional arrangements .....	1-2
1.3. Inventory preparation process .....	1-4
1.4. Methods and data sources .....	1-5
1.5. Key categories .....	1-7
1.5.1. Level assessment .....	1-7
1.5.2. Trend assessment .....	1-10
1.5.3. Summary of key category analysis .....	1-12
1.5.4. General remarks .....	1-15
1.6. QA/QC and Verification methods .....	1-18
1.7. General uncertainty evaluation .....	1-18
1.8. General assessment of completeness .....	1-20
Chapter 2. Explanation of key trends .....	2-1
2.1. National total emission trends .....	2-1
2.2. Trends/Time series inconsistencies: general explanations .....	2-3
2.3. Trends in key sectors of main pollutants, CO, PM10, Pb, dioxins and PAH .....	2-5
2.3.1. NO <sub>x</sub> .....	2-5
2.3.2. NMVOC .....	2-6
2.3.3. SO <sub>x</sub> .....	2-8
2.3.4. NH <sub>3</sub> .....	2-9
2.3.5. CO .....	2-10
2.3.6. PM10 .....	2-11
2.3.7. Pb .....	2-13
2.3.8. Dioxins and furanes .....	2-14
2.3.9. PAHs .....	2-15
Chapter 3. Energy (NFR sector 1) .....	3-1
3.1. Overview .....	3-1
3.2. Energy industries (1A1) .....	3-1
3.2.1. Source category description (1A1) .....	3-1
3.2.2. Methodological issues .....	3-2
3.3. Manufacturing Industries and Construction (1A2) .....	3-17
3.3.1. Source category description (1A2) .....	3-17
3.3.2. Methodological issues .....	3-18
3.4. Transport (sector 1A3, 1A5b and off-road) .....	3-30
3.4.1. Source category description .....	3-30
3.4.2. Methodological issues .....	3-31
3.4.2.1 Road transport (1A3b) .....	3-31
3.4.2.2 Air transport (1A3a) .....	3-32
3.4.2.3 Railways .....	3-33
3.4.2.4 Navigation .....	3-34
3.4.2.5 Other transportation (pipeline compressors 1A3ei and off-road 1A3eii) .....	3-36
3.5. Other sectors (sector 1A4) .....	3-37
3.5.1. Source category description (1A4) .....	3-37
3.5.2. Methodological issues .....	3-37
3.6. Other (category 1A5a and 1A5b) .....	3-47
3.7. Fugitive emissions from fuels (category 1B1 and 1B2) .....	3-48
3.7.1. Solid fuel transformation (category 1B1b) .....	3-48

3.7.2.	Fugitive emissions from oil (category 1B2a) .....	3-48
3.7.3.	Natural gas (category 1B2b).....	3-49
3.8.	Recalculations and planned improvements .....	3-50
3.9.	QA/QC .....	3-51
Chapter 4.	Industrial processes (NFR sector 2) .....	4-1
4.1.	Source category description .....	4-1
4.2.	Methodological issues .....	4-2
4.2.1.	Mineral products (category 2A) .....	4-2
4.2.1.1	Cement production (2A1).....	4-2
4.2.1.2	Lime production (2A2).....	4-3
4.2.1.3	Quarrying and mining of minerals other than coal (2A5a) .....	4-4
4.2.1.4	Other mineral products (2A6) .....	4-4
4.2.2.	Chemical industry (category 2B) .....	4-5
4.2.2.1	Ammonia production (2B1) .....	4-5
4.2.2.2	Nitric acid production (2B2) .....	4-5
4.2.2.3	Other chemical industry (2B10a) .....	4-6
4.2.3.	Metal production (category 2C) .....	4-6
4.2.3.1	Iron and steel production (2C1) .....	4-6
4.2.3.2	Other metal production (2C7c).....	4-7
4.2.3.3	Storage, handling and transport of metal products (2C7d) .....	4-7
4.2.4.	Solvent and product use .....	4-8
4.2.4.1	Domestic Solvent Use (category 2D3a) .....	4-8
4.2.4.2	Road paving with asphalt (2D3b).....	4-8
4.2.4.3	Coating Application: Construction, Building and Domestic Use, car reparing, wood, manufacture of automobiles, other industrial and non-industrial application (category 2D3d).....	4-8
4.2.4.4	Chemical Products, Manufacture and Processing (NFR 2D3g).....	4-13
4.2.4.5	Printing (category 2D3h).....	4-17
4.2.4.6	Application of Glues and Adhesives (category 2G/2D3i).....	4-18
4.2.4.7	Preservation of Wood (category 2G/2D3i) .....	4-18
4.2.4.8	Fat, edible and no-edible oil extraction (category 2D3i).....	4-19
4.2.5.	Pulp and paper (2H1) .....	4-19
4.2.6.	Food and drink (2H2).....	4-19
4.2.7.	Consumption of POPs and heavy metals (category 2K) .....	4-20
4.2.7.1	The use of PCB transformers and capacitors .....	4-20
4.2.8.	Other production, consumption, storage, transportation or handling of bulk products (category 2L) .....	4-21
4.3.	Recalculations .....	4-22
4.4.	QA/QC .....	4-23
4.5.	Planned improvements .....	4-23
Chapter 5.	Agriculture (NFR sector 3) .....	5-1
5.1.	Overview .....	5-1
5.1.1.	Allocation of emissions .....	5-1
5.1.2.	Description of the sector .....	5-1
5.1.3.	Climate: .....	5-1
5.1.4.	Data sources .....	5-1
5.1.4.1	Livestock .....	5-2
5.1.4.2	N-excretion factors .....	5-3
5.2.	Animal husbandary and manure management (category 3B) .....	5-5
5.2.1.	NH <sub>3</sub> .....	5-5

5.2.2.	Particulate matter.....	5-6
5.2.3.	NO <sub>x</sub> .....	5-7
5.2.4.	NMVOC .....	5-7
5.3.	Direct soil emission (category 3D).....	5-7
5.3.1.	Synthetic fertilizer use (category 3Da1).....	5-7
5.3.1.1	NH <sub>3</sub> .....	5-7
5.3.1.2	NO <sub>x</sub> .....	5-8
5.3.1.3	NMVOC & PM .....	5-8
5.3.2.	Manure processing (category 3Dd) .....	5-8
5.3.3.	Urine and dung deposited by grazing animals (category 3Da3) .....	5-9
5.4.	The use in pesticides (4G) .....	5-9
5.5.	Recalculations .....	5-10
5.6.	Planned improvements .....	5-10
Chapter 6.	Waste (NFR sector 5) .....	6-11
6.1.	Solid waste disposal on land (category 5A) .....	6-11
6.2.	Waste incineration (category 5C).....	6-11
6.3.	Wastewater treatment (category 5D).....	6-14
6.4.	Other.....	6-14
6.5.	Recalculation .....	6-14
6.6.	QA/QC .....	6-15
6.7.	Planned improvements .....	6-15
Chapter 7.	Land use and land use change .....	7-1
Chapter 8.	Other and natural emissions .....	8-1
8.1.	Biogenic emissions.....	8-1
Chapter 9.	Recalculations and improvements.....	9-1
9.1.	Recalculations and improvements in the energy sector .....	9-1
9.2.	Recalculations and improvements in the sector of industrial processes and products use	9-2
9.3.	Recalculations and improvements in the agricultural sector.....	9-4
9.4.	Recalculations and improvements in the waste sector .....	9-5
Chapter 10.	Projections .....	10-1
10.1.	Mobile sources .....	10-1
10.2.	Stationary sources .....	10-1
10.2.1.	Flanders: Flemish energy and greenhouse gas simulation model .....	10-1
10.2.2.	Walloon region .....	10-2
10.2.3.	Brussels capital region .....	10-3
Chapter 11.	Reporting of gridded emissions and LPS .....	11-1
Chapter 12.	Adjustments.....	12-1
12.1.	Adjustments - summary .....	12-1
References	.....	a
Annex 1: Key category analysis.....		- 1 -
1.A Level assessment .....		- 1 -
1.B Trend assessment.....		- 15 -
Annex 2: NMVOC emissions and TIER methods used in the Solvents sector.....		- 18 -

## Executive Summary

The Belgian Informative Inventory Report (IIR) is the descriptive report that accompanies the Belgian emission inventory of air pollutants submitted by 15 February 2015 under the Convention on Long Range Transboundary Air Pollution (CLRTAP) of the United Nations Economic Commission for Europe (UNECE).

This report follows the recommended structure for Informative Inventory Report (Annex II to the revised 2013 Guidelines). It provides background information on institutional arrangements for inventory preparation, methodologies, data sources, emission factors used, QA/QC activities, key source analyses, trend analyses, recalculations and improvement plans. Furthermore, for each sector more detail is given on the methodologies and assumptions made for estimating the Belgian air emissions. The improvement of the IIR is a constant and progressive work. This year, adaptations conform the new reporting template are added. In 2014, a stage 3 review was performed for Belgium. Recommendations made in this review are integrated in the emission inventory and the present IIR.

The emission data presented in this report were compiled according to the recommendations of the Guidelines for Estimating and Reporting Emission Data under CLRTAP (ECE/EB.AIR/97) revised in 2014 (ECE/EB.AIR/125). For the reporting, the new NFR14 templates provided by the EMEP Centre on Emission Inventories and Projections<sup>1</sup> were used. The submission of 15 February 2015 contains emissions from 1990, 1995, 2000, 2005 and 2010 to 2012 (recalculated) and 2013 (first reporting). In the resubmission of 15 March 2015 also the activity data were added to the inventory. Moreover, emission projections for 2020, 2025 and 2030 are provided. The 2015 submission includes emission data of the pollutants covered in the Convention and its Protocols. These are the main pollutants (NO<sub>x</sub>, SO<sub>x</sub>, NMVOC, NH<sub>3</sub>, CO), particulate matter (PM<sub>2.5</sub>, PM<sub>10</sub> and TSP), heavy metals (Pb, Cd, Hg, As, Cr, Cu, Ni, Se, Zn) and persistent organic pollutants (POPs – PCDD/PCDF, PAHs, HCB, PCB). Belgium reported also the black carbon emissions.

For the first time, Belgium reported road transport emissions based on fuels sold. However, the emissions based on fuels used will be used for compliance checking in accordance with the EMEP reporting guidance.

### **Main differences from last submission - recalculations**

- To continue the quality improvement, the notation keys for the NEC pollutants (NO<sub>x</sub>, NMVOC, SO<sub>x</sub> and NH<sub>3</sub>) and the other LRTAP pollutants (CO, particulate matter, heavy metals, POPs) have been revised. Special attention was given this year to 'IE'.
- For the road transport sector, the modelling is harmonized between the three regions. A common fleet module and the same module for the assignment of mobility are used as input for the COPERT4 v11 model applied in the three regions. For the first time, emissions based on fuels sold were reported.
- Emissions of NO<sub>x</sub> and NMVOC from the agricultural sectors: manure management and agricultural soils are implemented in the emission inventory since this submission.
- As a result of the in depth review (Stage 3) of the Belgian LRTAP inventory in June 2014, several suggestions made by the review experts, regarding allocation matters, gap filling, further explanations and interpretations to enhance the transparency, as well as the inclusion of an uncertainty analysis are taken into account in the current IIR.

---

<sup>1</sup> [www.ceip.at](http://www.ceip.at)

- As a result of the amended Gothenburg Protocol, BC is reported in the three regions.

### **Plans for improvement**

In spite of the intensive work that is already carried out and is still going on to improve estimates of air emissions in Belgium, it remains a continuous task of all members involved to improve the quality of emission inventories. The tuning and information exchange between the regions is taken care of in the bosom of the CCIEP Working Group 'Emissions'. Additional to the 'routine' consultation moments concerning the practical harmonisation of emission calculations between the regions, emphasis is laid on the reduction of the notation key 'NE'.

A tool that calculates the emissions of relevant air pollutants from aviation, according to the most recent guidelines for the international and European reports, will be developed (including exhaust and non-exhaust emissions).

The methodology to estimate emissions from the residential sector (1A4bi) will be upscaled from Tier 1 to Tier 2. Particularly the emission factors of the EMEP/EEA emission inventory guidebook 2013 (tier 2) will be used for wood burning in Flanders and Wallonia.

# Chapter 1. Introduction

## 1.1. National inventory background

The increasing problems of transboundary air pollution led to the signature of the Convention on Long Range Transboundary Air Pollution (CLRTAP) by the United Nations Economic Commission for Europe (UNECE). This Convention was adopted in November 1979 in Geneva and is ratified by Belgium in July 1982. The Convention came into force in March 1983.

The CLRTAP, together with the 8 Protocols that followed, is a framework for international scientific collaboration and policy negotiation to combat air pollution including long range transboundary air pollution. The 51 member parties to the CLRTAP commit themselves to develop policies and strategies to reduce air pollutants which threaten human health and ecosystems. The different Protocols that followed the Convention aim at the reduction of specific pollutants like SO<sub>x</sub>, heavy metals, POPs, and emissions leading to acidification, eutrophication and ground level ozone. Table 1-1 gives an overview of the ratification status of Belgium to the Convention and its Protocols.

Table 1-1 Belgian ratification status on the CLRTAP and its Protocols

<b>Convention on Long Range Transboundary Air Pollution</b>	<b>Signature</b>	<b>Ratification</b>
1979 CLRTAP	13/11/1979	15/07/1982
<b>Protocol</b>	<b>Signature</b>	<b>Ratification</b>
1984 EMEP Protocol	25/02/1985	5/08/1987
1985 Sulphur Protocol	9/07/1985	9/06/1989
1988 NO <sub>x</sub> Protocol	1/11/1988	31/10/2000
1991 VOC Protocol	19/11/1991	31/10/2000
1994 Sulphur Protocol	14/06/1994	31/10/2000
1998 POP Protocol	24/06/1998	8/06/2005
1998 Heavy Metals Protocol	24/06/1998	25/05/2006
1999 Gothenburg Protocol	4/02/2000	13/09/2007

In order to fulfil the obligations of the Protocols under the Convention, annual reporting of emission data to the Executive Body of the Convention on Long Range Transboundary Air Pollution is required.

The Belgian national emission data reported under CLRTAP are established using the Guidelines for Estimating and Reporting Emission Data under CLRTAP (ECE/EB.AIR/97), revised in 2014 (ECE/EB.AIR/125) for application in 2015 and subsequent years. The revised Nomenclature For Reporting (NFR14) was used as template for the reporting. The submission of the Belgian emission

inventory under CLRTAP of 2015 contains emission and activity data of the years 1990, 1995, 2000, 2005, 2010 to 2012 (recalculated) and 2013 (new). Currently a new datawarehouse is developed in the Flemish region. This will lead to an improvement in consistency and completeness of the whole time series of the national inventory. This year, a new inventory software was developed in Wallonia (WAPI) improving the quality of the regional and the national inventory.

The Belgian inventory contains emission estimates for NO<sub>x</sub>, SO<sub>x</sub>, NMVOC, NH<sub>3</sub>, CO, particulate matter (PM<sub>2.5</sub>, PM<sub>10</sub>, TSP, BC), heavy metals (Pb, Cd, Hg, As, Cr, Cu, Ni, Se, Zn), dioxins, PAH, HCB and PCBs.

The key information needed to establish the emission inventories are energy balances (at regional level), national statistics, annual reports of industrial facilities, transport statistics, etc. For several sectors (in particular key sources) national or regional methodologies are developed to give the best emission estimates. Other methodologies and emission factors are taken from the EMEP/EEA Air Pollutant Emission Inventory Guidebook.

## **1.2. Institutional arrangements**

In the Belgian federal context, major responsibilities related to environment lie with the regions. Compiling atmospheric emissions inventories is one of these responsibilities. Each region implements the necessary means to establish their own emission inventory in accordance with the EMEP/EEA Emission Inventory Guidebook. The emission inventories of the three regions are subsequently combined to compile the national atmospheric emission inventory. Since 1980, the three regions have been developing different methodologies (depending on various external factors) for compiling their atmospheric emission inventories. During the last years important efforts are made to harmonise these different methodologies, especially for the most important (key) sectors. Obviously, this requires some coordination to ensure the consistency of the data and the establishment of the national inventory. This coordination is one of the permanent tasks of the Working Group on 'Emissions' of the Coordination Committee for International Environmental Policy (CCIEP), where the different actors decide how the regional data will be aggregated to a national total, taking into account the specific characteristics and interests of each region as well as the available means. This working group consists of representatives of the 3 regions and of the federal public services. The Belgian Interregional Environment Agency (CELINE - IRCEL) is responsible for integrating the emission data from the inventories of the three regions and for compiling the national inventory.

The Inter-ministerial Conference for the Environment is one of the permanent working groups of the Concerted Action Committee and is composed of representatives of the several Belgian governments authorized for environmental matters. Decisions that have an impact on all regions are discussed and taken in consensus to guarantee a coherent Belgian policy.

Since environmental policy is a very specific matter, the federal estate and the 3 regions have entered into a cooperation treaty (5 April 1995, publication in the Belgian Law Gazette on 13 December 1995) on international environmental policy within the scope of the Inter-ministerial Conference for the Environment. A preliminary coordination prior to the Belgian position at international fora is necessary given the complexity of the Belgian competence distribution. The cooperation treaty provides for the establishment of the Coordination Committee of International Environmental Policy (CCIEP). The CCIEP is composed of representatives of the federal and the regional administrative departments and the governmental services with environmental competences. Consistent with the cooperation treaty and depending on particular needs, the CCIEP establishes expert working groups, with a specific mandate, e.g. to discuss and harmonise emission data. All matters related to the national emission

inventory (compilation, harmonisation between the regions, information exchange,...) are discussed during regular meetings of the CCIEP Working Group on Emissions.

Entities responsible for the performance of the main functions of the Belgian National Inventory System, as well as main institutional bodies in relation with the decision process as regards this system, are presented hereafter.




Figure 1.1 Overview of the entities responsible for the constitution and performance of the Belgian Inventory System

As decided by the legal arrangements, the 3 regions are responsible for delivering their atmospheric gas inventories, which are then compiled to produce the Belgian inventory. The main regional institutions involved are :

- The Department Air, Environment and Communication of the Flemish Environment Agency (VMM) in the Flemish Region (Flemish emissions) and Flemish Environment, Nature and Energy Department (DLNE) (Flemish projections);
- The Walloon Agency for Air and Climate (AWAC) in the Walloon Region;
- Brussels Environment (IBGE-BIM) in the Brussels Capital Region.

Each region has its own legal and institutional arrangements, which are detailed in the NIS (2010).

The institutions involved in the constitution (compilation and coordination) of the national emission inventory are:

- The Working group on Emissions of the Coordination Committee for International Environmental Policy (CCIEP) (referred to below as 'CCIEP-WG Emissions') plays a central role in the coordination of the national atmospheric emissions inventory.
- The Belgian Interregional Environment Agency (IRCEL-CELINE) is the single national entity with overall responsibility for the preparation of the Belgian atmospheric emissions inventory. IRCEL-CELINE operates as national compiler of the emissions inventories in Belgium.

### 1.3. Inventory preparation process

The regional atmospheric emissions inventories and projections are transmitted by 1 December (NEC pollutants) and by 1 February (all pollutants for LRTAP reporting) in the shape of NFR-tables to IRCEL-CELINE, the national inventory compiler. IRCEL-CELINE compiles the three regional inventories into the national one, in the right template by 20 December (NEC) and by 10 February (LRTAP). This implies coordination with all regions, within the context of the CCIEP-WG Emissions. The compiled data are fed back to the regions for cross-check. After approval by the regions, the data are submitted to the EU Commission via the Permanent Representation of Belgium to the European Union by 31 December (upload to CDR with notification mail and officially sent to the EC) and to the UNECE secretariat by 15 February (upload to CDR with notification mail to the UNECE secretariat). An overview of the inventory preparation process in Belgium is given in Figure 1.2.




Figure 1.2 Overview of the inventory preparation process in Belgium

## 1.4. Methods and data sources

As a consequence of the responsibility of the regions in preparing the emission inventories, concomitant methodologies have been developed by the three regions for compiling their inventory from basic data. Where it is possible, the existing methodologies are tuned within the regions. When optimisation of a methodology or the development of new methodologies occur, the regions aim at the use of the same methods. This section describes per sector the general approach developed by each region. The text box below gives some more specific detail on the data sources per region.

The emissions of the **industrial sector** (including the **waste** sector) are obtained from the annual industrial reports, submitted by the plant manager to the competent authorities. When this detailed information is not available, the emission data from this sector are based on calculations using the EMEP/EEA methodology or on plant specific emission factors (see also text box below). Energy data are provided in the regional energy balances of Flanders, Wallonia and Brussels.

To have a total picture of all emissions by industrial activities, also activities with emissions below the threshold (see text box Flanders for more information) have also to be taken into account. These emissions are estimated in a collective way. The collective estimation of the emissions is done by multiplying the appropriate activity data with default emission factors.

A detailed description of the methodologies used in the energy and industrial sectors is given in Chapters 3 and 4, the waste sector is described in Chapter 6.

Emissions by **heating systems of buildings** are calculated on a collective basis. A distinction is made between emissions due to residential combustion (heating by households) and tertiary combustion (heating by hotels/restaurants, medical services, education, offices and administrative activities, trade, other). Emissions are calculated by multiplying the energy use and emission factors. A more detailed description of the methodologies used can be found in 3.5.2.

The methodologies that are used to calculate **transport** emissions are described in 3.4. Emissions of road transport are calculated with a harmonized methodology between the 3 regions (based on COPERT IV v11.1). Air transport emissions are calculated by emission factors from the EMEP/EEA Guidebook or other internationally accepted emission factors. The emissions of railway traffic are estimated by a region specific approach. Flanders uses the EMMOSS model, whereas emissions in Wallonia and Brussels are calculated by multiplying the train's fuel consumption by fuel specific emission factors. Emissions from maritime navigation (only in Flanders) are calculated with the emission model EMMOSS.

**Off-road** emissions are calculated by the same mathematical model OFFREM (Off-road emission model) in the three regions. Emissions are calculated for machinery used in industry and building (category 1A2gvii), for machinery used in defence (category 1A5b), in harbours, airports and transshipment companies (category 1A3eii), in households (category 1A4bii) and in agriculture, forestry and green area (category 1A4cii). Exhaust emissions as well as non-exhaust emissions are calculated. Activity data as input for the model are data from the energy balance, statistics from harbours and airports, information about households and data on sales of machinery.

In Belgium the emissions of NMVOC in the source category '**Solvent and other product use**' come from a number of subsectors. The regions in Belgium are using comparable methodologies to estimate the emissions of solvent and other product use in their region. The sector is discussed in detail in Chapter 4.

The **agricultural sector** includes the emissions originating from animal manure, the use of synthetic N-fertilizer, N-excretion on pasture and from manure processing and emissions from agricultural soils. The methodologies that are used to calculate the emission data are given in detail in Chapter 5. The main activity data are the livestock figures, N-excreted and amount of synthetic fertilizer use. In

Flanders, the EMAY-model is used to calculate the emissions for the entire time series. In Wallonia, the emissions are calculated using a model developed by a consultant agency Siterem.

More detailed information on emissions due to fuel use in the agricultural sector (category 1A4c) is included in Chapter 3 Energy (3.5.2). Stationary emissions (1A4ci) are calculated by multiplying the activity data (energy consumption data from the regional energy balances) of the sector with emission factors (e.g. from the EMEP/EEA Guidebook or region specific emission factors). Emissions by tractors (category 1A4cii) are calculated by EMMOSS or by multiplying the activity data (energy consumption data from the regional energy balances) of the sector with emission factors (from the EMEP/EEA Guidebook).

Although NMVOC emissions of **biogenic** nature are not included in the national total, the methodology is written out in detail in chapter 7 due to the importance of the emissions in absolute figures.

---

#### Data sources per region

##### *Flanders*

*Since the reporting year of 1993 most important industrial companies in the Flemish region in terms of air pollution are obliged to report annually about their emissions when exceeding a threshold value, as defined in Vlarem, the Flemish (regional) environmental legislation. From 2006 on this reporting obligation was harmonized in the Flemish region with the EPER-decision (2000/479/EC) and with the EPRTR-regulation (166/2006/EC). In total nearly 1000 industrial companies are registered in a database, due to this obliged emission reporting by the industrial companies. Mainly for the refineries, iron and steel sectors and the chemical industry (process emissions) this obliged reporting of emissions remains since that time an important source of information for the European and international reporting obligations.*

*Energy data are obtained from the Energy Balance for Flanders, made by the Flemish Institute for Technological Research (VITO, Vlaams Instituut voor Technologisch Onderzoek.)*

##### *Wallonia*

*The emission inventories of the Walloon region are compiled using the EMEP/EEA methodology. Emission factors used, are performed for all industrial sectors. In some cases as agriculture and forestry, the emissions estimates are based on a specific study reflecting the Walloon environment.*

*One main data source for the inventory preparation is the energy balance delivered yearly by the Energy and Sustainable Building Department. The energy balance describes the quantities of energy imported, produced, transformed and consumed in the Walloon Region in a given year. In 2003, an environmental integrated survey has been created which includes all pertinent environment-related reporting requirements for 300 companies. The environmental integrated survey is personalised to the 300 operators of the activities/installations pointed out by one or several regulations (four international Conventions and their protocols, seven European Directives, three European Regulations, two European Decisions, one European Recommendation, two Walloon laws, one Walloon Decree and several non- legally binding agreements).*

##### *Brussels*

*The emission inventory in the Brussels-Capital region is compiled by IBGE-BIM (the Brussels Institute for Environmental Management) using the EMEP/EEA methodology. The emissions are calculated by multiplying activity data by an emission factor. The activity data are mostly coming from the Energy Balance of the region performed annually by the ICEDD (Institut de Conseil et d'Etudes en Développement Durable).*

*The different sectors taken into account in the Brussels emissions inventory reflect the characteristics of a strict urban environment. Nearly all the emissions of this urban region originate from energy consumption (Residential, Commercial and Road Transport).*

---

## 1.5. Key categories

A key category is one that is prioritised within the national inventory system because its emission estimate has a significantly influence, for one or a number of air pollutants, on the level of the national total inventory in terms of the absolute level, or the trend in emissions, or both.

The identification of the key categories is performed according to 'Approach 1' as described in the EMEP/EEA emission inventory guidebook 2013 (see Chapter 2: 'Key category analysis and methodological choice', 2.4.1) for both the level assessment and trend assessment. The key category analysis (level and trend) is performed for all reported gases at the least aggregated level of NFR categories.

### 1.5.1. Level assessment

The level assessment is a quantitative analysis of the magnitude of emissions in one year of each category compared to the total national emissions. For each pollutant, the contribution of each source category estimate to the absolute total national estimate is calculated. The source categories are sorted in descending order of contribution magnitude and then summed together. Source categories are identified as 'key source' when 80% of the national total emissions is covered.

Table 1-2 to Table 1-6 show the results of the level assessment for 2013 for the main pollutants and PM10. For the results of the key source level assessment of all the pollutants, we refer to Annex 1.

Table 1-2 Key source level assessment for NO<sub>x</sub>, 2013

NO <sub>x</sub> Source Code	Source Category	Gg NO <sub>2</sub>	Level ass.	2013 Cum.Total
1A3bi	Road transport: Passenger cars	46.275	22.5%	22.5%
1A3biii	Road transport: Heavy duty vehicles and buses	40.621	19.8%	42.3%
1A4bi	Residential: Stationary	11.868	5.8%	48.1%
1A3bii	Road transport: Light duty vehicles	10.692	5.2%	53.3%
1A1a	Public electricity and heat production	10.34	5.0%	58.4%
2A1	Cement production	8.829	4.3%	62.7%
3Da2a	Animal manure applied to soils	5.654	2.8%	65.4%
2B10a	Chemical industry: Other (please specify in the IIR)	5.516	2.7%	68.1%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	5.027	2.4%	70.5%
3Da1	Inorganic N-fertilizers (includes also urea application)	4.73	2.3%	72.8%
2C1	Iron and steel production	4.264	2.1%	74.9%
1A4ai	Commercial/institutional: Stationary	4.027	2.0%	76.9%
1A3dii	National navigation (shipping)	4.022	2.0%	78.8%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	3.881	1.9%	80.7%

Table 1-3 Key source level assessment for NMVOC, 2013

NMVOC Source Code	Source Category	Gg NMVOC	Level ass.	2013 Cum.Total
3B1b	Manure management - Non-dairy cattle	17.903	13.0%	13.0%
1A4bi	Residential: Stationary	16.971	12.3%	25.4%
2D3a	Domestic solvent use including fungicides	14.888	10.8%	36.2%
2D3d	Coating applications	12.308	9.0%	45.2%
2B10a	Chemical industry: Other (please specify in the IIR)	9.687	7.0%	52.2%
3B1a	Manure management - Dairy cattle	8.304	6.0%	58.3%
2D3g	Chemical products	4.533	3.3%	61.6%
3B3	Manure management - Swine	4.132	3.0%	64.6%
1A3bi	Road transport: Passenger cars	3.448	2.5%	67.1%
1B2aiv	Fugitive emissions oil: Refining / storage	3.11	2.3%	69.3%
1B2b	Fugitive emissions from natural gas (exploration, production, processing, transmission, storage, distribution and other)	3.075	2.2%	71.6%
2H2	Food and beverages industry	2.971	2.2%	73.7%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	2.66	1.9%	75.7%
3B4gii	Manure management - Broilers	2.514	1.8%	77.5%
2D3h	Printing	2.438	1.8%	79.3%
1A3bv	Road transport: Gasoline evaporation	2.359	1.7%	81.0%

Table 1-4 Key source level assessment for SO<sub>x</sub>, 2013

SO <sub>x</sub> Source Code	Source Category	Gg SO <sub>2</sub>	Level ass.	2013 Cum.Total
1A4bi	Residential: Stationary	8.468	18.6%	18.6%
1A1b	Petroleum refining	6.831	15.0%	33.6%
2C1	Iron and steel production	6.024	13.2%	46.8%
2B10a	Chemical industry: Other (please specify in the IIR)	3.71	8.1%	54.9%
2A1	Cement production	3.199	7.0%	62.0%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	2.585	5.7%	67.6%
1A1a	Public electricity and heat production	1.966	4.3%	72.0%
1B2c	Venting and flaring (oil, gas, combined oil and gas)	1.791	3.9%	75.9%
2A6	Other mineral products (please specify in the IIR)	1.756	3.9%	79.7%
1A4ai	Commercial/institutional: Stationary	1.256	2.8%	82.5%

Table 1-5 Key source level assessment for NH<sub>3</sub>, 2013

NH <sub>3</sub> Source Code	Source Category	Gg NH <sub>3</sub>	Level ass.	2013 Cum.Total
3B3	Manure management - Swine	19.962	32.1%	32.1%
3B1b	Manure management - Non-dairy cattle	7.411	11.9%	44.0%
3Da1	Inorganic N-fertilizers (includes also urea application)	7.234	11.6%	55.6%
3Da2a	Animal manure applied to soils	6.632	10.7%	66.3%
3Da3	Urine and dung deposited by grazing animals	5.38	8.6%	74.9%
3B1a	Manure management - Dairy cattle	4.834	7.8%	82.7%

Table 1-6 Key source level assessment for PM10, 2013

PM10 Source Code	Source Category	Gg PM10	Level ass.	2013 Cum.Total
1A4bi	Residential: Stationary	21.255	49.4%	49.4%
2A5a	Quarrying and mining of minerals other than coal	2.376	5.5%	54.9%
1A3bvi	Road transport: Automobile tyre and brake wear	2.002	4.7%	59.6%
1A3bi	Road transport: Passenger cars	1.399	3.3%	62.8%
3B4gii	Manure magement - Broilers	1.395	3.2%	66.1%
3B4gi	Manure magement - Laying hens	1.333	3.1%	69.2%
3Da1	Inorganic N-fertilizers (includes also urea application)	1.113	2.6%	71.8%
1A3bvii	Road transport: Automobile road abrasion	1.096	2.5%	74.3%
3B3	Manure management - Swine	0.974	2.3%	76.6%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.874	2.0%	78.6%
2C1	Iron and steel production	0.791	1.8%	80.5%

Table 1-7 gives an overview of the key source level assessment for 2013. The results of the key source analysis (level assessment) of other years are presented in Annex 1.

Table 1-7 Key source analysis (level assessment) for 2013

2013	Key source categories (sorted from high to low, from left to right)										Total
NO <sub>x</sub>	1A3bi	1A3biii	1A4bi	1A3bii	1A1a	2A1	3Da2a	2B10a	1A4cii	3Da1	80.73%
	22.54%	19.79%	5.78%	5.21%	5.04%	4.30%	2.75%	2.69%	2.45%	2.30%	
NMVOC	2C1	1A4ai	1A3dii	1A2f							80.99%
	2.08%	1.96%	1.96%	1.89%							
SO <sub>x</sub>	3B1b	1A4bi	2D3a	2D3d	2B10a	3B1a	2D3g	3B3	1A3bi	1B2aiv	82.50%
	13.03%	12.35%	10.83%	8.96%	7.05%	6.04%	3.30%	3.01%	2.51%	2.26%	
NH <sub>3</sub>	1B2b	2H2	1A4cii	3B4gii	2D3h	1A3bv					82.68%
	2.24%	2.16%	1.94%	1.83%	1.77%	1.72%					
PM <sub>2.5</sub>	1A4bi	1A1b	2C1	2B10a	2A1	1A2f	1A1a	1B2c	2A6	1A4ai	80.47%
	18.59%	14.99%	13.22%	8.14%	7.02%	5.67%	4.32%	3.93%	3.85%	2.76%	
PM <sub>10</sub>	3B3	3B1b	3Da1	3Da2a	3Da3	3B1a					80.46%
	32.08%	11.91%	11.62%	10.66%	8.65%	7.77%					
TSP	1A4bi	1A3bi	1A3bvi	1A2d	2A5a	1A3biii	1A3bii	1A2gviii			80.62%
	61.95%	4.18%	3.21%	2.54%	2.39%	2.17%	2.12%	1.91%			
BC	1A4bi	2A5a	1A3bvi	1A3bi	3B4gii	3B4gi	3Da1	1A3bvii	3B3	1A2d	81.91%
	49.42%	5.52%	4.65%	3.25%	3.24%	3.10%	2.59%	2.55%	2.26%	2.03%	
CO	2C1										80.62%
	1.84%										
CO	1A4bi	2A5a	1A3bvi	1A3bvii	3B3	1A3bi	3B4gii	3B4gi	1A2gvii	3Da1	80.54%
	39.78%	12.33%	4.68%	3.89%	3.80%	2.48%	2.47%	2.37%	2.04%	1.97%	
CO	2C1	1A2d	2B10a								80.54%
	1.69%	1.65%	1.47%								
CO	1A4bi	1A3bi	1A3bii	1A3biii							80.54%
	41.63%	19.96%	9.95%	9.00%							
CO	2A2	1A4bi	2C1	1A3bi							81.91%
	30.81%	23.55%	20.39%	7.16%							

Pb	2C1 54.50%	1A3bvi 8.63%	2C7c 8.16%	1A4bi 4.52%	1B2aiv 4.12%	1A2a 3.92%					83.85%
Cd	2C7c 34.91%	1A4bi 14.49%	1B2aiv 9.56%	2C1 9.55%	1A1a 6.39%	1B1b 3.44%	1A2d 3.36%				81.71%
Hg	2B10a 18.40%	2A1 14.35%	1B2aiv 13.98%	1A1a 10.36%	2C1 9.69%	1A4bi 7.49%	1A2c 4.02%	2C7c 3.30%			81.58%
As	2C7c 38.17%	1A1a 12.84%	2C1 10.26%	2A1 6.88%	1A2c 5.73%	1A2f 4.38%	1B2aiv 3.27%				81.54%
Cr	2C1 22.55%	1A2a 17.27%	1A3bvi 14.09%	1A4bi 10.97%	1A1a 8.70%	1B1b 3.36%	1A2f 3.26%				80.20%
Cu	1A3bvi 58.31%	1A3c 16.00%	2G 9.31%								83.62%
Ni	1A3bvi 33.59%	2C1 13.80%	1A2c 6.61%	1B2aiv 5.49%	1A2f 4.97%	1A1a 4.62%	1A1b 3.21%	2B10a 2.65%	2A1 2.51%	1A2b 2.37%	82.15%
	2C7c 2.33%										
Se	1A2f 46.22%	1A1a 21.99%	2A1 14.07%								82.28%
Zn	2C7c 22.42%	1A4bi 17.17%	1A3bi 10.72%	1A3bvi 10.47%	2C1 9.31%	1A3biii 4.77%	1A2d 4.18%	1A2a 4.18%			83.24%
PCDD/PCDF	1A4bi 53.10%	5C2 19.42%	2C1 8.99%								81.51%
PAHsTotal1-4	2D3a 43.36%	1A4bi 36.26%	1B1b 14.47%								94.09%
HCB	2L 38.87%	1A1a 27.21%	2A1 17.73%								83.81%
PCBs	2A1 56.76%	2C1 30.56%									87.32%

### 1.5.2. Trend assessment

The trend assessment is a quantitative analysis of the change in emission year to year of each category compared to the change in total national emissions. As emissions for the base year as well as the last year are provided, a trend key category analysis could be performed. The trend assessment identifies categories as key sources when they have a trend that significantly differs from the trend of the national total inventory. Key sources are those categories whose trend differences are, when summed together in descending order of magnitude, cover 80% of the total of all source trend differences.

Table 1-8 to Table 1-12 show the key source trend analyses for the main pollutants and PM10 (base year – 2013). The results for all pollutants are presented in Annex 1.B.

Table 1-8 Key source trend assessment for NO<sub>x</sub>, 1990-2013

NO <sub>x</sub>	Source Code	1990	2013	trend Ass.	% contrib	cum.total
1A1a	Public electricity and heat production	60.73	10.34	0.049	25.20%	25.20%
1A3bi	Road transport: Passenger cars	111.27	46.28	0.023	11.90%	37.10%
1A3bii	Road transport: Light duty vehicles	8.86	10.69	0.015	7.90%	45.00%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	13.56	1.5	0.013	6.70%	51.60%
1A3biii	Road transport: Heavy duty vehicles and buses	73.2	40.62	0.01	5.00%	56.60%
1A4bi	Residential: Stationary	16.53	11.87	0.009	4.50%	61.10%
2B10a	Chemical industry: Other (please specify in the IIR)	5.12	5.52	0.007	3.70%	64.80%
1A4ai	Commercial/institutional: Stationary	3.49	4.03	0.006	2.90%	67.70%
1A3di(ii)	International inland waterways	2.67	3.6	0.006	2.80%	70.50%
2A2	Lime production	2.71	3.55	0.005	2.80%	73.30%
1A4ci	Agriculture/Forestry/Fishing: Stationary	3.2	3.37	0.004	2.20%	75.50%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	6.79	5.03	0.004	2.00%	77.60%
1A3dii	National navigation (shipping)	4.84	4.02	0.004	2.00%	79.60%
2C1	Iron and steel production	5.38	4.26	0.004	2.00%	81.50%

Table 1-9 Key source trend assessment for NMVOC, 1990-2013

NMVOC	Source Code	1990	2013	trend Ass.	% contrib	cum.total
1A3bi	Road transport: Passenger cars	65.03	3.45	0.067	21.70%	21.70%
1A4bi	Residential: Stationary	13.01	16.97	0.034	11.20%	32.80%
2D3a	Domestic solvent use including fungicides	9.41	14.89	0.033	10.60%	43.40%
3B1b	Manure management - Non-dairy cattle	21.45	17.90	0.027	8.80%	52.20%
2D3d	Coating applications	49.72	12.31	0.023	7.30%	59.60%
1A3bv	Road transport: Gasoline evaporation	17.88	2.36	0.014	4.60%	64.20%
1B2aiv	Fugitive emissions oil: Refining / storage	16.93	3.11	0.011	3.50%	67.70%
2D3h	Printing	13.96	2.44	0.009	3.00%	70.80%
1B2av	Distribution of oil products	11.37	1.80	0.008	2.60%	73.40%
3B3	Manure management - Swine	4.54	4.13	0.007	2.20%	75.60%
3B1a	Manure management - Dairy cattle	15.04	8.30	0.007	2.10%	77.70%
2B10a	Chemical industry: Other (please specify in the IIR)	29.56	9.69	0.006	2.10%	79.90%
2H2	Food and beverages industry	2.10	2.97	0.006	2.00%	81.90%

Table 1-10 Key source trend assessment for SO<sub>x</sub>, 1990-2013

SO <sub>x</sub>	Source Code	1990	2013	trend Ass.	% contrib	cum.total
1A1a	Public electricity and heat production	95.2	1.97	0.027	24.90%	24.90%
2C1	Iron and steel production	10.51	6.02	0.013	11.90%	36.80%
1A4bi	Residential: Stationary	31.24	8.47	0.013	11.50%	48.30%
1A4ci	Agriculture/Forestry/Fishing: Stationary	28.41	0.71	0.008	7.10%	55.40%
2A1	Cement production	4.33	3.2	0.007	6.70%	62.10%
1B2c	Venting and flaring (oil, gas, combined oil and gas)	0.25	1.79	0.005	4.40%	66.60%
1A1b	Petroleum refining	40.85	6.83	0.005	4.40%	70.90%
2A6	Other mineral products (please specify in the IIR)	0.46	1.76	0.005	4.30%	75.20%

1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	16.96	0.83	0.004	3.20%	78.40%
1A2e	Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	16.03	1.03	0.003	2.40%	80.90%

Table 1-11 Key source trend assessment for NH<sub>3</sub>, 1990-2013

NH3	Source Code	1990	2013	trend Ass.	% contrib	cum.total
3B3	Manure management - Swine	49.85	19.96	0.055	29.70%	29.70%
3Da1	Inorganic N-fertilizers (includes also urea application)	7.27	7.23	0.029	15.50%	45.20%
3Da2a	Animal manure applied to soils	9.04	6.63	0.016	8.40%	53.70%
3B1a	Manure management - Dairy cattle	11.85	4.83	0.012	6.70%	60.30%
1A4bi	Residential: Stationary	0.71	1.82	0.012	6.60%	67.00%
3B4gii	Manure management - Broilers	1.39	1.92	0.01	5.40%	72.40%
3B4gi	Manure management - Laying hens	4.52	1.26	0.01	5.20%	77.60%
5D1	Domestic wastewater handling	3.36	0.96	0.007	3.80%	81.40%

Table 1-12 Key source trend assessment for PM<sub>10</sub>, 2000-2013

PM10	Source Code	1990	2013	trend Ass.	% contrib	cum.total
1A4bi	Residential: Stationary	11.9	21.26	0.216	37.80%	37.80%
2C1	Iron and steel production	9.67	0.79	0.118	20.60%	58.40%
1A3bi	Road transport: Passenger cars	4.41	1.4	0.035	6.20%	64.60%
1A1a	Public electricity and heat production	2.52	0.3	0.029	5.10%	69.70%
1A3biii	Road transport: Heavy duty vehicles and buses	2.43	0.73	0.02	3.50%	73.20%
2A1	Cement production	1.32	0.03	0.017	3.10%	76.20%
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	1.33	0.26	0.014	2.40%	78.60%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.2	0.87	0.013	2.20%	80.90%

### 1.5.3. Summary of key category analysis

Key categories are identified by means of their contribution to the national total emissions (level assessment) and according to the difference in their trend compared to the trend of the national total emissions (trend assessment). Key source categories identified by the approach 1 level assessment (L1) or trend assessment (T1) are summarized in Table 1-13.

Table 1-13 Key category analysis for 2013 based on level (L1) or trend (T1) assessment

2013	NOx	NM VOC	SOx	NH3	PM2.5	PM10	TSP	BC	CO	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	PCDD/ PCDF	PAH	HCB	PCB	#
1A1a	L1,T1		L1,T1		T1	T1	T1				L1	L1,T1	L1,T1	L1	T1	L1,T1	L1,T1		T1		L1,T1		14
1A1b			L1,T1													L1							2
1A2a	T1								T1	L1		T1		L1,T1				L1,T1					6
1A2b															T1	L1							2
1A2c			T1								T1	L1,T1	L1,T1			L1,T1	T1						6
1A2d					L1,T1	L1,T1	L1				L1,T1							L1,T1					5
1A2e			T1													T1							2
1A2f			L1								T1		L1	L1		L1	L1,T1	T1					7
1A2gviii					L1						T1					T1					T1		4
1A3bi	L1,T1	L1,T1			L1,T1	L1,T1	L1,T1	L1,T1	L1,T1	T1								L1,T1					9
1A3bii	L1,T1				L1			L1															3
1A3biii	L1,T1				L1,T1	T1	T1	L1,T1										L1					6
1A3bv		L1,T1																					1
1A3bvi					L1	L1	L1			L1,T1				L1,T1	L1,T1	L1,T1		L1,T1					8
1A3bvii						L1	L1																2
1A3c															L1,T1								1
1A3di(ii)	T1																						1
1A3dii	L1,T1																						1
1A4ai	L1,T1		L1																				2
1A4bi	L1,T1	L1,T1	L1,T1	T1	L1,T1	L1,T1	L1,T1	L1,T1	L1,T1	L1,T1	L1,T1	L1,T1		L1,T1				L1,T1	L1,T1	L1,T1			16
1A4ci			T1																				1
1A4cii	L1,T1	L1																					2
1B1b						T1	T1	T1			L1,T1			L1							L1,T1		6
1B2aiv		L1,T1								L1,T1	L1	L1	L1			L1							6
1B2av		T1																					1
1B2b		L1																					1
1B2c			L1,T1																				1
2A1	L1		L1,T1			T1	T1					L1,T1	L1,T1			L1	L1,T1				L1,T1	L1,T1	10
2A2	T1						T1		L1,T1			T1											4
2A5a					L1	L1	L1,T1																3
2A6			L1,T1																				1
2B10a	L1,T1	L1,T1	L1				L1				T1	L1				L1		T1					8
2C1	L1,T1		L1,T1		T1	L1,T1	L1,T1		L1	L1,T1	L1,T1	L1,T1	L1,T1	L1,T1	T1	L1		L1,T1	L1,T1	T1		L1,T1	17
2C7c										L1	L1	L1,T1	L1,T1			L1		L1,T1					6
2D3a		L1,T1																		L1,T1			2

2D3d		L1,T1																				1
2D3g		L1																				1
2D3h		L1,T1																				1
2H2		L1,T1																				1
2L																				L1,T1		1
3B1a		L1,T1		L1,T1																		2
3B1b		L1,T1		L1																		2
3B3		L1,T1		L1,T1		L1	L1															4
3B4gi				T1		L1	L1															3
3B4gii		L1		T1		L1	L1															4
3Da1				L1,T1		L1	L1															3
3Da2a	L1			L1,T1																		2
3Da3				L1																		1
5C1a																			T1			1
5C1bi															T1							1
5C1bv											T1											1
5C2																			L1,T1			1
5D1				T1																		1

#### 1.5.4. General remarks

To evaluate the key sources in time, the level assessments for the base years 1990 (all pollutants except particulate matter) and 2000 (particulate matter) are actualized as well. The summary of these key source analyses can be found in Annex 1.

The absolute change in emission values of key source categories for a particular pollutant over the period 1990-2013 will be discussed in Chapter 2.

By comparing the key sources (level assessment) between 1990 and 2013 some remarks can be made. Besides some (smaller) shifts in the order of ranking, a number of more structural shifts in the following sectors can be seen:

- *1A1a Public electricity and heat production*: disappears as a key source for Cu, Zn, dioxins and particulate matter in 2013 compared to 1990. Also for SO<sub>x</sub>, NO<sub>x</sub> and the other metals, the relative contribution of this sector to the national total decreases. Emissions of NO<sub>x</sub>, SO<sub>x</sub>, particulate matter and (heavy) metals decrease with the termination of some coal power plants, the use of environment-friendlier fuels (minimal use of liquid fuels, application of renewable sources), the higher efficiency of existing plants and the application of new technologies.
- *1A1b Petroleum refining*: disappears for NO<sub>x</sub> and PM<sub>2,5</sub> in 2013 compared to 1990. In 1990, the refining plants (all situated in the Flemish region) were not yet obliged to report their emissions (obligation from 1993 described in the Flemish environmental law *Vlaem II*). As a result, very little information on emissions at plant level is available before 1993. Emissions were only estimated collectively based on the existing knowledge.
- *1A2a Iron and steel*: appears as a key source in 2013 for Pb, Cr and Zn due to a part of process emissions allocated in the combustion sector.
- *1A2b Non-ferrous metals*: for Cu, the non-ferro industry disappears as key source in 2013 compared to 1990. This is due to the efforts of one *Flemish company with several exploitations*.
- *1A2c Chemicals*: is as a key source in 2013 for Hg, As and Ni. Relative changes in the key sources for heavy metals can be attributed to an optimised methodology that is applied from 2000 on in Flanders.
- *1A2d Pulp, paper and print*: appears as a key source for particulate matter in 2013 compared to 2000 due to the increased use of renewable solid fuels.
- *1A2e Food processing, beverages and tobacco*: Is no longer a key source for SO<sub>x</sub> and Ni in 2013. The proportion of the Ni emissions from this sector to the national total decreases strongly. The reduction of Ni emissions is due to the reduction of the residual oil as fuel.
- *1A2f Non-metallic minerals*: is a key source for NO<sub>x</sub>, SO<sub>x</sub>, As, Cr, Ni and Se in 2013. For Se it became the largest source of emissions as some polluting plants are now in this sector (cement plants, lime plants and glass plants).
- *1A2gviii Other*: disappears as a key source for Ni and HCB in 2013 compared to 1990 due to a significant decrease in absolute emission values as some polluting plants are now in the sector 1A2f (cement plants, lime plants and glass plants).
- *1A3bi Road transport (Passenger cars)*: Disappears as key sector in 2013 for Pb and has a very much lower relative importance for NMVOC and CO in 2013 compared to 1990 due to the increasing use of catalytic converters and other technical measures. The absolute Pb emissions of passenger cars have strongly decreased due to the removing of leaded petrol. It stays the largest source of NO<sub>x</sub> emissions.
- *1A3biii Road transport (Heavy duty vehicles)*: This is a key sector for NO<sub>x</sub> and particulate matter but no great changes are seen in the relative importance of this sector between 1990 and 2013 for these pollutants.

- *1A3bv Gasoline evaporation:* The relative importance for NMVOC decreases in 2013 due to the decrease of gasoline use between 1990 and 2013.
- *1A3bvi Automobile tyre and brake wear:* this is the most important key source for Cu and Ni emissions. The sector appears as a key source for Pb, Cr and Zn in 2013 compared to 1990. This is due to the increase in mobility and the optimised methodology to estimate heavy metals from the year 2000 on in Flanders. Regarding the Ni, there was an error in the calculation that explains the increase of the emissions, this will be corrected for the next submission.
- *1A3bvii Automobile road abrasion:* The relative importance of the sector increases slightly for PM10 and TSP. This is due to the increase in mobility and so the increase in road distance travelled.
- *1A3c Railways:* appears as a key source for Cu in 2013 compared to the base year. This can be attributed to the optimised methodology to estimate heavy metals since 2000 in Flanders. Cu emissions, due to the abrasion of overhead wires and electric contact withdrawals were estimated in 2013, but not in 1990.
- *1A4ai Commercial/Institutional:* appears as key source in 2013 for NO<sub>x</sub> and SO<sub>x</sub> due to absolute emission changes in other sectors which become relatively less important.
- *1A4bi Residential: Stationary plants:* appears as a key source for Pb, Cd, Hg, Cr and Zn in 2013. Also the relative importance of this sector for NMVOC, SO<sub>x</sub>, CO and particulate matter increases in 2013 compared to 1990. Since the absolute heavy metal emissions remain rather stable, this is mainly due to emission changes in other sectors. There is a significant absolute decrease in dioxin emissions due to the reduction of coal consumption. However, the sector becomes the principal key source of dioxins due to the huge emission decline in the electricity sector and the sector of waste incineration. This sector is also the most important key source for particulate matter due to the high consumption of wood for residential heating.
- *1A4ci Agriculture/Forestry/Fishing: Stationary:* disappears as key source for SO<sub>x</sub> due to the decreasing emissions in the greenhouse culture (more natural gas and less heavy fuel).
- *1A4cii Agriculture/Forestry/Fishing: Off-road vehicles and other machinery:* appears as a key source for NO<sub>x</sub> and NMVOC, due to emission changes in other sectors.
- *1B1b Solid fuel transformation:* appears as a key source for Cd and Cr and the proportion of PAH's emissions to the total PAH's emissions decreases strongly because the activities of the Brussels and Flemish coke ovens have been terminated respectively in 1993 and 1996. The sector disappears as a key source for TSP in comparison with the base year 2000 due to emission changes in other sectors.
- *1B2aiv Refining/storage:* appears as a key source for SO<sub>x</sub>, Pb, Cd, Hg, As and Ni in 2013 compared to 1990. In 1990, the refining plants (all situated in the Flemish region) were not yet obliged to report their emissions (obligation from 1993 described in the Flemish environmental law Vlarem II). As a result, very little information on emissions at plant level is available before 1993. Emissions were only estimated collectively based on the existing knowledge.
- *1B2av Distribution of oil products:* is not longer a key source for NMVOC in 2013 due to the obliged vapour recycling during the refuelling of petrol stations and during tanking (the so-called Stage I and Stage II measures)
- *1B2b Natural gas:* appears as a key source for NMVOC in 2013 due to the change in emissions in other sectors.
- *1B2c Venting and flaring:* appears as a key source for SO<sub>x</sub> in 2013 due to the change in emissions in other sectors
- *2A1 Cement production:* disappears as key source for PM10 and TSP. A significant emission reduction was obtained due to new dust purification systems of some plants in 2008, 2010 and 2012. The sector becomes key source for SO<sub>x</sub>, Hg, As, Se, HCB and PCB in 2013 compared to 1990. It becomes the most important source for PCB emissions due to the large decrease of PCB emissions from the iron and steel sector. The absolute SO<sub>2</sub>

and Hg emissions remain stable between 1990 and 2013 but the emissions of other sectors have decreased.

- *2A2 Lime production:* disappears for Se and TSP and appears as key sector for CO. The CO emissions in the lime production is very important in 2013 (169.9 tons) as the lime production of one plant occurred without oxygen (reducing atmosphere). The change in relative importance of this sector can be attributed to emission changes in the other sectors.
- *2A6 Other Mineral products:* appears as key source for Se in 2013 compared to 1990. This can be attributed to the optimised methodology to estimate heavy metals for the years from 2000 on in Flanders. The sector disappears as key source for PM2.5 and PM10 in 2013 compared to 2000 due to lower emissions from bricks and tiles production. Lower activity data and a lower emission factor were provided by the brick federation.
- *2B10a Other chemical industry:* disappears as a key source for Zn. The emissions in 1990 originate from a number of sulphuric acid production installations in a (Flemish) non-ferro plant. These installations have been shut down in 1992. The sector becomes a key source for NOx, Ni and TSP due to changes in other sectors.
- *2C1 Iron and steel production:* disappears as a key sector for Cu, PAH and HCB in 2013 compared to 1990. For Cu, this is because of a different emission estimation method before and after 1993 (obligation from 1993 for Flemish plants to report their emissions as described in Vlarem II). In the Walloon region, all the blast furnace plants and basic oxygen plants have been closed since 2011. These were an emission source of PAH and HCB. 2C1 appears as a key sector for Hg and remains an important (key) source for most metals and dioxins. This sector remains an important sector in Belgium.
- *2C7c Other metal production:* appears as a key source for Hg and Ni due to the optimised methodology to estimate heavy metals for the years from 2000 on in Flanders. 2C7c disappears as key source for Cu and dioxins due to changes in other sectors.
- *2D3a Domestic solvent use including fungicides:* is a key sector for NMVOC and PAHs. Increasing absolute emissions for NMVOC and slightly increasing emissions for PAHs. Because of the fact that the emission is largely depending on the population, the emission of NMVOC and PAHs increases.
- *2H2 Food and beverages industry:* appears as a key source for NMVOC. This can be attributed to emission changes in the other sectors.
- *2L Other production, consumption, storage, transportation or handling of bulk products:* this sector becomes the most important key source for HCB. The emissions originating from the production of solvents used to be allocated in NFR09 category 3C (solvent use), but it was aimed to better allocate these emissions in NFR14 category 2L, which refers more to production.
- *3B1b Manure management Dairy Cattle:* this becomes the most important key sector for NMVOC, because absolute emissions from the chemical and coating sector decreased strongly since 1990.
- *3B3 Manure management Swine:* stays the most important key source for NH<sub>3</sub> emissions.
- *3D1a Inorganic N-fertilizers:* appears as a key source for NH<sub>3</sub> and TSP due to changes in other sectors.
- *5C1a Municipal waste incineration:* In 1994, this sector has undergone a (structural) reorganisation, which included also air purification measures. Moreover, the majority of the intermunicipal waste incinerators recuperate their energy nowadays. As a consequence their emissions are reported under the sector 1A1a. For dioxins, the sector disappears as key source because of air purification measures.
- *5C1bi Industrial waste incineration:* disappears as a key sector in 2013 compared to 1990 for several heavy metals (Pb, Cd, Hg, Cu, Cr, Ni, Zn) and dioxins. A (structural) reorganisation of the sector in 1994 led to a strong decline of the emissions. From 2006 on intermunicipal waste incinerators recuperate their energy; the emissions are then allocated in sector 1A1a.

- *5C2 Open burning of waste:* appears as a key source in 2013 for dioxins. Emissions in this sector declined less than in other sectors.

It can be assumed that most categories with a notation key NE will not bring big differences in the ranking of the key sources if they would be estimated, since most emissions are relatively low or even not existing. More information on the reasons for not estimating the emissions in a sector are given in 1.8 (table 1.15).

The emissions of the categories that are IE (included elsewhere) are explained in 1.8 (Table 1-16).

## **1.6. QA/QC and Verification methods**

In the framework of the European and international obligations with respect to the greenhouse gas emission inventory, Belgium has developed a QA/QC-plan.

Although this plan is focused on greenhouse gas emissions, a lot of these issues are also appropriate for the air pollutants.

Information about the developed QA/QC-plan in Belgium and all procedures involved can be found in the NIR (National Inventory Report) of March 2014, more specifically in chapter 1.6. 'Information on the QA/QC plan including verification and treatment of confidentiality issues where relevant'.

The three regions have their own QA/QC procedures. The regional inventories are compiled by the Belgian Interregional Environment Agency (IRCEL-CELINE), which is responsible for the international emission reporting obligations. The national inventory compiler is not involved in the development of the regional inventories.

Before compilation at the national level, the regional inventories are again controlled by the national compiler (as an additional control from an external person). The regional emission inventories are compared with the regional inventories used in the former submission, compared with the reporting under the NEC Directive and checked for sudden dips or jumps in the time series. Remarkable results of this review are fed back to the regions in order to obtain confirmation or adjustments on the emission data.

The same control processes are applied for the compiled national inventory. An additional check is made on the consistency in allocation of source categories of the 3 regional inventories. Also a cross-check is performed of the national aggregated data with the sum of the data from the input inventories to ensure that emissions are correctly aggregated from a lower reporting level to a higher reporting level. Any changes in the emission inventory at the national level is conducted by IRCEL-CELINE after coordination with the regional contact persons.

At last, the compiled national inventory is tested with the electronic RepDab-tool, on-line available at the ceip website before submission.

## **1.7. General uncertainty evaluation**

For all emission measurements or estimations, a particular uncertainty can be determined, that is inseparably related to the emission value. In 2014, a study for calculating uncertainty values related to the emissions reported for NEC and LRTAP is conducted in the three Belgian regions by an independent consultant. Uncertainty analysis was done for the emission levels in 2010 and for the 1990-2010 trend in emissions on Tier 1 and Tier 2 level for the pollutants covered in the NEC directive, for the key sectors. Uncertainty for the other LRTAP pollutants was done on Tier 1 level for

the key sectors. The results are available in the technical report 'Uncertainty Analysis of Emission Inventories of NEC/LRTAP Air Pollutants'. The methodology used in this report was the basis for the uncertainty analysis of 2013.

To assess the uncertainty in the air pollutant emission inventory, the methodology provided in the *EMEP/EEA emission inventory Guidebook (2013)* and the *IPCC Guidelines for National Greenhouse Gas Inventories chapter 3 (2006)* were used. The uncertainty calculation is applied on the three regional air pollution emission inventories for the year 2013 and base year-2013 for the trend uncertainty. Subsequently, the uncertainties were aggregated on the national level by the error propagation equation from the Good Practice Guidance, in order to produce one single table 6.1 per pollutant (as expressed in the guidelines).

As most of the data suppliers in Belgium do not provide any information on the associated uncertainty, inventory experts were consulted to give their expert estimation. If this information was not available, either the consortium members' expert judgement was applied or default uncertainties were applied as described in the EMEP/EEA Guidelines.

A comparison of the Tier 1 and Tier 2 results for uncertainty in annual emissions show that there is only a minor difference for the mean emissions. Therefore, no further investments were made for uncertainty calculations on Tier 2 level.

According to the available references, in most member states the ultimate choice of an uncertainty estimate is often based on expert judgement and is therefore also rather uncertain. However, as stressed by the IPCC Good Practice Guidance, uncertainty calculation is a mean to provide inventory users with quantitative judgements on the inventory quality and enables the inventory preparation team to identify and prioritise improvement activities.

The results of the Tier 1 analysis for 2013 for the overall uncertainty per pollutant are given in Table 1-14.

Table 1-14 : Summary of uncertainties in the national total emissions per pollutant

Pollutant	Total Emissions in Base Year	Total Emissions in Reporting Year	Change in total emissions (Reporting Year - Base Year)	Uncertainty in Reporting Year inventory (%)	Uncertainty in trend (Reporting Year - Base Year) (%)
NOx (as NO <sub>2</sub> )	409.90	205.30	-204.60	29.24	3.77
NMVO	341.37	137.43	-203.94	26.39	8.23
SOx (as SO <sub>2</sub> )	365.51	45.56	-319.95	10.69	1.14
NH <sub>3</sub>	117.32	62.23	-55.08	72.99	12.55
CO	1414.64	553.03	-861.61	27.96	19.30
Pb	253.96	26.39	-227.57	275.49	22.50
Cd	6.17	2.32	-3.85	187.66	19.39
Hg	5.68	1.66	-4.02	129.58	17.02
As	6.22	1.62	-4.60	200.71	39.64
Cr	32.85	5.93	-26.92	111.04	18.02
Cu	40.14	31.25	-8.89	220.37	96.00
Ni	74.10	6.78	-67.32	137.38	11.30
Se	4.98	3.52	-1.45	112.39	67.66
Zn	234.34	80.86	-153.47	129.66	35.07
PCDD	576.75	43.08	-533.67	197.58	15.33
Total PAH	79.10	27.94	-51.15	250.98	71.06
HCB	97.82	14.11	-83.71	219.34	14.19
PCB	105.16	4.71	-100.44	323.45	18.49
PM 2.5	41.32	33.45	-7.86	20.19	12.85
PM 10	56.16	43.01	-13.15	20.51	10.01
TSP	78.69	56.37	-22.32	19.30	8.52
BC	8.87	5.54	-3.33	35.91	13.68

## **1.8. General assessment of completeness**

The Belgian emission inventory covers all pollutants of the CLRTAP and its Protocols, i.e. main pollutants (NO<sub>x</sub>, SO<sub>x</sub>, NMVOC, NH<sub>3</sub>, CO), particulate matter (PM<sub>2,5</sub>, PM<sub>10</sub>, TSP, BC), heavy metals (Pb, Cd, Hg, As, Cr, Cu, Ni, Se, Zn) and POP's (PCDD/PCDF, PAH, HCB). Estimations of the individual PAH's could not be made for the Walloon region, so in the national inventory only the totals of the 4 PAH's are reported. In the 2015 submission, recalculations were made for 1990, 1995, 2000, 2005 and 2010 up to 2012. 2013 was reported for the first time.

The Belgian emission inventory covers all relevant sources specified in the CLRTAP. However, it is not always possible to estimate the emissions of all subsectors in detail. Therefore, notation keys have been used. An overview and explanation of the notation keys NE and IE used in the 2013 emission inventory, as well as the sub-sources accounted for in reporting codes 'other' are summarized in Table 1-15 to Table 1-17.

An overview of the basis that is used to estimate emissions from mobile sources (fuels sold versus fuels used) is given in Table 1-18.

Table 1-15 Explanation to the Notation key NE

<b>NFR14 code</b>	<b>Substance(s)</b>	<b>Reason for not estimated</b>
All	benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene, indeno(1,2,3-cd)pyrene	No information available by PAHs specified in Walloon region
1A2f	PCBs	No emission factors available to calculate the emissions
1A2gvii	Hg, As, PCDD/F	No emission factors available to calculate the emissions
1A3ai(i) 1A3aii(i)	NH3, Hg, dioxins, PAH	No emission factors available to calculate the emissions
1A3aii(ii) 1A3ai(ii)	NH3, PM, heavy metals, dioxins, PAH	No emission factors available to calculate the emissions
1A3b (i to iv)	Hg, As, HCB, PCBs	No emission factors available to calculate the emissions
1A3bvi	Hg, As, HCB, PCBs	Hg, As, no emission factors
1A3bvii	heavy metals, HCB, PCB's	Considering the diversity of the road coatings, no estimate was made
1A3di(ii)	Cr, dioxins	No emission factors available to calculate the emissions
1A3ei	NMVOC, SOx, NH3	No data available from the facilities
1A3eii	Hg, As, dioxins	No emission factors available to calculate the emissions
1A4aii	Hg, As, dioxins	No emission factors available to calculate the emissions
1A4bii	Hg, As, dioxins, HCB	No emission factors available to calculate the emissions
1A4ciii	NH3, Cd, Hg, As, Cr, Cu, Ni, Se, Zn, dioxins, PAH's	No emission factors available to calculate the emissions
1A5a	all	There are no data available. The activity data are coming from the military sector (confidentiality).
1A5b	Hg, As, dioxins	No emission factors available to calculate the emissions
1B2aiv	CO	No data available from the facilities
1B2c	NH3, particulate matter	No detailed data available
1B2d	NMVOC, CO	No detailed data available
2A6	NH3, Cu, Zn, PAH	There are no data available or the EF aren't available
2B1	SOx	No data available from the facilities
2B2	CO	There are no data available or the EF aren't available
2B10a	Se, PAH's	No emission factors available to calculate the emissions
2B10b	Heavy metals, dioxins, PAHs	There are no data available or the EF aren't available
2C3	Se, PAHs	No data available from the facilities or no emission factors available

2C5	Se, PAHs	No data available from the facilities or no emission factors available
2C6	Se, PAHs	No data available from the facilities or no emission factors available
2C7a	Se, PAHs	No data available from the facilities or no emission factors available
2C7b	Se, PAHs	No data available from the facilities or no emission factors available
2C7c	NH3	No data available from the facilities or no emission factors available
2C7d	BC, Hg, Ni, Se, dioxins, PAH	No data available from the facilities or no emission factors available
2D3b	NMVOC, dioxins and PAHs	No activity data available
2D3c	NMVOC, particulate matter, CO, dioxins, PAHs	No activity data available
2G	PAHs	No data available from the facilities or no emission factors available
2H1	SOx, NH3, particulate matter	No data available from the facilities or no emission factors available
2H3	all	No activity data available
2I	SOx, BC	No data available from the facilities or no emission factors available
2J	PAHs, HCB, PCBs	No activity data available
2K	Heavy metals, dioxins, PAHs, HCB, PCBs	No data available from the facilities or no emission factors available, POPs emissions probably not relevant
2L	Se, PCBs	No activity data available
3B4giv	particulate matter	No activity data available from 2007 on
3Da2b	NH3	No activity data available
3Da2c	NOx, NMVOC, NH3	No activity data available
3Da4	NOx, NH3	No activity data available
3Db	NOx, NH3	No activity data available
3Dc	NH3, particulate matter	No data available
3Dd	particulate matter	No data available
3De	NOx, NH3	No data available
3Df	Dioxins, PAH, HCB, PCB	No activity data available or no emission factors available
3I	NH3	No activity data available
5A	SOx, NH3, BC, CO	No emission factors available to calculate the emissions
5B1	NOx, particulate matter, CO	No activity data available or no emission factors available
5B2	Particulate matter	No activity data available or no emission factors available
5C1bi	PM2.5, BC, Pb, Cd, Hg, As, Cr, Cu, Se, Zn, PAHs, HCB, PCBs	There are no data available or the EF aren't available; emissions of installations with

		energy recuperation are allocated in 1A1a, POP's emissions probably not relevant
5C1bii	PAHs	There are no detailed data available or the EF aren't available
5C1biii	Dioxins, PAHs	There are no detailed data available or the EF aren't available
5C1biv	Dioxins, PAHs	There are no detailed data available or the EF aren't available
5C1bv	NH3, BC	No emission factors available to calculate the emissions
5C1bvi	all	There are no detailed data available or the EF aren't available
5C2	main pollutants, CO, heavy metals, HCB, PCBs	No emission factors available to calculate the emissions , POP's emissions probably not relevant
5D1	NMVOC	There are no data available or the EF aren't available
5D2	NMVOC, SOx, NH3	There are no data available or the EF aren't available or data not provided by the facility
5D3	Main pollutants, CO	There are no data available or the EF aren't available or data not provided by the facility
6A	all	Should be NA
1A3d i(i)	dioxins, Cr	No emission factors available

Table 1-16 Explanation to the Notation key IE

NFR14 code	Substance(s)	Included in NFR code
1A1c	SOx, particulate matter, heavy metals	1B1b
1A3a ii(i)	heavy metals except Hg	1A3ai(i)
1A4aii	all except Hg, As, HCB and PCBs	1A3eii
1B2ai	NMVOC	1B2av
2A3	all	1A2f
2A5b	particulate matter	2A6
2A5c	particulate matter	2A6
2A6	Heavy metals except Cu and Zn	1A2f
2B1	NMVOC	2B10a
2B6	Particulate matter	2B10a
2B10b	Main pollutants, particulate matter, CO	2B10a
2C3	all except Se, PAHs	2C7c
2C4	all except Se, PAHs	2C7c
2C5	all except Se, PAHs	2C7c
2C6	all except Se, PAHs	2C7c
2C7a	all except Se, PAHs	2C7c
2C7b	all except Se, PAHs	2C7c
2D3e	Pb, As, Cr, Ni	2C7c
2H1	BC	1A2d
2I	Particulate matter	2L
3B4f	NOx, NMVOC, NH3, particulate matter	3B4e
3B4giii	NOx, NMVOC, NH3, particulate matter	3B4gi
3Da3	NOx	3Da2a
5B2	Main pollutants, CO	1A1a
5C1bii	NOx, SOx, NH3, particulate matter, CO, heavy metals	5C1bi or 1A1a (E-recup)
5C1biii	All except dioxins, PAH, PCB	5C1bi or 1A1a (E-recup)
5C1biv	All except dioxins, PAH, PCB	5C1bi or 1A1a (E-recup)
5C1a	NOx, NMVOC, SOx, CO, dioxins	5C1b (Flanders)

Table 1-17 Sub-sources accounted for in reporting codes 'other'

NFR14code	Substance(s) reported	Sub-source description
1 A 2 gviii	all	Non-metallic mineral products, (cement, lime, asphalt concrete, glass, mineral wool, bricks and tiles, fine ceramic materials), metal products, textile, leather and clothing and other industry (wood industry, rubber and synthetic material, manufacturing of furniture, recycling and construction included)
1 A 3 eii	all except Hg, As, dioxins, HCB, PCBs	Off-road emissions of (air)ports
1 A 5 a	-	NE, cfr. Table 1-15, military source
1 A 5 b	all except Hg, As, dioxins, HCB, PCBs	Military aviation in Wallonia and in the Flemish Region+off-road defence
1 B 1 c	-	NO
1 B 2d		
2 A 6	NO <sub>x</sub> , SO <sub>x</sub> , NMVOC, particulate matter, CO, heavy metals, dioxins	Manufacture and processing of flat and hollow glass, glass fibres and other glass, manufacture of bricks, tiles and construction products in baked clay, manufacture of articles of concrete, plaster and other non-metallic products, manufacture of ceramic household and ornamental articles
2 B 10 a	all except Se, PAH, HCB, HCH, PCB's	Production of sulfuric acid, ammonium nitrate, ammonium phosphate, vinylchloride, PEHD, polypropylene, PVC, polystyrene, phthalic anhydride, titanium dioxide, processes in organic chemical industry (excl. adipic acid)
2 B 10 b	pollutants included in 2B5a	IE or NE
2 C 7 c	all except NH <sub>3</sub> , HCB	galvanization, non-ferro
2 C 7d	NO <sub>x</sub> , NMVOC, SO <sub>x</sub> , particulate matter, CO, Pb, Cd, As, Cr, Cu, Zn	metallurgic activities, including (iron) foundries and galvanization activities
2D3i	NMVOC	Process emissions of vegetable oil extraction, gluing, wood preservation, recuperation of waste solvents
2G	NO <sub>x</sub> , NMVOC, SO <sub>x</sub> , particulate matter, CO, Cu, dioxins	application of glues and adhesives, plant oil extraction, wood preservation, recuperation of waste solvents, Flemish estimation of tobacco smoke (PM) and fireworks (Cu)
2H3		NE
2 L	NMVOC, Particulate matter, all heavy metals except Se	construction, manufacture of other non-metallic mineral products including asphalt production, manufacture of man-made fibres, surface treatment and casting of metals, manufacture of fabricated metal products, machinery and equipment, electrical and optical equipment, transport equipment, manufacture of textile and textile products, leather and leather products, manufacture of wood and wood products incl. furniture, manufacture of rubber and plastic products, manufacture of mattresses, recycling of metal and non-metal waste and scraps, industrial cleaning,

3B4giv	NOx, NMVOC, NH3	hens for multiplication and austriches
3B4h	NOx, NMVOC, NH3, particulate matter	rabbits and minks
3 D a2c		NE
3I	-	NE
5C1biv		NE
5D3		NE
5E	all except Zn, dioxins, PAH, HCP, PCBs	Waste recuperation, compost
6 B		NO
11 C	NMVOC	Forest and grassland

Table 1-18 Basis for estimating emissions from mobile sources

NFR09 code	Description	Fuel sold	Fuel used	Comment
1 A 3 a i (i)	International aviation (LTO)		X	
1 A 3 a i (ii)	International aviation (Cruise)		X	
1 A 3 a ii (i)	1 A 3 a ii Civil aviation (Domestic, LTO)		X	
1 A 3 a ii (ii)	1 A 3 a ii Civil Aviation (Domestic, Cruise)		X	-
<b>1A3b</b>	<b>Road transport</b>	<b>X</b>	x	First reporting of emissions of road transport based on fuel sold, emissions based on fuel used are also supplied for compliance purposes,
1A3c	Railways		X	
1A3di (i)	International maritime navigation		X	
1A3di (ii)	International inland waterways		X	
1A3dii	National navigation		X	
1A4ci	Agriculture		X	
1A4cii	Off-road vehicles and other machinery		X	
1A4ciii	National fishing		X	
1 A 5 b	Other mobile (Including military)		X	

## Chapter 2. Explanation of key trends




### 2.1. National total emission trends

The Belgian absolute total emissions per pollutant are summarized in Table 2-1 for the years recalculated in the 2015 LRTAP-submission. The absolute difference as well as the relative difference are calculated between 2013 and the base year. For all pollutants the base year is 1990, except for particulate matter the base year is 2000. The emissions of all pollutants have a downward trend between 1990 (2000) and 2013. Main reasons for this are the great emission reduction efforts made by the industrial and transport sectors as well as the changeover to less polluting fuels. The larger decrease between 2008 and 2009 is mainly due to the crisis that hit the heavy industry in Belgium. Emissions of most pollutants increased again slightly in 2010 after which the reduction is continued in 2011, except for particulate matter, which increased again in 2012 and 2013 due to the cold winter periods.

Figure 2.1 shows the trends of the national total emissions per pollutant group. The years 1991-1994, 1996-1999, 2001-2004 and 2006-2009 were taken from the 2014 LRTAP submission. Those years were not recalculated this year. Reasons for the changes in the time series are given in the next paragraphs.

Table 2-1 Absolute total emissions and absolute and relative differences for the time series 1990-2013

Pollutant	Unit	1990	1995	2000	2005	2010	2011	2012	2013	absolute difference base-2013	relative difference base-2013
NOx	Gg as NO2	410	381	344	317	250	232	214	205	-205	-50%
NM VOC	Gg	341	291	227	186	155	143	141	137	-204	-60%
SOx	Gg as SO2	366	258	174	143	61	53	47	46	-320	-88%
NH3	Gg	117	113	83	69	65	64	63	62	-55	-47%
PM2.5	Gg	NR	NR	41	36	37	30	32	33	-8	-19%
PM10	Gg	NR	NR	56	49	47	39	41	43	-13	-23%
TSP	Gg	NR	NR	79	69	60	52	55	56	-22	-28%
BC		NR	NR	9	8	7	6	6	6	-3	-38%
CO	Gg	1415	1129	942	766	526	423	374	553	-862	-61%
Pb	Mg	254	187	107	75	41	30	30	26	-228	-90%
Cd	Mg	6	5	3	2	3	2	2	2	-4	-62%
Hg	Mg	6	3	3	2	2	2	1	2	-4	-71%
As	Mg	6	6	5	3	2	2	2	2	-5	-74%
Cr	Mg	33	28	18	17	13	11	11	6	-27	-82%
Cu	Mg	40	40	39	37	35	34	33	31	-9	-22%
Ni	Mg	74	69	35	28	12	11	8	7	-67	-91%
Se	Mg	5	6	6	27	11	4	3	4	-1	-29%
Zn	Mg	234	189	183	131	110	104	89	81	-153	-65%
PCDD/ PCDF	g I-Teq	577	399	117	61	53	44	50	43	-534	-93%
PAH	Mg	79	66	43	37	32	31	29	28	-51	-65%
HCB	kg	98	218	73	28	27	40	29	14	-84	-86%
PCB	kg	105	88	93	71	95	57	9	5	-100	-96%






Figure 2.1 Time trends of Belgian total emissions

## 2.2. Trends/Time series inconsistencies: general explanations

Below, some general explanations are given for the occurring inconsistencies and changes in the time trends caused by the changes in emissions of the different sectors.

- In 1993 environmental legislation (Vlarem II) came into force in Flanders. This included a reporting obligation for class 1 industrial plants, which induced in some cases a difference in methodology to calculate/estimate emissions before and after 1993. In 1995 Vlarem II was extended with class 2 industrial plants and thresholds per pollutant. In 2004 the emission reporting (as part of the reporting of environmental data) was established by decree in the integrated annual environmental report (IMJV). The modification of some thresholds can result in the incomparability of emission data from 2004 on compared to the period before 2004 for e.g. some heavy metals. In Wallonia, IPPC plants have had to report their emissions since 2001 and it's sometimes difficult to make a recalculation before 2001 because of the lack of data.
- In Flanders, there is a different level of data handling in some years (1990- 1993, 1995, 1996, 1998, 2000, 2001, 2008-2013) compared to the other years (1994, 1997, 1999, from 2001 to 2007). In the former years emissions are available on installation level (NFR code) whereas in the latter years the emissions are available on a less detailed level (facility/sector level). A partition key based on the most recent detailed data (e.g. for emission data of 1999, the partition used in 1998 is applied for 1999) is used to attribute the emissions to the appropriate NFR code for the year where less detailed emission data are available.
- *Public electricity and heat production:* decrease of the emissions because of the introduction of highly performant power stations, application of technical measures and changeover to natural gas, use of fossil fuels with less sulphur, opting for renewable/less polluting fuels. In the Walloon region, there are no more coal power plant as they were progressively replaced by gas turbines. Emissions of waste incineration with energy recuperation and emissions of CHP installations are allocated to the electricity sector. The decrease of emissions is mainly observed in Flanders. In Flanders, less solid and fluid fuels and more gaseous fuels were used. The use of 'classic' fuels is decreased in 1999 with nearly 9 % compared to 1998, partly due to the good functioning of nuclear units. The choice for a type of fuel depends mostly on the prices and the goals that are assumed in the Environmental Policy Agreement (e.g. coal with a low S content <1%, purchase of

extra heavy fuel with maximal S content of 1 %). Increasing use of natural gas due to better prices and the continuation of the STEG and CHP programme. Installations are modernised and old coal driven installations are replaced by STEG's. Also technical measures were taken to decrease the NO<sub>x</sub> emissions (SCR, specific local measures per installation, old units were closed).

In the Walloon region, a coal power plant was replaced by a gas turbine in 1999.

- *Petroleum refining*: decrease of SO<sub>x</sub> and NO<sub>x</sub> to meet the bubble emission thresholds for 2010 as imposed by the Flemish Government (e.g. by desulphurization of the fuels used or by technical measures). The strong decrease in emissions, particularly from 2008 on, is related to the more stringent emission limit values for SO<sub>2</sub> and NO<sub>x</sub> that became valid on 1 January 2010 as one of the main measures that the Flemish Government has taken in the framework of the European national emission ceilings directive (NECD or National Emission Ceilings Directive). Refineries made heavy investments in purification technology (also of influence on the PM emissions) the years before to be compliant with the NECD. Also a more stringent monitoring and control on the contribution of the emissions through flaring and the switchover of high to low sulphur fuel was mentioned as a measure to lower SO<sub>2</sub> emissions. During the years 2004-2006, one refinery had very limited refining activities
- *Manufacture of solid fuels*: decreasing emissions due to the closure of coke ovens in Brussels and Flanders, respectively in 1993 and 1996 and closure of the last Flemish mines in 1992.
- *Stationary combustion in manufacturing industries*: in general decreasing emission trends between 1990 and 2013 due to important efforts to reduce emissions. The decrease between 2008 and 2009 is mainly due to the crisis in the industry in this period. In category 1A2b strong decrease of some heavy metals because in 1993 a new gas purification installation on a blast furnace of the most polluting facility in this sector reduced strongly the Pb and Cd emissions.
- *Residential sector*: emissions are highly climate related. Fluctuations in emissions can also be attributed to a shift towards natural gas, the increasing number of households (with fewer persons per household), the limited isolation degree of the houses and the low compactness. Emissions of NMVOC and particulate matter increase due to the increased consumption of wood for heating.
- *Commercial/institutional sector*: as for the residential sector, emissions are highly climate related.
- *Road transport*: decrease of emissions of SO<sub>x</sub> due to the use of fuels with low sulphur content (from 2003 on). A significant decline in Pb emissions occurs due to the use of unleaded petrol (from 2000 on), but the emissions of the other heavy metals increase due to a higher fuel use. Due to the enhanced application of catalytic converters NO<sub>x</sub>, CO and NMVOC emissions decrease, but NH<sub>3</sub> emissions increase. More stringent emission standards for diesel cars from 2005 induced lower emissions of particulate matter.
- *Railways*: decreasing emissions due to the gradual change of diesel trains towards less polluting alternatives. Decreasing emissions in particular for freight trains due to increased efficiency (more wagons per engine, better loading, ...).
- *Inland shipping*: decrease of the emissions in 2009 due to the lower economic activity (crisis).
- *Maritime navigation*: gradual increase of emissions of most pollutants due to the expansion of the merchant fleet (number of services and magnitude of ships). Decrease of most emissions in 2009 due to the economic crisis, decrease of SO<sub>2</sub> emissions in international maritime navigation, as determined by the Marpol Convention (more stringent sulphur limits in 2008 and 2010).
- *National fishing*: decreasing emissions due to the scaling down of the sector.
- *Off-road*: decrease of SO<sub>2</sub> and Pb emissions due to the lower S and Pb content of the fuels used.
- *Manure management*: significant decreases of NH<sub>3</sub> emissions in 1991 (Flemish Manure Decree of 23/1/1991), 2000 (MAP 2bis), 2003 (more stringent legislation) and 2007 (MAP 3, particular influence on emissions from cattle). Decrease of NH<sub>3</sub> emissions of poultry in 2003 due to the brake-out of bird flu and the subsequent extermination of poultry by the authorities. From 1990 to 1999 activity data are obtained from the yearly count of cattle, from 2000 on data are available from the Manure Bank of the Flemish Land Agency. In Wallonia, the reduction of livestock is a main driver for the decrease of emissions.
- *Fertilizer*: emissions are related to the amount of fertilizer used (depending on the price) and the type of fertilizer used (liquid fertilizer, ureum,...).

- *Fuel combustion in agriculture*: decreasing emissions (in particular emissions of heavy metals) due to the switchover towards less polluting fuels. Decrease of SO<sub>2</sub> emissions due to the lower S content in gasoil from 2008 on.
- Dioxin emissions of the *metallurgical sector* have decreased significantly due to emission reduction measures.
- *Cement production*: decrease of CO emissions from 2002 onwards as old kiln generating high CO emissions has been stopped in 2002, decrease of dust emissions from 2004 onwards as one plant generating high dust emissions has installed a new filtration system in 2004, PCB emissions in one plant were very high in 2010 and 2011 because of the use of an alternative raw material containing high concentrations of PCB. The removal of the raw material causing high PCB emissions at the end of 2011 has allowed returning to a normal level of emissions.
- *Lime production*: decrease of SO<sub>x</sub> emissions from 2004 onwards as since 2004, there is a progressive reduction of the use of petroleum coke in a lime plant.
- PAH emissions of *wood preservation* have decreased significantly due to emission reduction measures in the sector.
- *Waste incineration*: global emissions have decreased significantly due to the (structural) reorganisation of the sector in 1994, which included also air purification measures. Moreover, in Belgium the emissions of waste incinerators with energy recuperation are reported under the sector 1A1a.
- An optimised methodology to estimate heavy metals emissions in Flanders is applied for the years 2000 to 2013. For some sectors, this might cause an inconsistency between the years before 2000 and the years from 2000 on.

## **2.3. Trends in key sectors of main pollutants, CO, PM<sub>10</sub>, Pb, dioxins and PAH**

A great part of the trend in the absolute total emissions can be explained by the changes in key sector emissions. Therefore, an analysis was made of the key sector emissions throughout the time series for the NEC pollutants NO<sub>x</sub>, NMVOC, SO<sub>x</sub> and NH<sub>3</sub>, for CO and PM<sub>10</sub>.

### **2.3.1. NO<sub>x</sub>**

The greatest contributors to NO<sub>x</sub> emissions are the transport (passenger cars and heavy duty vehicles) and energy sector. The largest absolute emission reductions are made in these sectors. Consequently, this led to the decrease in total NO<sub>x</sub> emissions from 410 Gg in 1990 to 205 Gg in 2013, which is a decline of 50% (figs. 2.2 and 2.3).




Figure 2.2 Trends in NOx emissions for the key sectors




Figure 2.3 Absolute NOx emission differences from 1990 to 2013 for all key sectors

### 2.3.2. NMVOC

The emissions of NMVOC show a steady decrease between 1990 and 2013, from 341 Gg to 137 Gg (-60%, figs. 2.4 and 2.5). The largest absolute emission reductions are made in the transport sector (passenger cars). An explanation is the shift of fuel (gasoline to diesel oil). Other sectors with significant emission reductions are *coating applications* and *Other chemical industry*. A minor increase in the NMVOC emissions over the 24-years period is observed in the *Residential: stationary plants, domestic solvent use* and the *food and beverage industry*.




Figure 2.4 Trends in NMVOC emissions for the key sectors




Figure 2.5 Absolute NMVOC emission differences from 1990 to 2013 for all key sectors

### 2.3.3. SO<sub>x</sub>

SO<sub>x</sub> emissions declined from 366 Gg in 1990 to 46 Gg in 2013, a reduction of 88% (figs. 2.6 and 2.7). This is largely due to the use of fuels with less sulphur in combustion in the energy and manufacturing industries.




Figure 2.6 Trends in SO<sub>x</sub> emissions for the key sectors




Figure 2.7 Absolute SO<sub>x</sub> emission differences from 1990 to 2013 for all key sectors

### 2.3.4. NH<sub>3</sub>

In Belgium, over 90% of the NH<sub>3</sub> emissions are attributed to agricultural activities. Due to the successive Flemish Manure Decrees (1991, 2000, 2003 and 2007), focusing on including manure application and a reduction of the livestock population, the ammonia emissions show a reduction of 47% between 1990 and 2013 (figs. 2.8 and 2.9). In Flanders, more than half this reduction is attributed to the emission reduction of swine. In Wallonia, the decrease of emissions is driven by the reduction of livestock on the one hand and on the reduction of use of mineral fertilizer on the other hand. The latter is linked to the implementation of the Nitrates Directive (EC 91/676) and the Sustainable Nitrogen management program put in place for supervising and advising farmers with their formalities and ensuring compliance with the Directive objectives (<http://www.nitrawal.be/101-documents-anglais.htm>).




Figure 2.8 Trends in NH<sub>3</sub> emissions for the key sectors




Figure 2.9 Absolute NH<sub>3</sub> emission differences from 1990 to 2013 for all key sectors.

### 2.3.5. CO

CO emissions decreased from 1415 Gg in 1990 to 553 Gg in 2013, a reduction of 61% (figs. 2.10 and 2.11). This is mainly due to the reductions realized in the road transport sector and the iron and steel industry. The drop between 2008 and 2009 is mainly due to the closure of some iron and steel plants in Wallonia during 2008 (one coking plant, one sinter plant and one blast furnace plant). There is still one coking plant in Wallonia in 2012. The last sinter plant and the last blast furnace closed in 2011. The sudden increase in 2013 is due to 1 plant where the lime production occurred without oxygen (reducing atmosphere).




Figure 2.10 Trends in CO emissions for the key sectors




Figure 2.11 Absolute CO emission differences from 1990 to 2013 for all key sectors

### 2.3.6. PM10

PM10 emissions between 2000 and 2013 declined with 23%, from 56 Gg to 43 Gg (figs. 2.12 and 2.13). Many sectors contribute to the dust emissions. The sources with the largest absolute emission

reductions are the iron and steel production, road transport (passenger cars and heavy duty vehicles) and the energy sector. The reduction in the transport sector is due to more stringent emission standards for diesel cars. Between 2008 and 2009 the emissions of the iron and steel production have been reduced significantly due to the closure of some iron and steel plants in Wallonia during 2008 (one coking plant, one sinter plant and one blast furnace plant). There is still one coking plant in Wallonia in 2012. The last sinter plant and the last blast furnace closed in 2011.

The emissions from the residential sector increase significantly over the last 14 years due to the increased use of wood for residential heating.




Figure 2.12 Trends in PM10 emissions for the key sectors




Figure 2.13 Absolute PM10 emission differences from 1990 to 2013 for all key sectors

### 2.3.7. Pb

Emissions of Pb decreased strongly between 1990 and 2013 with a decline of 90% from 254 tonnes to 26 tonnes (Figure 2.154 and 2.15). The use of unleaded petrol from 2000 on made Pb emissions originated from road transport exhaust very small. *Iron and steel production* and *other metal production* are the sectors with the greatest emission decreases.




Figure 2.14 Trends in Pb emissions for the key sectors




Figure 2.15 Absolute Pb emission differences from 1990 to 2013 for all key sectors

### 2.3.8. Dioxins and furanes

PCDD-PCDF emissions were high in the early nineties (577 g I-teq), but are greatly reduced in 2013 (43 g I-teq), with a decline of 93% (Figures 2.16 and 2.17). The greatest absolute emission reductions are made in the *energy sector*, the *cement production sector* and the *municipal waste incineration*.




Figure 2.16 Trends in PCDD-PCDF emissions for the key sectors




Figure 2.17 Absolute PCDD-PCDF emission differences from 1990 to 2013 for all key sectors

### 2.3.9. PAHs

Emission of PAHs decreased from 79 tonnes in 1990 to 28 tonnes in 2013, a reduction of 65% (figs. 2.18 and 2.19). This is largely due to reductions in the iron and steel sector. In the Walloon region, all the blast furnace plants and basic oxygen plants have been closed since 2011. PAHs emissions from solid fuel transformations decreased strongly because the activities of the Brussels and Flemish coke ovens have been terminated respectively in 1993 and 1996.




Figure 2.18 Trends in PCDD-PCDF emissions for the key sectors




Figure 2.19 Absolute PCDD-PCDF emission differences from 1990 to 2013 for all key sectors

## **Chapter 3. Energy (NFR sector 1)**

### **3.1. Overview**

This sector includes all combustion emissions (stationary and mobile combustion emissions). Furthermore, it includes fugitive emissions from the energy sector.

The emission data from this sector are based on calculations (fuel consumed x default emission factors) or on direct emission measurements.

To prepare the Belgian inventory for the energy sector, the regional energy balances of Flanders, Wallonia and Brussels are the prime source of activity data.

The main source of information on the industrial emissions is also obtained from the annual industrial reports.

To have a total picture of all emissions by industrial activities, also activities with emissions below the threshold have to be taken into account. These emissions are estimated in a collective way. The collective estimation of the emissions due to combustion processes is done by multiplying the energy data with default emission factors.

### **3.2. Energy industries (1A1)**

#### **3.2.1. Source category description (1A1)**

The energy industries contain the following sectors: the public electricity and heat production, petroleum refining and the manufacture of solid fuels and other energy industries.

The category 'Public Electricity and Heat production (1A1a)' includes fuel combustion emissions associated with the generation of electricity for commercial, industrial or public sale. The emissions of auto-generators are allocated to the category 1A1 (refineries, solid fuel producer), 1A2 'Manufacturing Industries and Construction' and 1A4 'Other sectors', depending on the type of the sector or industry where the energy is used. Some CHP (Combined Heat and Power) units are in joint venture with the energy sector. For these installations, all heat is delivered to the industrial plant and most electricity produced, is sold to the energy sector. In these cases, all fuel in the energy balance and the associated emissions are included in the energy sector, category 1A1a.

The following chart (fig. 3.1) shows the trend of the energy consumption in this sector :




Figure 3.1 Trend of fuel consumption in the energy industries.

The emissions of the refineries, an activity which takes place only in the Flemish region, are allocated in the category 1A1b (combustion emissions) and in the category 1B2a (oil) (diffuse emissions) and in the category 1B2c (flaring emissions). The emissions of CHP of the refinery sector are allocated in 1A1a.

The emissions reported in category 1A1c 'Manufacture of Solid Fuels and Other Energy Industries' are the emissions coming from the combustion in the cokes ovens. Also the emissions of some energetic activities in the mines (mainly an auto-generator) in the Flemish and Brussels region during the beginning of the nineties and emissions due to some gas transport activities are included in this category 1A1c. Fugitive emissions are reported in category 1B1b.

### 3.2.2. Methodological issues

#### **Public electricity and heat production (1A1a)**

This category contains the power installations for the production of electricity and heat, including turbojets, and the (other) combined heat-power (CHP) installations (in joint venture with the electricity producers). These latter installations are located in different sectors in Belgium (refineries, industry, agriculture and service sector). Also included in this sector are the waste incineration installations with energy recuperation (waste incineration installations without energy recuperation are allocated in the sector 5C).

Category 1A1a is a key category of NO<sub>x</sub>, SO<sub>x</sub>, Hg, As, Ni, Se and HCB emissions in terms of emissions level and trend and a key category of PM<sub>2.5</sub>, PM<sub>10</sub>, TSP, Cd, Cr, Cu and PCDD/F emissions in terms of emissions level or trend.

The activity data reported in this sector are the fuel consumption data as reported in the regional energy balances.

The share of the regional emissions of NO<sub>x</sub>, SO<sub>x</sub> and TSP compared to the national emissions for emission year 2013 is presented in the following table (Table 3-1):

Table 3-1 Share to national emissions by regions (%).

	Flanders	Brussels	Wallonia
NO <sub>x</sub>	72.1	3.7	25.9
SO <sub>x</sub>	84.3	3.3	14.6
TSP	22.0	0.4	77.3

The emission data are based on environmental annual reports submitted by the operator of the power plants, the waste incinerators and the industrial plants owning a CHP installation. If the installation is equipped with continuous measuring devices, the SO<sub>2</sub>, NO<sub>x</sub>, TSP and CO emissions are based on continuous analyses in the chimneys.

The emissions of the public power plants are based on continuous measurements for 71 % for NO<sub>x</sub>, 22 % for SO<sub>2</sub> and for 21 % for TSP in Wallonia. Main part of the SO<sub>2</sub> emissions are coming from the combustion of biogas in waste plants (waste disposals, wastewater treatment plants,...) where emission factors are used.

For the estimation of other air pollutants and when there aren't plant specific data or the installation is not equipped with continuous measurement devices, emission factors were used. Emission factors used in the three regions are given below (Table 3-2 to Table 3-20).

Table 3-2 Emission factors for the sector 1A1a Public electricity and Heat Production in the Walloon region (EMEP/EEA Guidebook 2013 – NH<sub>3</sub> : EPA and Emep 1996 – HM for GC and BFG : average of measurements between 2005 and 2011 on boilers)

FUEL	SO <sub>2</sub>	NO <sub>x</sub>	NM VOC	CO	NH <sub>3</sub>	TSP	PM <sub>10</sub>	PM <sub>2,5</sub>	BC
	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ
Natural gas	PS	PS	2.60	39.00	0.60	0.89	0.89	0.89	0.02
Natural gas (in gas turbine)			1.6	4.8	0.60	0.2	0.2	0.2	0.005
GC			2.60	39.00	0.87	2.60	1.18	1.18	0.03
BFG			2.60	39.00	0.60	2.60	1.18	1.18	0.03
Diesel oil			0.80	16.20	0.10	6.50	3.20	0.80	0.27
Diesel oil (in gas turbine)			0.19	1.49	0.10	1.95	1.95	1.95	0.65
Fuel			2.30	15.10	0.10	35.40	25.20	19.30	1.08
Coal	PS	PS	1.00	8.70	0.40	PS	PS	PS	2.2%ofPM <sub>2,5</sub>
Biogas (in stat.engine)	43.70	88	2.50	13	15				
Wood	10.80	81	7.31	90	7	172	155	133	4.39

FUEL	As	Cd	Cu	Cr	Ni	Pb	Sé	Zn	Hg	Dio	HAP (4)	HCB	PCB
	mg/GJ	mg/GJ	mg/GJ	mg/GJ	mg/GJ	mg/GJ	mg/GJ	mg/GJ	mg/GJ	ng/GJ	mg/GJ	yg/GJ	ng/GJ
Natural gas	0.12	0.50	0.40	0.001	0.001	0.002	0.01	0.002	0.10	0.50	0.0031		
Natural gas (in gas turbine)	0.12	0.0003	7.6E-05	0.0008	0.0005	0.002	0.0112	0.0015	0.1		0.0116		
GC	5.40	2.60	5.30	9.00	7.50	8.40	0.30	9.20	0.10	1.90	0.1500		
BFG	5.40	2.60	5.30	9.00	7.50	8.40	0.30	9.20	0.10	1.90	0.1500		
Diesel oil	1.81	1.36	2.72	1.36	1.36	4.07	6.79	1.81	1.36	0.50	0.01		
Diesel oil (in gas turbine)	0.002	0.0012	0.17	0.28	0.0023	0.007	0.0023	0.44	0.053				
Fuel	3.98	1.20	5.31	2.55	255	4.56	2.06	87.80	0.34	2.50	0.02		
Coal	PS	PS	PS	PS	PS	PS	PS	PS	PS	10	0.07	6.70	3.30
Wood	9.46	1.76	21.10	9.03	14.20	20.60	1.20	181	1.51	50	1.22	5.00	3500

Table 3-3 shows the emission factors used for the 1A1a Public electricity and Heat Production in Brussels Capital region.

Table 3-3 Emission factors for the sector 1A1a Public electricity and Heat Production in the Brussels region.

Fuel	UNIT	NO <sub>x</sub>	NM VOC	SO <sub>x</sub>	NH <sub>3</sub>	PM <sub>2.5</sub>	PM <sub>10</sub>	TSP	BC (EC)	CO	PCDD/PCDF*
Natural gas	g/GJ	89	2,6	0,28	0,6	0,89	0,89	0,89	0,022	39	0,5
Gas oil	g/GJ	398	0.19	46,5	0.1	1,95	1,95	1,95	0,65	1,49	0,5
Waste	g/tonne	271,77	20	45,78	9	3,65	3,65	3,65	0,12	51,74	19,88
Fuel	UNIT	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Total HAP
Natural gas	mg/GJ	0,002	0,00025	0,1	0,12	0,0008	7,6E-05	0,0005	0,0112	0,002	0,00308
Gas oil	mg/GJ	0,007	0,0012	0,05	0,002	0,28	0,17	0,0023	0,0023	0,44	
Waste	mg/tonne	1109	100	66,1	79,1	91,54	245,5	14,2	1,2	181	12,16

\* ng-TEQ/GJ or ng-TEQ/tonne

The Flemish emission factors to calculate NO<sub>x</sub> emissions are given below in Table 3-4. The emissions are calculated based on electricity production only (final net).

Table 3-4 Emission factors of NO<sub>x</sub> for the sector 1A1a Public electricity and Heat Production in the Flemish region

Installation	Unit	Emission factor	Source
CHP (power plant) - existing	g/MWhe	280	MBO*
Gas motor - existing	g/MWhe	1.400	MBO
Diesel motor - existing	g/MWhe	5.000	MBO
CHP (power plant) - new	g/MWhe	100	MBO
Gas motor - new	g/MWhe	1.000	MBO
Diesel motor - new	g/MWhe	1.500	MBO
Turbojets	g/MWhe	3000	Emission measurements
Open cycle gasturbines	g/MWhe	1500	Emission measurements
Steam power plant Alost	kg/kNm3	1.6	US-EPA

\*MBO: Environmental Policy Agreement

The calculation of SO<sub>2</sub> emissions originating from installations not equipped with continuous measurements is not applicable: it concerns gas turbines, CHP, gas motors (all burnt on gas) or turbojets (use of fuel with very low sulphur content).

The emission factors used to calculate the emissions of NMVOC are adjusted with rest factors for FGD (flue gas desulphurisation) and SCR (selective catalytic reduction) (Table 3-5). A distinction is made between normal boilers and gas turbines (GT).

Emissions calculation:

$$Em(kg) = \frac{M (GJ) \times EF \left( \frac{g}{GJ} \right)}{1000} \times RF_{FGD} \times RF_{SCR}$$

Table 3-5 Emission factors of NMVOC for the sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel	Unit	Emission factor NMVOC - uncontrolled	Emission factor HCB-uncontrolled	RF-FGD	RF-SCR	Source
Coal	g/GJ	0,4	0.0000062	1	0,3	Eurelectric
Water treatment sludge	g/GJ	10	0.000006	1	0,3	VMM + Eurelectric for reduction
Olive stones	g/GJ	10	0.000006	1	0,3	VMM + Eurelectric for reduction
Wood dust	g/GJ	10	0.000006	1	0,3	VMM + Eurelectric for reduction
Wood chips	g/GJ	10	0.000006	1	0,3	VMM + Eurelectric for reduction
Wood pellets	g/GJ	10	0.000006	1	0,3	VMM + Eurelectric for reduction
Biodust	g/GJ	10	0.000006	1	0,3	VMM + Eurelectric for reduction
Fuel A	g/GJ	0,6		1	0,3	Eurelectric
Gas oil	g/GJ	7,5		1	1	VMM
Gasoil – gas turbine	g/GJ	1,5		1	1	Eurelectric
Paraffin	g/GJ	3		1	1	CITEPA
Natural gas	g/GJ	1		1	0,3	VMM + Eurelectric for reduction
Natural gas – gas turbine	g/GJ	0,5		1	1	Eurelectric
Blast-furnace gas	g/GJ	0		1	1	VMM

An emission factor of 8 mg CO/Nm<sup>3</sup> flue gas is applied for gas-fired installations not equipped with continuous measurement devices (based on continuous measurements of other similar installations).

Although the TSP emissions originating from installations not equipped with continuous measurement devices are very low per unit fuel (installation groups fed with natural gas, blast-furnace gas, gas oil and paraffin or lamp oil), the high volumes of fuel burnt cause a significant emission. The emission factors used to calculate the emissions of TSP are adjusted with rest factors for ESP (electrostatic precipitation for thermal power plants), FGD and SCR (Table 3-6).

Emissions calculation:

$$Em(ton) = \frac{M(GJ) \times EF\left(\frac{g}{GJ}\right)}{1.000.000} \times RF_{ESP} \times RF_{FGD} \times RF_{SCR}$$

Table 3-6 Emission factors of TSP for the sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel	Unit	Emission factor	RF <sub>ESP</sub>	RF <sub>FGD</sub>	RF <sub>SCR</sub>	Source
Gas oil	g/GJ	3	0,01	0,1	1	CORINAIR
Paraffin	g/GJ	3	0,01	0,1	1	CORINAIR

Fuel	Unit	Emission factor	RF <sub>ESP</sub>	RF <sub>FGD</sub>	RF <sub>SCR</sub>	Source
Natural gas	g/GJ	0,005	0,01	0,1	1	CORINAIR
Blast-furnace gas	g/GJ	0,1	0,01	0,1	1	CORINAIR

Emissions of PM10 and PM2,5 are calculated as a fraction of TSP:

$$Em_{PM10} (ton) = Em_{TSP} (ton) \times \frac{\%PM10}{100}$$

$$Em_{PM2,5} (ton) = Em_{TSP} (ton) \times \frac{\%PM2,5}{100}$$

The percentages applied per power plant are given below (Table 3-7):

Table 3-7 Percentages of PM10 and PM2,5 as a fraction of TSP per power plant and percentages of EC as a fraction of PM2,5 per power plant

Power Group	%PM10	%PM2,5	Source	%EC Source: TNO
EDF Luminus Gent	80	70	VITO	19.45
EDF Luminus Harelbeke	80	70	VITO	*
E.on Langerlo	100	100	CORINAIR	1.67
Kallo 1	100	100	CORINAIR	*
Kallo 2	100	100	CORINAIR	*
Mol 12	64,4	32,7	LBE-2001	*
Rodenhuize 2	100	100	CORINAIR	*
Rodenhuize 3	100	100	CORINAIR	*
Rodenhuize 4	78,2	32,3	LBE-2001	9.76
Ruien 3	67	29	EPA	1.68
Ruien 4	67	29	EPA	1.5
Ruien 5	46	33	LBE-2008	3.52
Ruien 6	100	100	CORINAIR	7
Ruien after deSO <sub>x</sub>	71	51	EPA	Equally Ruien 3, 4 and 5

Turbojets	100	100	CORINAIR	7
Gas groups	100	100	CORINAIR	7
A&S Energie	71.43	34.30	EMEP/EEA Guidebook	10
Electrawinds Biomassa	80	70	VITO	7.02

\*: Power plant not active in 2013

Heavy metals can come from various fuels. Depending on the fuel and the type of installation different techniques will be used and/or be combined.

In case one features analyses of the flue gases, these will be used at first to determine the emissions of heavy metals (this will particularly be the case at sites with flue gas desulphurisation (FGD)). In case no such measurements are available, or for certain heavy metals the emission was not determined by the flue gas analysis, one can use the following techniques:

- gaseous fuels (natural gas and blast furnace gas): use of emission factors
- liquid fuels (heavy fuel, gas oil and lamp oil): use of emission factors
- solid fuels (coal, biofuels): method based on the emission rates determined by Laborelec and elementary analyses of the solid fuels.

In certain cases, one shall combine techniques when:

- the flue gas analysis does not cover all the necessary parameters: combination of the flue gas analyses with 1 or more other techniques (emission factors/calculation on the basis of the analyses on the solid fuel). The missing parameters will be completely replaced by the alternative calculation.
- another emission point (chimney) is used for the same group, but no flue gas analyses are available: use of 1 or more other techniques for the whole calculation of the emissions through the other chimney, taking into account the utilization rate of the chimneys (split factor).

## Heavy metals from solid fuels

### Calculation based on flue gas analyses

Where analyses of the flue gases (min. 1 per year) are available for the installation, these measurements are used to determine the annual emissions of heavy metals. Emissions calculation:

$$Em_{ZM1} \left( \frac{kg}{y} \right) = \frac{AW_{ZM1} \left( \frac{mg}{Nm^3} \right) \times V_{RG} \left( \frac{Nm^3}{y} \right)}{1.000.000}$$

with :

- $Em_{ZM1}$  : annual emission of the heavy metal considered
- $AW_{ZM1}$  : average analysis value of the heavy metal in the dry flue gases at a specific oxygen content (e.g. 6% O<sub>2</sub>)
- $V_{RG}$  : Volume of the flue gases on yearly basis

## Calculation from fuel analyses

If no analyses of the flue gases are available or parameters are missing in the existing flue gas analyses, the emissions of the heavy metals are calculated using the fuel analyses.

Calculation of the emission per heavy metal per amount of fuel:

$$Em_{ZM1}(kg) = \left[ \frac{M(ton) \times 1000 \times \frac{As(\%)}{As_{std}(\%)} \times \frac{PM_{inst-av}(\frac{mg}{Nm^3})}{PM_{std}(\frac{mg}{Nm^3})} \times AW_{ZM1}(\frac{mg}{kg}) \times \frac{EP_{solid-ZM1}}{100}}{1.000.000} + \frac{M(ton) \times 1000 \times AW_{ZM1}(\frac{mg}{kg}) \times \frac{EP_{gas-ZM1}}{100}}{1.000.000} \right] \times \frac{RF_{FGD}}{100}$$

with :

- M : the amount of dry fuel expressed in tons. This may be the total annual quantity, or the quantity per batch delivered. The data comes from Michelangelo and is provided by TDM. However, the raw data is the wet quantity, so that it has to be converted first to the dry quantity by means of the moisture content.

$$M_{dry}(ton) = M_{wet}(ton) \times \frac{100 - moisture\ content}{100}$$

- As : the ash content of the fuel, either coming from Michelangelo and provided by TDM, either submitted by external analyses, provided by Fuel Procurement.
- As<sub>std</sub> : standard ash content that was used in the study of Laborelec to determine the emission rates. this default percentage should be calculated to the current ash-percentage. It amounts to 18.5%.
- PM<sub>inst-av</sub> : the (weighted) yearly average dust emission for the set of groups for which the calculation is performed, expressed in mg/Nm<sup>3</sup> at 0% O<sub>2</sub> and dry flue gases. The data is available in Image and is provided by TDM.
- PM<sub>std</sub> : standard dust emission that was used in the study of Laborelec to determine the emission rates. The default percentage should be calculated to the current dust emission (100 mg/Nm<sup>3</sup> at 0% O<sub>2</sub>)
- AW : the analysis value of the heavy metal in the solid fuel. This information is taken from the external analysis reports provided by Fuel Procurement
- EP<sub>solid</sub> : the emission rate for a particular heavy metal in terms of emissions in ash-bound state (Table 3-8). The bulk of the heavy metals emitted is adsorbed on the fly ashes.
- EP<sub>gas</sub> : the emission rate for a particular heavy metal in terms of emissions in the volatile state (Table 3-8). Only a few heavy metals are emitted in volatile state.
- RF<sub>FGD</sub> : the rest factor as a result of the presence of a FGD installation (flue gas desulphurisation)(Table 3-8). For the heavy metals this factor is put at 100% because the effect of the FGD is already taken into account by reduced dust emissions. Only heavy metals that are emitted in volatile state are even further reduced by the FGD.

Table 3-8 Factors to calculate emissions of heavy metals for the sector 1A1a Public electricity and Heat Production in the Flemish region based on fuel analyses

Parameter	EP <sub>solid</sub> (%)	EP <sub>gas</sub> (%)	RF <sub>FGD</sub> (%)
As	2,42	0	100
Cd	2,56	0	100
Cr	0,84	0	100
Cu	1,03	0	100
Ni	1,1	0	100
Pb	1,54	0	100
Se	1,69	20,2	55
Zn	1,96	0	100
Hg	-	100	15

Conversion of delivered to consumed solid fuels:

The quantities of solid fuels available in Michelangelo are delivered quantities. To know the exact emissions, these values must be converted into the amount of fuel consumed

$$Em_{ZM1-used} (kg) = Em_{ZM1} (kg) \times \frac{M_{total} (ton) - \Delta stock (ton)}{M_{total} (ton)}$$

with :

- $M_{total}$  : the total annual amount of delivered fuels
- $\Delta stock$  : the stock difference between the end of the year and the beginning:

$$\Delta stock = stock_{31/12/yyyy} - stock_{1/1/yyyy}$$

**Heavy metals from fluid fuels (fuel A and gas oil)**

The emission factors shown are intended for installations without any form of dust reduction measures or NO<sub>x</sub> or SO<sub>2</sub> reduction measures. So a rest factor must be used according to the availability of certain installations (Table 3-9):

- ESP : dust reduction via electrostatic dust filter (or sleeve filter)
- FGD : SO<sub>2</sub>-reduction via FGD
- SCR : NO<sub>x</sub>-reduction via Selective Catalytic Reduction (SCR)

Table 3-9 Emission factors for heavy metals from fluid fuels for the sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel		Emission factor - uncontrolled (g/ton)	RF <sub>ESP</sub>	RF <sub>FGD</sub>	RF <sub>SCR</sub>	source
Fuel A <sup>2</sup>	As	0,081	0,01	0,1	1	EPA (CORINAIR for reductions)
Fuel A	Cd	0,051	0,1	0,1	1	EPA (CORINAIR for reductions)
Fuel A	Cr	0,139	0,01	0,1	1	EPA (CORINAIR for reductions)
Fuel A	Cu	0,223	0,01	0,1	1	EPA (CORINAIR for reductions)
Fuel A	Hg	0,014	0,965	0,05	1	EPA (CORINAIR for reductions)
Fuel A	Ni	10,723	0,01	0,1	1	US-EPA + CORINAIR for reductions
Fuel A	Pb	0,192	0,1	0,1	1	US-EPA + CORINAIR for reductions
Fuel A	Se	0,087	0,235	0,24	1	US-EPA + CORINAIR for reductions
Fuel A	Zn	3,693	0,1	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	As	0,074	0,01	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	Cd	0,0555	0,1	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	Cr	0,0555	0,01	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	Cu	0,111	0,01	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	Hg	0,0555	0,965	0,05	1	US-EPA + CORINAIR for reductions
Gas oil	Ni	0,0555	0,01	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	Pb	0,1665	0,1	0,1	1	US-EPA + CORINAIR for reductions
Gas oil	Se	0,2775	0,235	0,24	1	US-EPA + CORINAIR for reductions
Gas oil	Zn	0,074	0,1	0,1	1	US-EPA + CORINAIR for reductions

Emissions calculation :

$$Em_{ZM1}(kg) = \frac{M(ton) \times EF\left(\frac{g}{ton}\right)}{1000} \times RF_{ESP} \times RF_{FGD} \times RF_{SCR}$$

#### Heavy metals from gaseous fuels (natural gas and blast-furnace gas)

Only mercury and selenium are considered, given their volatility.

<sup>2</sup> Fuel A = heavy fuel. The 'A' is an indication for the S-content (max 1%)

The emission factors shown are intended for installations without any form of dust reduction measures or NO<sub>x</sub> or SO<sub>2</sub> reduction measures. So a rest factor must be used according to the availability of specific installations (Table 3-10):

- ESP : dust reduction via electrostatic dust filter (or sleeve filter)
- FGD : SO<sub>2</sub>-reduction via FGD
- SCR : NO<sub>x</sub>-reduction via Selective Catalytic Reduction (SCR)

Table 3-10 Emission factors for heavy metals from gaseous fuels for the sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel		Emission factor - uncontrolled (g/kNm <sup>3</sup> )	RF <sub>ESP</sub>	RF <sub>FGD</sub>	RF <sub>SCR</sub>	Source
Natural gas	Hg	0,00416	0,965	0,05	1	US-EPA + CORINAIR for reductions
Natural gas	Se	0	0,235	0,24	1	US-EPA + CORINAIR for reductions
Blast-furnace gas	Hg	0,0000625	0,965	0,05	1	NPI + CORINAIR for reductions

Emission calculations:

$$Em_{ZM1}(kg) = \frac{M(kNm^3) \times EF\left(\frac{g}{kNm^3}\right)}{1.000} \times RF_{ESP} \times RF_{FGD} \times RF_{SCR}$$

The calculation of emissions of the PAHs benzo(a)pyrene, benzo(k)fluoranthene, indeno(1,2,3-cd)pyrene and benzo(b)fluoranthene is based on emission factors, which are given in Table 3-11.

Table 3-11 Emission factors for PAH(4) for the sector 1A1a Public electricity and Heat Production in the Flemish region

g/GJ	benzo(a)pyrene	benzo(k)fluoranthene	indeno(1,2,3-cd)pyrene	benzo(b)fluoranthene	source
coal	6,80E-07	0	1,09E-06	0	EPA
gas oil	0	0	0	0	EPA
gas oil gas turbine	0	0	0	0	Econotec
heavy fuel	0	0	6,40E-06	0	EPA
natural gas	2,55E-07	3,80E-07	3,80E-07	3,80E-07	Econotec
blast-furnace gas	0	0	0	0	Econotec
sludge, olive stones, wood dust, pellets, coffee, wood chips, biodust	8,00E-05	0	0	0	CORINAIR
biofuel	0	0	0	0	EPA

The calculation of emissions of dioxins and furans (PCDD/PCDF) is based on emission factors, representing the sum of PCDDs and PCDFs (Table 3-12). The emission is expressed in mg I-TEQ (International toxic equivalent). It is assumed that only FGD affects the PCDD/PCDF-emissions. A distinction should be made between normal boilers and gas turbines.

Table 3-12 Emission factors for PCDD/PCDF for the sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel	Emission factor (mg I-TEQ/TJ)	RF <sub>FGD</sub>	source
Coal	0,000417	0,0124	Analyses by the power plants
Water treatment sludge	0,000417	0,0124	Analyses by the power plants
Olive stones	0,000417	0,0124	Analyses by the power plants
Wood dust	0,00163	0,0124	ESI
Wood chips	0,00163	0,0124	ESI
Wood pellets	0,00163	0,0124	ESI
Biodust	0,00163	0,0124	ESI
Fuel A	0,00124	0,0124	ESI
Gas oil	0,0009	0,0124	ECONOTEC
Gas oil – gas turbine	0,0005	0,0124	ECONOTEC
Paraffin	-	-	ECONOTEC
Natural gas – gas turbine	0	-	-
Natural gas – gas turbine	0	-	-
Blast-furnace gas	0	-	-

Emission calculation:

$$Em (mg I - TEQ) = \frac{M (GJ) \times EF \left( \frac{mg I - TEQ}{TJ} \right)}{1000} \times RF_{FGD}$$

The calculation of the emissions of polychlorinated biphenyls (PCB) is based on emission factors, representing the sum of PCBs (Table 3-13).

It is assumed that neither FGD nor SCR affect the PCB emissions.

Table 3-13 Emission factors for PCBs for the sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel	Emission factor (mg/TJ)	Source
Coal	0,04	ESI
Water treatment sludge	0,0456	ESI
Olive stones	0,0456	ESI
Wood dust	0,0456	ESI
Wood chips	0,0456	ESI
Wood pellets	0,0456	ESI
Biodust	0,0456	ESI

Fuel	Emission factor (mg/TJ)	Source
Fuel A	0,0415	ESI
Gas oil	-	-
Paraffin	-	-
Natural gas	0	-
Blast-furnace gas	-	-

Emission calculation :

$$Em(kg) = \frac{M (GJ) \times EF \left( \frac{mg}{TJ} \right)}{1.000.000.000}$$

The combined heat-power (CHP) installations (in joint venture with the electricity producers) are located in different sectors in Belgium (refineries, industry, agriculture and service sector).

Emissions of CHP installations in the refinery sector are reported in the environmental annual reports submitted by the operator of the refinery.

Emissions of industrial installations are mainly reported in the environmental annual reports submitted by the operator of the plant. The missing emissions are estimated based on the energy data per CHP installation multiplied by an emission factor, as given below in Table 3-14, Table 3-15 and Table 3-16.

Table 3-14 Emission factors of NO<sub>x</sub>, CO, SO<sub>2</sub> and NH<sub>3</sub> for the industrial CHP installations in joint-venture with the power plants in the Flemish region

Installation	Fuel	Unit	NO <sub>x</sub>	CO	SO <sub>2</sub>	NH <sub>3</sub>	Source
Gas turbine	Natural gas	g/GJ	48	4.8	0.281	0.6	EMEP/EEA Guidebook 2013
Gas motor	Natural gas	g/GJ	135	56	0.5	0.6	EMEP/EEA Guidebook 2013
Gas turbine	Gas oil	g/GJ	398	1.49	46.5	0.1	EMEP/EEA Guidebook 2013
Gas motor	Gas oil	g/GJ	942	130	46.5	0.1	EMEP/EEA Guidebook 2013
Gas turbine	Biogas/waste gas	g/GJ	88	13	43.7	15	EMEP/EEA Guidebook 2013
Gas motor	Biogas/waste gas	g/GJ	88	13	43.7	15	EMEP/EEA Guidebook 2013

Table 3-15 Emission factors of TSP and the percentages of PM10 and PM2,5 as a fraction of TSP and the EC fraction of PM2.5 for the industrial CHP installations in joint-venture with the power plants in the Flemish region

Sector	Fuel	Emission factor TSP	Unit e.f.	% PM10	% PM2,5	Fraction EC
Chemical industry	Coal	35,0	ton/PJ	71%	34%	0,1
	heavy fuel	20,0	ton/PJ	75%	45%	0,1
	Natural gas	0,200	ton/PJ	100%	100%	0,07
	Gas oil	5,00	ton/PJ	100%	100%	0,45
	Biogas	0,200	ton/PJ	100%	100%	0,07
	Ind. Waste	0,200	ton/PJ	100%	100%	0,07
Food, drinks and beverages	Natural gas	0,200	ton/PJ	100%	100%	0,07

	biogas	0,200	ton/PJ	100%	100%	0,07
Paper	Natural gas	0,200	ton/PJ	100%	100%	0,07
Textile, leathere and clothing	Natural gas	0,200	ton/PJ	100%	100%	0,07
Gas distribution	Natural gas	0,200	ton/PJ	100%	100%	0,07
	Gas oil	5,000	ton/PJ	100%	100%	0,45

Table 3-16 Emission factors of heavy metals for the industrial CHP installations in joint-venture with the power plants in the Flemish region

CHP installations	Fuel	Unit	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Chemical industry	coal	mg/GJ	8,1	1	1,6	8	5	4,8	5,4	25	19	EMEP/EEA Guidebook 2009
Chemical industry	heavy fuel	mg/GJ	4,9	1,3	0,4	4,3	2,7	5,7	273	2,2	94	EMEP/EEA Guidebook 2009
Chemical industry	natural gas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Chemical industry	gas oil	mg/GJ	4,1	1,4	1,4	1,8	1,4	2,7	1,4	6,8	1,8	EMEP/EEA Guidebook 2009
Chemical industry	biogas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Chemical industry	industrial waste	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Food, drinks and tobacco	natural gas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Food, drinks and tobacco	biogas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Paper industry	natural gas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Textile, leather and clothing industry	natural gas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009
Gas distribution	natural gas	mg/GJ	0,2	0,5	0,2	0,09	0,7	0,4	1	0,01	14	EMEP/EEA Guidebook 2009

Emissions of waste incineration installations with energy recuperation are generally reported in the environmental annual reports submitted by the operator of the installation. In the Flemish region the waste incineration with energy recuperation includes the incineration of industrial and domestic waste.

The PCDD/F emissions of 1990 and 1995 (industrial and domestic waste) are based on the results of a study performed by VITO under the authority of VMM (Polders et al., 2003). Since 2000 the emissions of domestic waste incineration are reported in the yearly environmental reports. Since 2000 the emissions of industrial waste incineration are calculated by using activity data and emission factors. The activity data are the amount of waste obtained from OVAM (Public Waste Agency of Flanders). The emission factors are taken from the UNEP Standardized Toolkit for PCDD/F (Table 3-17).

The HCB emissions are calculated by using activity data and emission factors. The activity data are the amount of waste obtained from OVAM (Public Waste Agency of Flanders). The emission factors are taken from the EMEP/CORINAIR Guidebook for HCB (Table 3-18).

Table 3-17 Emission factors of PCDD/F for the sector 1A1a Incineration of waste in the Flemish region

Fuel	Unit	Value	Reference
Industrial waste	µg TEQ/tonne	0.5	UNEP Standardized Toolkit; Category 1a4: Waste incineration; Municipal solid waste incineration; High tech. combustion, sophisticated APCS
Hazardous waste	µg TEQ/tonne	0,75	UNEP Standardized Toolkit; Category 1b4: Waste incineration; Hazardous waste incineration; High tech. combustion, sophisticated APCS
Clinical waste	µg TEQ/tonne	1	UNEP Standardized Toolkit; Category 1c4: Waste incineration; Medical/hospital waste incineration; High tech, continuous, sophisticated APCS
Sewage sludge	µg TEQ/tonne	0,4	UNEP Standardized Toolkit; Category 1e3: Waste incineration; Sewage sludge incineration; State-of-the-art, full APCS

Table 3-18 Emission factors of HCB for the sector 1A1a Incineration of waste in the Flemish region

Fuel	Unit	Value	Reference
Industrial waste	g/tonne	0.0001	EMEP/CORINAIR Guidebook (2005)
Hazardous waste	g/tonne	0.01	EMEP/CORINAIR Guidebook (2005)
Clinical waste	g/tonne	0.019	EMEP/CORINAIR Guidebook (2005)
Sewage sludge	g/tonne	0.002	EMEP/CORINAIR Guidebook (2009)
Domestic waste	g/tonne	0.001	EMEP/CORINAIR Guidebook (2005)

Emissions of CHP installations in the service and agricultural sector are calculated based on the energy data (energy balance) and emission factors, as given below in Table 3-19 and Table 3-20.

Table 3-19 Emission factors for the sector CHP installations in the service sector, reported in sector 1A1a Public electricity and Heat Production in the Flemish region

Fuel	Unit	SO <sub>2</sub>	NO <sub>x</sub>	COV	CO	NH <sub>3</sub>	TSP/PM10/PM2,5	EC
renewable fuels	g/GJ	0.5	165	50	56	-	2	0.14
diesel oil	g/GJ	48	307	46	130	-	30	3
natural gas	g/GJ	0.5	158	47	56	-	2	0.14
Source		EMEP/EEA or S-content (diesel oil)	Emission limit values	Emission limit values	EMEP/EEA		EMEP/EEA	TNO

Fuel	Unit	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Natural gas	g/GJ	0.00004	0.000003	0.0001	0.00005	0,00005	0.00001	0.00005	0.0002	0.0029	EMEP/EEA
Diesel oil	g/GJ	0.00015	0.00001	0.00011	0.00006	0.00020	0.0003	0.00001	0.00022	0.058	EMEP/EEA
Biogas	g/GJ	0.00004	0.000003	0.0001	0.00005	0.00005	0.00001	0.00005	0.0002	0.0029	EMEP/EEA

Fuel	benzo(a)pyrene (kg/TJ)	benzo(b)fluoranthene (kg/TJ)	benzo(k)fluoranthene (kg/TJ)	indeno(1,2,3-cd)pyrene (kg/TJ)	PCDD/F (mg TEQ/TJ)
Natural gas					0,0005
diesel oil	0,000101	5,38E-05	8,19E-05	0,000201	0,0005
biogas/waste gas					0,008
Source	TNO	TNO	TNO	TNO	UNEP

Table 3-20 Emission factors for the CHP installations in the agricultural sector, reported in sector 1A1a Public electricity and Heat Production in the Flemish region

CHP agriculture	Unit	natural gas	other (biogas)	diesel oil
CO	kg/TJ	407	407	385
NO <sub>x</sub>	kg/TJ	158	165	307
C <sub>x</sub> H <sub>y</sub>	kg/TJ	47	50	46
dust	kg/TJ	0,2	0,2	5
PM10	kg/TJ	0,2	0,2	5
SO <sub>x</sub>	kg/TJ	0	0	48
Pb	kg/TJ	0,000234	0,000234	0,004
NH <sub>3</sub>	kg/TJ	/	/	/
As	kg/TJ	0,000	0,000	0,002
Cd	kg/TJ	0,001	0,001	0,001
Cr	kg/TJ	0,001	0,001	0,001
Cu	kg/TJ	0,000	0,000	0,003
Hg	kg/TJ	0,000	0,000	0,001
Ni	kg/TJ	0,001	0,001	0,001
Se	kg/TJ	0,000	0,000	0,007
Zn	kg/TJ	0,014	0,014	0,002
PM <sub>2,5</sub>	kg/TJ	0,200	0,200	5,000

### **Petroleum refining (category 1A1b)**

Petroleum refining activities (5 refineries) only take place in Flanders.

Category 1A1b is a key category of SO<sub>x</sub> emissions in terms of emissions level and trend and a key category of Ni emissions in terms of emissions level.

The activity data of the petroleum refining are taken from the Flemish energy balance as petroleum refining only occurs in Flanders.

Combustion emissions of SO<sub>2</sub>, NO<sub>x</sub>, CO, NMVOC, TSP and heavy metals are directly reported by the refineries in the environmental annual reports submitted by the operator of the plant.

SO<sub>2</sub> emissions are calculated based on the sulphur content of the fuel or on measurements. NO<sub>x</sub>, CO, NMVOC, TSP and heavy metals emissions are calculated with emission factors provided by the refineries or emissions are calculated based on measurements.

Emissions of PM10 and PM2,5 are calculated as a fraction of TSP (Schrooten & Van Rompaey, 2002) (Table 3-21):

Table 3-21 Percentages of PM10 and PM2,5 as a fraction of TSP for petroleum refineries

Fuel	%PM10	%PM2,5	Source
Refinery gas	100	100	VMM
Light fuel	100	100	VMM
Heavy fuel	80	70	VMM
Cokes	60	35	VMM

### **Manufacture of solid fuels and other energy industries (category 1A1c)**

The emissions originating from category 1A1c 'Manufacture of Solid Fuels and Other Energy Industries' are the emissions coming from the combustion in the coke ovens. Nowadays 2 plants are still operational in Belgium instead of 8 plants in the beginning of the nineties. One plant was closed in the Flemish region in 1996, 4 plants closed in the Walloon region (one in 1995, a second in 2000, a third in 2005 and a fourth in 2010) and the only plant active in the Brussels region was closed in 1994.

In Wallonia, the emission factors for all pollutants are plant specific. The non-diffuse NH<sub>3</sub> emissions are estimated by using the default emission factors (coke gas, 0,87 g/GJ).

There was a coke plant in the Brussels region until 1993. The emission factors used are those included in the EMEP/EEA guidebook 2013; the emission factor for NH<sub>3</sub> is the same used in Wallonia.

In Flanders the last coke plant has closed in 1996. But there are still cokes ovens as part of the iron and steel sector. The emission factors for SO<sub>x</sub> are based on the sulphur content of the fuel. The emission factors for NO<sub>x</sub> and CO are based on literature data.

The emission factors for heavy metals are given in Table 3-22.

Table 3-22 Emission factors of heavy metals for the cokes ovens in Flanders in the sector 1A1c

	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
g/Mg produced coke	1,2	0,03	0,02	0,1			0,3			EMEP/EEA Guidebook 2008

Emissions of coal mining activities were reported until the beginning of the nineties. The emission factors for SO<sub>x</sub> are based on the sulphur content of the fuel. CO and NO<sub>x</sub> emissions are calculated with emission factors provided by the facilities. The mining industries have disappeared with the closure of the last coalmines in 1992.

Also some emissions due to gas transport activities are included in this sector. The emission data are provided by the facilities.

## **3.3. Manufacturing Industries and Construction (1A2)**

### **3.3.1. Source category description (1A2)**

The structure of the industrial sector has undergone profound changes over recent decades. The metallurgy and textile sectors have been relatively stable, after several waves of closures and restructuring. The metallurgical industry nevertheless remains one of the key sectors of Belgian industry, both in terms of employment and turnover. The two other key sectors of industrial activity are

the chemical industry and the food processing industry. These three sectors each contribute about 10% of gross value added of the industrial sector.

The category 1A2 'Manufacturing industries and construction' contains the energetic emissions of the industrial sector of the 3 regions in Belgium. The following sectors are involved: iron and steel (1A2a), non-ferrous metals (1A2b), chemicals (1A2c), pulp, paper and print (1A2d), food processing, beverages and tobacco (1A2e), non-metallic minerals (1A2f) and other industries (1A2g).

The following industries are integrated in category 1A2g (Other industries): metal products, textile, leather and clothing and other industry (wood industry, rubber and synthetic material, manufacturing of furniture, recycling and construction).

The industrial sector is not very developed in the Brussels region, mainly due to its urban features. The only big industry is a car manufacturer. The other industries are (very) small companies specialised in high added value products and/or located close to the final consumer. All these industries are classified in the 1A2g category (Other industries).

The emissions originating from the use of recovered fuels from cracking units or other processes where a fuel is used as a raw material and where a part of this fuel (or transformed product) is recovered for energy purposes is allocated to category 1A2c (other fuels).

Emissions of industrial combined heat-power installations in joint venture with the energy sector are allocated in the category 1A1a.

Emissions of the combustion of blast furnace gas, produced in the steel plants and delivered to the energy sector, are allocated in the category 1A1a. The following chart (fig. 3.2) shows the trend of the energy consumption in this sector :




Figure 3.2 Trend of fuel consumption in the manufacturing Industries and construction.

### 3.3.2. Methodological issues

#### Default emission factors

Pollutant emissions are mostly reported directly by the individual large companies on the basis of analyses. For most sectors the remainder of the emissions is calculated on the basis of the remaining fuel consumption (estimated as the difference between energy consumption reported in the regional

energy statistics for the whole sector and the fraction reported by the large companies) and standard emission factors listed in tables below.

The energy consumption data originate from the regional energy balances in the 3 regions, supplemented with specific information from the companies themselves, for example activity data from iron and steel industry.

Generally in the combustion processes, the SO<sub>2</sub> emissions are mainly based on the sulphur content of the fuel and the NO<sub>x</sub> emissions vary with the fuel and the sector.

The following tables (Table 3-23 and Table 3-24) give the default emission factors used in the Walloon and Brussels region. Estimated emissions in individual plants in Flanders are based on plant-specific emission factors per installation.

Table 3-23 Emission factors for the sector 1A2 Manufacturing Industries and Construction in the Walloon region (EMEP/EEA Guidebook 2013 – NO<sub>x</sub> diesel : calculated with the maximum norm in the plants permits for diesel boilers - NH<sub>3</sub> : EPA and Emep 1996 – HM for GC and BFG : average of measurements between 2005 and 2011 on boilers)

	SO <sub>2</sub>	NO <sub>x</sub>	COVNM	CO	TSP	PM <sub>10</sub>	PM <sub>2,5</sub>	BC	NH <sub>3</sub>
	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ
Natural gas	0.5	74	2	29	0.45	0.45	0.45	0.02	0.6
Biogas	43.7	88	2.5	13					15
Diesel oil	47	163	25	66	20	20	20	11.2	0.1
Fuel		163	25	66	20	20	20	11.2	0.1
Coal	900	173	88	931	124	117	108	6.912	0.4
Coke	540	173	88	931	124	117	108	6.912	0.4
Wood	11	91	300	570	150	143	140	39.2	37
BFG	70	74	2.5	25	2.60	1.18	1.18	0.03	
Coke gas	70	74	2.5	25	2.60	1.18	1.18	0.03	0.87
LPG		74	2	29	0.45	0.45	0.45	0.02	0.6
Petroleum coke	540	173	88	931	124	117	108	6.912	

	As	Cd	Cu	Cr	Ni	Pb	Sé	Zn	Hg	Dioxins	PAH	PCB	HCB
	mg/GJ									ng/GJ	mg/GJ	yg/GJ	yg/GJ
Natural gas	0.1	0	8E-05	8E-04	0	0	0.011	0.002	0.1	0.5	0.003		
Diesel oil	1.8	1.36	2.72	1.36	1.36	4.1	6.79	1.81	1.36	1.4	20.1		
Fuel	4	1.2	5.31	2.55	255	4.6	2.06	87.8	0.34	1.4	20.1		
Coal	4	1.8	17.5	13.5	13	134	1.8	200	7.9	203	146.6	170	0.62
Coke	4	1.8	17.5	13.5	13	134	1.8	200	7.9	203	146.6	170	0.62
Wood	0.2	13	6	23	2	27	0.5	512	0.56	100	35	0.06	5
BFG	5.40	2.60	5.30	9.00	7.50	8.40	0.30	9.20	0.10	1.90	0.1500		
Coke gas	5.40	2.60	5.30	9.00	7.50	8.40	0.30	9.20	0.10	1.90	0.1500		
LPG	0.1	0	8E-05	8E-04	0	0	0.011	0.002	0.1	0.5	0.003		
Petroleum coke	4	1.2	12	15	1030	4.6		49	0.11				

SO <sub>x</sub>		S content	EF
		%	g/GJ
FUEL	<1993	3	1400
	1994	2.17	1085
	1995	1.05	520
	1996	1.0	495
gasoil	<1993	0.5	160
	1994		
	1995	0.2	95
	1996	0.2	95
	2008	0.1	48

Table 3-24 Emission factors for the sector 1A2 Manufacturing Industries and Construction in the Brussels region.

Fuel	UNIT	NO <sub>x</sub>	NM VOC	SO <sub>x</sub>	NH <sub>3</sub>	PM <sub>2.5</sub>	PM <sub>10</sub>	TSP	BC (EC)	CO	PCDD/PCDF*
Natural gas	g/GJ	74	23	0,67	0,6	0,78	0,78	0,78	0,031	29	0,52

Gas oil	g/GJ	513	25	47	0,1	20	20	20	11,2	66	1,4
Butane/Propane	g/GJ	74	23	0,67	0,6	0,78	0,78	0,78	0,031	29	0,52
<b>Fuel</b>	<b>UNIT</b>	<b>Pb</b>	<b>Cd</b>	<b>Hg</b>	<b>As</b>	<b>Cr</b>	<b>Cu</b>	<b>Ni</b>	<b>Se</b>	<b>Zn</b>	<b>Total HAP</b>
Natural gas	mg/GJ	0,011	0,0009	0,54	0,1	0,013	0,0026	0,013	0,058	0,73	31,9
Gas oil	mg/GJ	0,08	0,006	0,12	0,03	0,2	0,22	0,008	0,11	29	20,1
Butane/Propane	mg/GJ	0,011	0,0009	0,54	0,1	0,013	0,0026	0,013	0,058	0,73	31,9

\* ng-TEQ/GJ

### **Iron and steel sector (category 1A2a)**

Category 1A2a is a key category of Cr and Zn emissions in terms of emissions level and trend and a key category of NO<sub>x</sub>, CO, Pb and Hg emissions in terms of emissions level or trend.

In the Flemish region there is one integrated steel plant, one plant that produces stainless steel and one that handles molybdenum to be used in the production of stainless steel. In the Walloon region, there are 5 electric arc furnace plants. No iron and steel activities take place in the Brussels region.

Because different approaches approved by the different companies involved (a.o. based on historical background) it is not possible to harmonize completely these methodologies between the 2 regions involved (Flanders and Wallonia).

The emissions from the iron and steel sector are partly put in category 1A2a (energetic part / except for the emissions from the cokes ovens which are allocated in the category 1A1c in Wallonia) and partly in category 2C1 (process part).

In the Walloon region, the last integrated iron and steel plant (blast furnace-oxygen furnace) was closed in 2011. Five electric arc furnaces are operational in 2013.

Since 2004, all the IPPC companies are obliged to report their energy consumptions, their productions and their emissions of IPPC pollutants on a website (Regine). IPPC companies which are also emission trading companies are obliged to report on the same way. This plant information is compared and combined with the energy balance of the sector. The remainder of the emissions is calculated on the basis of the remaining fuel consumption (energy balance of the sector minus plant energy consumptions) and by using the default emission factors of the sector 1A2.

In Flanders, emissions are reported directly by the individual companies. SO<sub>x</sub> emissions are calculated based on the sulphur content of the fuel or on (continuous) measurements. NO<sub>x</sub>, CO and NMVOC emissions are generally calculated based on measurements.

TSP emissions are based on calculations (fuel consumed x emission factors per fuel type). Emissions of PM<sub>10</sub> and PM<sub>2,5</sub> are calculated as a fraction of TSP (Table 3-25).

Table 3-25 Emission factors of TSP and fractions of PM<sub>10</sub> and PM<sub>2,5</sub> and EC fraction of PM<sub>2,5</sub> for the sector 1A2a Iron and steel

Iron and steel		Emission factor	% PM <sub>10</sub>	% PM <sub>2,5</sub>	EC fraction	Source
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%	EMEP/EEA Guidebook 2013
Gas- en diesel oil	ton/PJ	6,47	50%	12,5%	33,50%	EMEP/EEA Guidebook 2013

LPG	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Natural gas	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Cokes gas	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Renewables - solid	ton/PJ	150,00	95%	93%	28,00%	EMEP/EEA Guidebook 2013

Also heavy metals emissions are based on calculations. The emission factors to calculate the heavy metals emissions for the iron and steel sector are given in Table 3-26.

Table 3-26 Emission factors of heavy metals for the sector 1.A.2.a Iron and steel production

Iron and steel		Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Heavy fuel	mg/GJ	4,56	1,2	0,341	3,98	2,55	5,31	255	2,06	87,8	EMEP/EEA Guidebook 2013
Gas-en diesel oil	mg/GJ	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
LPG	mg/GJ	0,0015	0,00025	0,1	0,12	0,00076	0,000076	0,00051	0,011	0,0015	EMEP/EEA Guidebook 2013
Natural gas	mg/GJ	0,0015	0,00025	0,1	0,12	0,00076	0,000076	0,00051	0,011	0,0015	EMEP/EEA Guidebook 2013
Cokes oven gas	mg/GJ	0,0015	0,00025	0,1	0,12	0,00076	0,000076	0,00051	0,011	0,0015	EMEP/EEA Guidebook 2013
renewables - solid	mg/GJ	27	13	0,56	0,19	23	6	2	0,5	512	EMEP/EEA Guidebook 2013

To calculate the remainder of the emissions (emissions not reported directly by the individual companies) from the iron and steel sector in Flanders a new methodology is used (Sleeuwaert et al., 2010). For this methodology (used for the first time during this submission) 3 types of activity data are important: the total energy consumption reported in the regional energy statistics for the iron and steel sector (for each fuel type), the energy consumption reported by the individual companies in this sector (for each fuel type), the pollutants reported by each individual company in the sector. This methodology calculates in the first place, for each fuel type, the difference between the energy consumption reported in the regional energy statistics for the iron and steel sector and the energy consumption reported by the individual companies in this sector. Furthermore this difference is calculated for each pollutant separately on the level of the company. This results for each pollutant and each fuel type, in a percentage of the total energy consumption from which emissions have to be estimated. In combination with a region specific corresponding emission factor (see Table 3-27) the estimated emission is calculated.

Table 3-27 Emission factors of CO, SO<sub>x</sub> and NO<sub>x</sub> in the iron and steel sector used in the collective approach

Iron and steel	Unit	CO (g/GJ)	SO <sub>x</sub> (g/GJ)	NO <sub>x</sub> (g/GJ)
Coal	g/GJ	82	683	242
Cokes	g/GJ	82	683	242
LPG	g/GJ	62	0.0000435	90
Gas and diesel oil	g/GJ	67	47	166
Heavy fuel	g/GJ	67	493	180
Natural gas	g/GJ	59	0.0000450	46
Cokes gas	g/GJ	40	0.5	58

### **Category 1A2b to 1A2e**

Category 1A2b is a key category of Ni and Cu emissions in terms of emissions trend or level.

Category 1A2c is a key category of Hg, As and Ni emissions in terms of emissions level and trend and a key category of SO<sub>2</sub>, Cd and Se emissions in terms of emissions trend.

Category 1A2d is a key category of PM<sub>2,5</sub>, PM<sub>10</sub>, Zn and Cd emissions in terms of emissions level and trend and a key category of TSP emissions in terms of emissions level.

Category 1A2e is a key category of SO<sub>2</sub> and Ni emissions in terms of emissions trend.

In Flanders, emissions of the main pollutants are reported directly by the individual companies. SO<sub>x</sub> emissions are calculated based on the sulphur content of the fuel or on measurements. NO<sub>x</sub>, CO and NMVOC emissions are measured, calculated or estimated based on plant specific information. To calculate the remainder of the emissions (emissions not reported directly by the individual companies) from the categories 1A2b – 1A2e in Flanders a new methodology is used (Sleeuwaert et al., 2010). For a description of this methodology see above in chapter 1A2a, iron and steel. For this collective approach, for each sector in these categories and each fuel type a specific corresponding emission factor is used.

TSP emissions are based on calculations (fuel consumed x emission factors per fuel type). Activity data are taken from the Flemish energy balance. Emissions of PM<sub>10</sub> and PM<sub>2,5</sub> are calculated as a fraction of TSP. The EC emissions are calculated as a fraction of PM<sub>2,5</sub>.

The emission factors used to calculate the TSP emissions and the fractions of PM<sub>10</sub> and PM<sub>2,5</sub> and the EC fraction of PM<sub>2,5</sub> of the key categories is given below in Table 3-28:

Table 3-28 Emission factors used to calculate the TSP emissions and the fractions of PM10 and PM2,5 and EC fraction of PM2.5 for combustion in the sectors of non-ferro, chemistry, pulp and paper and food and drinks.

Non-ferro 1A2b	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction
Cokes	ton/PJ	124	94%	87%	6,40%
Coal	ton/PJ	124	94%	87%	6,40%
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%
Petrol cokes	ton/PJ	20	75%	45%	5,60%
Gas-and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%
LPG	ton/PJ	0,78	100%	100%	4,00%
Natural gas	ton/PJ	0,78	100%	100%	4,00%
Other fuels	ton/PJ	3,625	75%	56%	18,75%
Renewables - solid	ton/PJ	150	95%	93%	28,00%

Chemical sector 1A2c	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction
Petroleum cokes	ton/PJ	20	75%	45%	5,60%
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%
Gas and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%
LPG	ton/PJ	0,78	100%	100%	4,00%
Natural gas	ton/PJ	0,78	100%	100%	4,00%
Other fuels	ton/PJ	3,63	75%	56%	18,75%
Renewable fuels - solid	ton/PJ	150	95%	93%	28,00%
Renewable fuels - liquid	ton/PJ	6,47	50%	12,5%	33,50%
Renewable fuels - gaseous	ton/PJ	0,78	100%	100%	4,00%

Pulp and paper 1A2d	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction
Coal	ton/PJ	2,14	94%	87%	6,40%
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%
Gas and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%
LPG	ton/PJ	0,78	100%	100%	4,00%

Natural gas	ton/PJ	0,78	100%	100%	4,00%
Other fuels	ton/PJ	0,78	100%	100%	4,00%
Renewable fuels - liquid	ton/PJ	150	95%	93%	28,00%
Renewable fuels - gaseous	ton/PJ	0,78	100%	100%	4,00%
Food, drinks and tobacco 1A2e	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction
Cokes	ton/PJ	124	94%	87%	6,40%
Coal	ton/PJ	124	94%	87%	6,40%
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%
Gas-and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%
Lamp petrol	ton/PJ	5	100%	100%	45,00%
LPG	ton/PJ	0,78	100%	100%	4,00%
Natural gas	ton/PJ	0,78	100%	100%	4,00%
Renewables - solid	ton/PJ	150	95%	93%	28,00%
Renewables - gaseous	ton/PJ	0,780	100%	100%	4,00%

In Wallonia, all the plants which are under the IPPC directive report their annual emissions. The remainder of the emissions is calculated on the basis of the energy balance and the default emissions factors (Table 3-23).

#### **Non-metallic minerals (category 1A2f)**

Category 1A2f is a key category of Se emissions in terms of emissions level and trend and a key category of SO<sub>2</sub>, Cd, As, Cr, Ni and Zn in terms of emissions level or trend.

The sector 1A2f includes combustion emissions of the glass sector, the ceramic sector, the lime production in a chemical plant, in sugar plants and in a paper pulp plant. Also process emissions of the glass sector are included in category 1A2f instead of category 2A3.

All the emissions of the cement plants and the lime plants are in the category 2A and are plant specific.

In Flanders, emissions of the main pollutants are reported directly by the individual companies. SO<sub>x</sub> emissions are calculated based on the sulphur content of the fuel or on measurements. NO<sub>x</sub>, CO and NMVOC emissions are measured, calculated or estimated based on plant specific information. To calculate the remainder of the emissions (emissions not reported directly by the individual companies) from the categories 1A2f in Flanders a new methodology is used (Sleeuwaert et al., 2010). For a description of this methodology see above in chapter 1A2a, iron and steel. For this collective approach, for each sector in these categories and each fuel type a specific corresponding emission factor is used (Table 3-29).

Table 3-29 Emission factors of CO, SO<sub>x</sub> and NO<sub>x</sub> in the Non- metallic minerals used in the collective approach

Non- metallic minerals	Unit	CO (g/GJ)	SO <sub>x</sub> (g/GJ)	NO <sub>x</sub> (g/GJ)
Coal	g/GJ	82	683	242
Cokes	g/GJ	82	683	242
LPG	g/GJ	62	0.0000435	90
Gas and dieseloil	g/GJ	67	47	166
Heavy fuel	g/GJ	67	493	180
Natural gas	g/GJ	59	0.0000450	46

TSP emissions are based on calculations (fuel consumed x emission factors per fuel type). Activity data are taken from the Flemish energy balance. Emissions of PM10 and PM2,5 are calculated as a fraction of TSP and EC emissions are determined as a fraction of PM2.5.

The emission factors used to calculate the TSP emissions, the fractions of PM10 and PM2,5 and the EC fraction of PM2.5 of the key categories is given below in Table 3-30:

Table 3-30 Emission factors used to calculate the TSP emissions, the fractions of PM10 and PM2,5 and the EC fraction of PM2.5 for combustion in the sectors of non-metallic mineral product activities

Non-metallic mineral products 1A2f	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction	Source
Cokes	ton/PJ	124	94%	87%	6,40%	EMEP/EEA Guidebook 2013
Coal	ton/PJ	124	94%	87%	6,40%	EMEP/EEA Guidebook 2013
Petrol cokes	ton/PJ	20	75%	45%	5,60%	EMEP/EEA Guidebook 2013
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%	EMEP/EEA Guidebook 2013
Gas and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%	EMEP/EEA Guidebook 2013
LPG	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Petrol	ton/PJ	5,0	100%	100%	25,00%	EMEP/EEA Guidebook 2013
Natural gas	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Other petrol products	ton/PJ	70	80%	70%	-	EMEP/EEA Guidebook 2013
Other fuels	ton/PJ	124	94%	87%	10,00%	EMEP/EEA Guidebook 2013

Heavy metals emissions are based on calculations (fuel consumed x emission factors per fuel type). Activity data are taken from the Flemish energy balance. Table 3-31 gives an overview of the emission factors that are used to calculate the emissions of the sectors included in category 1A2f.

Table 3-31 Emission factors used to calculate the heavy metals emissions for combustion in the sectors of non-metallic mineral product activities

Mineral non-metallurgical products	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Cokes	134	1,8	7,9	4	13,5	17,5	13	1,8	200	EMEP/EEA Guidebook 2013
Coal	134	1,8	7,9	4	13,5	17,5	13	1,8	200	EMEP/EEA Guidebook 2013
Heavy fuel	4,56	1,2	0,34 1	3,98	2,55	5,31	255	2,06	87,8	EMEP/EEA Guidebook 2013
Gas-and diesel oil	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
LPG	0,0015	0,00 025	0,1	0,12	0,00 076	0,00 0076	0,00 051	0,01 1	0,00 15	EMEP/EEA Guidebook 2013
Petrol										EMEP/EEA Guidebook 2013
Natural gas	0,0015	0,00 025	0,1	0,12	0,00 076	0,00 0076	0,00 051	0,01 1	0,00 15	EMEP/EEA Guidebook 2013
Other petrol products										EMEP/EEA Guidebook 2013
Other fuels	134	1,8	7,9	4	13,5	17,5	13	1,8	200	EMEP/EEA Guidebook 2013

In Wallonia, all the plants which are under the IPPC directive report their annual emissions.

#### **Other industries (category 1A2gviii)**

Category 1A2gviii is a key category of PM2.5, Cd, Ni and HCB emissions in terms of emissions level or trend.

In Flanders, emissions are reported directly by the individual companies. SO<sub>x</sub> emissions are calculated based on the sulphur content of the fuel or on measurements. NO<sub>x</sub>, CO and NMVOC emissions are measured, calculated or estimated based on plant specific information. To calculate the remainder of the emissions (emissions not reported directly by the individual companies) from the category 1A2gviii in Flanders a new methodology is used (Sleeuwaert et al., 2010). For a description of this methodology see above in chapter 1A2a, iron and steel. For this collective approach, for each sector in these categories and each fuel type a specific corresponding emission factor is used (see Table 3-32).

Table 3-32 Emission factors of CO, SO<sub>x</sub> and NO<sub>x</sub> in the other industries used in the collective approach

Metal products 1A2gviii	Unit	CO (g/GJ)	SO <sub>x</sub> (g/GJ)	NO <sub>x</sub> (g/GJ)
Cokes	g/GJ	82	683	204
LPG	g/GJ	44	0.0000435	73
Gas and dieseloil	g/GJ	59	47	126
Gasoline		57	46	123
Heavy fuel	g/GJ	58	493	170
Natural gas	g/GJ	43	0.0000450	43
Renewable fuels - solid	g/GJ	156	13	260
Renewable fuels - fluid	g/GJ	67	0.53	143
Textile, leather and clothing 1A2gviii	Unit	CO (g/GJ)	SO <sub>x</sub> (g/GJ)	NO <sub>x</sub> (g/GJ)
LPG	g/GJ	66	0.0000435	94
Gas and dieseloil	g/GJ	70	47	176
Heavy fuel	g/GJ	69	493	183
Natural gas	g/GJ	63	0.000045	47
Other industries 1A2f				
Coal	g/GJ	82	683	233
LPG	g/GJ	58	0.0000435	86
Gas and dieseloil	g/GJ	65	47	156
Heavy fuel	g/GJ	65	493	178
Natural gas	g/GJ	55	0.000045	46
Renewable fuels - solid	g/GJ	156	13	260

TSP emissions are based on calculations (fuel consumed x emission factors per fuel type). Activity data are taken from the Flemish energy balance. Emissions of PM<sub>10</sub> and PM<sub>2,5</sub> are calculated as a fraction of TSP and EC emissions are determined as a fraction of PM<sub>2,5</sub>.

The emission factors used to calculate the TSP emissions, the fractions of PM<sub>10</sub> and PM<sub>2,5</sub> and the EC fraction of PM<sub>2,5</sub> of the key categories is given below in Table 3-33:

Table 3-33 Emission factors used to calculate the TSP emissions, the fractions of PM10 and PM2,5 and the EC fraction of PM2.5 for combustion in the sectors of metal products, textile, leather and clothing and other industries in the Flemish region

Metal products 1A2gviii	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction	Source
Cokes	ton/PJ	124	94%	87%	6,40%	EMEP/EEA Guidebook 2013
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%	EMEP/EEA Guidebook 2013
Gas and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%	EMEP/EEA Guidebook 2013
LPG	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Natural gas	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Other fuels (i.e. H <sub>2</sub> )	ton/PJ				-	EMEP/EEA Guidebook 2013
Renewable solid	ton/PJ	150	95%	93%	28,00%	EMEP/EEA Guidebook 2013
Renewable liquid	Ton/PJ	6,47	50%	12,5%	33,50%	EMEP/EEA Guidebook 2013
Textile, leather and clothing 1A2gviii	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction	Source
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%	EMEP/EEA Guidebook 2013
Gas and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%	EMEP/EEA Guidebook 2013
Lamp petrol					45,00%	EMEP/EEA Guidebook 2013
LPG	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Natural gas	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Renewable fuels - solid	ton/PJ	150	95%	93%	28,00%	EMEP/EEA Guidebook 2013
Other industries 1A2gviii	Unit	Emission factor TSP	%PM10	%PM2,5	EC fraction	Source
Coal	ton/PJ	124	94%	87%	6,40%	EMEP/EEA Guidebook 2013
Heavy fuel	ton/PJ	35,40	71%	55%	5,60%	EMEP/EEA Guidebook 2013
Gas and diesel oil	ton/PJ	6,47	50%	12,5%	33,50%	EMEP/EEA Guidebook 2013
Petrol	ton/PJ	7,47	50%	12,5%	133,50%	EMEP/EEA Guidebook 2013

Lamp petrol	ton/PJ	5,00	100%	100%	45,00%	EMEP/EEA Guidebook 2013
LPG	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Natural gas	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Other fuels	ton/PJ	0,78	100%	100%	4,00%	EMEP/EEA Guidebook 2013
Renewable - solid	ton/PJ	150	95%	93%	28,00%	EMEP/EEA Guidebook 2013

Heavy metals emissions are based on calculations (fuel consumed x emission factors per fuel type). Activity data are taken from the Flemish energy balance. Table 3-34 gives an overview of the emission factors that are used to calculate the emissions of the sectors included in category 1A2f.

Table 3-34 Emission factors used to calculate the emissions of heavy metals for combustion in the sectors of metal products, textile, leather and clothing and other industries

<b>Metallurgical manufacturing</b>	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Cokes	134	1,8	7,9	4	13,5	17,5	13	1,8	200	EMEP/EEA Guidebook 2013
Heavy fuel	4,56	1,2	0,341	3,98	2,55	5,31	255	2,06	87,8	EMEP/EEA Guidebook 2013
Gas-and diesel oil	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
LPG	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Natural gas	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Other fuels										EMEP/EEA Guidebook 2013
Renewables - solid	27	13	0,56	0,19	23	6	2	0,5	512	EMEP/EEA Guidebook 2013
renewables - gaseous	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
<b>Textile, leather and clothing</b>	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Heavy fuel	4,56	1,2	0,341	3,98	2,55	5,31	255	2,06	87,8	EMEP/EEA Guidebook 2013
Gas-and diesel oil	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
LPG	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Natural gas	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Renewables - solid	27	13	0,56	0,19	23	6	2	0,5	512	EMEP/EEA Guidebook 2013
<b>Other industries (including wood manufacturing)</b>	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Source
Coal	134	1,8	7,9	4	13,5	17,5	13	1,8	200	EMEP/EEA Guidebook 2013
Heavy fuel	4,56	1,2	0,341	3,98	2,55	5,31	255	2,06	87,8	EMEP/EEA

										Guidebook 2013
Gas-and diesel oil	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
Petrol	4,07	1,36	1,36	1,81	1,36	2,72	1,36	6,79	1,81	EMEP/EEA Guidebook 2013
Lamp petrol										EMEP/EEA Guidebook 2013
LPG	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Natural gas	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Other fuels	0,001 5	0,000 25	0,1	0,12	0,000 76	0,000 076	0,000 51	0,011	0,001 5	EMEP/EEA Guidebook 2013
Renewables - solid	27	13	0,56	0,19	23	6	2	0,5	512	EMEP/EEA Guidebook 2013

In Wallonia, all the plants which are under the IPPC directive report their annual emissions. The emissions of the area source is calculated on the basis of the energy balance and the default emission factors.

The calculation of industry emissions in Brussels is based on the energy consumptions described in the regional energy balance and the emission factors mentioned in Table 3-24.

#### **Mobile Combustion in manufacturing industries and construction (category 1A2gvii)**

Off-road emissions are calculated by the same mathematical model OFFREM (Off-road emission model) (Schrooten et al., 2009) in the three regions. Emissions are calculated for machinery used in industry and building (category 1A2gvii). In Wallonia, some plants (cement plant, carriers,...) report their off-road emissions which are also included in 1A2gvii.

### **3.4. Transport (sector 1A3, 1A5b and off-road)**

#### **3.4.1. Source category description**

Belgium is provided with a very dense road (3.94 km/km<sup>2</sup>) and rail (117 m/km<sup>2</sup>) network (2009). These densities of road and rail networks should be looked at in conjunction with the very high density of population in Belgium: relative to the number of inhabitants the infrastructure is close to the European average. The port of Antwerp, located in the Flemish region, is very important for Belgium. It is the second largest European seaport, and one of the 5 largest in the world. The port of Antwerp benefits from excellent connections to the hinterland and the large French and German industrial basins by waterway (1500 km of navigable routes). It has also been decided to strengthen the rail infrastructure giving access to the port of Antwerp. Road transport is the mean of transport the most generally used in Belgium, both for the transport of goods and passengers, generating severe traffic congestion. Damages to the environment resulting from fuel use in road traffic are considerable. Goods (without pipelines) are transported by railways for 10.2% of total achieved tonne-kilometres in Belgium, on navigable waterways for 12.1% and by road transport for 77.7% (2009<sup>3</sup>).

<sup>3</sup>[http://economie.fgov.be/fr/modules/publications/statistiques/circulation\\_et\\_transport/transports\\_routiers\\_de\\_marchandises\\_-\\_overview.jsp](http://economie.fgov.be/fr/modules/publications/statistiques/circulation_et_transport/transports_routiers_de_marchandises_-_overview.jsp)

The reported emissions in the transport sector are reported in the categories 1A3a Civil aviation, 1A3b Road transport, 1A3c Railways, 1A3d Navigation and 1A3e Other transportation.

In the category 1A3e the emissions originating from the transport of natural gas through pipelines are allocated as well as emissions of off-road machinery in harbours, airports and due to storage and handling.

No civil aviation takes place in the Brussels region, Brussels Airport is located on the Flemish territory.

Emissions of the military aviation are allocated to the category 1A5b.

Sea navigation takes only place in the Flemish region.

### 3.4.2. Methodological issues

#### 3.4.2.1 Road transport (1A3b)

Until the 2013 submission, the 3 regions used COPERT 4 methodologies in specific regional models (previous versions of COPERT 4 were used in the Walloon and the Brussels regions, MIMOSA was used in Flemish region). Moreover the process to transfer the basic data of the Belgian vehicle fleet to a regional fleet file that serves as input for the regional models was performed separately for the 3 regions.

Since 2014, regional submissions are almost fully harmonised:

- Each region uses a common fleet module (produced by Transport & Mobility Leuven (TML)) which provides harmonised regional fleet files as input for COPERT. The vehicles fleet is based on actual registration data for the period 2007-2013 and on a combination of fleet data and backwards extrapolations for the period 1990-2006.
- Each region uses the same module for the assignment of mobility (produced by IRCEL-CELINE).
- Each region uses directly COPERT 4 v11.1bis to produce regional emissions and “fuel used” consumptions.

The 2 major determinants of COPERT modelling (fleet and mobility) are now fully harmonized across the 3 regions. However, for some parameters regional data based on specific measurements and/or modelling have been used (average trip length and duration, driving modes and corresponding average speed values).

The emissions from road transport have been recalculated for the whole time series 1990-2013 using the COPERT IV v11.1bis software. For PM, a post process correction was applied to account for the retrofitted particulate filters for Euro 4 passenger cars and Euro II and Euro III urban busses.

Because the COPERT results for non-exhaust PM emissions do not allow to differentiate “tyre & brake wear” and “road abrasion” emissions, the non-exhaust PM emissions are calculated based on the emission factors from the 2013 EMEP Guidebook.

The consumption modelling in COPERT is entirely based on the energy content of fossil fuels. The use of air-conditioning is included.

Since 2015, the emissions are reported as “fuel sold”. The fuel sold emission values are obtained by scaling up all the emissions with a specific scaling factor per year and per fuel. The scaling factor is obtained by dividing for each year and given fuel (Gasoline, Diesel and LPG), the total amount of fuel from the federal fuel balance report (FOD/SPF Mobility & Transport) by the total fuel used of the 3 regions.

Figure 3.3 shows the trend of the energy consumption in the road transport sector.




Figure 3.3 Trend of fuel used in road transport.

#### 3.4.2.2 Air transport (1A3a)

In the two regions (Flemish and Walloon region) where air transport is relevant, a slightly different approach was applied in estimating the emissions from air transport.

In Flanders, emissions of civil aviation are calculated for Landing and Take-Off cycles (LTO) based on flight data (data per flight, statistics delivered by the airports) and emission factors. These emissions are calculated for 4 airports for civil aviation (Antwerp, Ostend, the international airport Brussels Airport and from 2008 on also Kortrijk-Wevelgem) and for 6 airports for military aviation between 1990 and 1996 (Kleine Brogel, Brasschaat, Koksijde, Sint-Truiden, Goetsenhove and Melsbroek) and 4 airports for military aviation from 1997 until 2011 (Kleine Brogel, Brasschaat, Koksijde, Melsbroek). The airport of origin and airport of destination are known, a division in domestic/international traffic can be made. Emission factors from the EMEP/EEA Guidebook (2009 table 3-5) are used as well as emission factors from EPA.

In Wallonia, the emissions from civil aviation (category 1A3a) are estimated on the basis of the number of LTO of the two main airports (Liège and Charleroi) and the amount of LTO in small airports. Each year, the two main airports report the jet fuel and the gasoline consumptions for the domestic and for the international activities. They also report the amount of domestic LTO and the amount of international LTO. Some information on the Walloon total number of LTO is available (Liège and Charleroi airports flights, training flights,...). Unfortunately, there is no general knowledge about the aircraft types carrying out the aviation activities. Emission factors used to estimate emissions from domestic and international traffic are based on the table 3-3 in the EMEP/EEA guidebook 2009. The specific energy consumption by LTO is 105 kg fuel/LTO for the domestic flight.

For the international flight in one of the two international airports, the specific energy consumption by LTO is 3400 kg fuel/LTO instead of 825 kg fuel/LTO as the planes are mainly cargo planes.

The emissions from domestic LTO and international LTO are reported under the category 1A3ai(i) and 1A3aii(i), while emissions from cruise activities are reported under 'Memo items' 1A3ai(ii) and 1A3aii(ii).

### 3.4.2.3 Railways

The emissions of railway traffic are estimated by a region specific approach.

Flemish emissions of railway traffic are estimated by the EMMOSS model (Vanherle et al., 2007, 2010). The basis for the calculations is gross ton kilometres driven by trains.

Emission calculation:

$$EM(g) = \text{gross ton kilometers} \left( \frac{\text{ton}}{\text{km}} \right) \times \text{specific end - energy use} \left( \text{kWh} \cdot \frac{\text{km}}{\text{ton}} \right) \times EF \left( \frac{\text{g}}{\text{kWh}} \right)$$

Emission factors are derived from ISO 8178/F test cycles for CO, NO<sub>x</sub>, TSP and VOC (Table 3-35).

Table 3-35 Emission factors for different train types (g/kWh) to calculate emissions from rail transport in Flemish Region

	Type HLD77	Type MW41	old locos	old railcars
CO	0.73	1.07	10.70	10.70
NO <sub>x</sub>	11.70	8.74	18.20	18.20
TSP	0.20	0.15	0.60	0.60
VOC	0.11	0.61	1.60	1.60

Emissions for NH<sub>3</sub> and PAH were taken over from Klein, 2006 (The Netherlands) (Table 3-36). SO<sub>2</sub> and heavy metals are fuel-specific (SO<sub>2</sub> calculated dependent on content of S in fuel).

Table 3-36 Emission factors from Klein, 2006 (NL) to calculate emissions from rail transport in Flemish Region

Pollutant	EF(g/g or %)	calculation base off
NH <sub>3</sub>	0,00001	kgFC
Cd	0,00000001	kgFC
Cu	0,0000017	kgFC
Cr	0,00000005	kgFC
Ni	0,00000007	kgFC
Se	0,00000001	kgFC
Zn	0,000001	kgFC
benz(b)fluoranteen	0,0000169	fractionVOC
benz(k)fluoranteen	0,00000643	fractionVOC
benz(a)pyreen	0,0000169	fractionVOC

Indeno(1,2,3-cd)-pyrene	0	fractionVOC
PM2.5	0,95	fractionPM
PM10	1	fractionPM

Emissions for shunting trains are also calculated. Emissions are reported in the NFR category 1A3c (railways).

The emissions from railways in the Walloon region and in the Brussels region are estimated by multiplying the train's fuel consumption by the fuel specific emission factors. The emission factors are those described in table 8.1 of the EMEP/EEA guidebook 2013.

In Wallonia and Brussels, the data from the National Society of the Belgian Railways (SNCB-NMBS) are used to calculate the energy consumption for the train services in Belgium. These data are available for the transport of persons and goods and for electricity and gasoil driving. The total consumption of gasoil in the Walloon and Brussels regions is based on the Belgian data of gasoil consumption and the regional information on driven train- and ton-kilometres of persons and goods. The emissions are estimated by multiplying the train's fuel consumption by the fuel specific emission factors (Table 3-37). Following the study of VITO on Heavy Metals, it must also take into account the wear catenary (Cu : 961 mg/GJ).

Table 3-37 Emission factors in the Walloon region (EMEP/EEA Guidebook 2013)

		<b>Gasoil</b>
<b>SO2</b>	g/GJ	48.00
<b>NOx</b>	g/GJ	1227.17
<b>COVNM</b>	g/GJ	108.90
<b>CO</b>	g/GJ	250.59
<b>Dust</b>	g/GJ	35.60
<b>PM10</b>	g/GJ	33.72
<b>PM2.5</b>	g/GJ	32.08
<b>BC</b>	g/GJ	23.14
<b>Cd</b>	mg/GJ	0.23
<b>Cr</b>	mg/GJ	1.17
<b>Cu</b>	mg/GJ	39.81
<b>Ni</b>	mg/GJ	1.64
<b>Pb</b>	mg/GJ	
<b>Se</b>	mg/GJ	0.23
<b>Zn</b>	mg/GJ	23.42
<b>NH3</b>	mg/GJ	0.16
<b>HAP</b>	mg/GJ	2.78
<b>Dioxines</b>	ng/GJ	1442.86

#### 3.4.2.4 Navigation

Category 1A3di(ii) International inland waterways is a key category of NO<sub>x</sub> emissions in terms of emissions trend.

For navigation, fuel consumption is taken from the regional energy balances. In Flanders, emissions from maritime navigation are calculated with the emission model EMMOSS. The emissions originating from maritime shipping starting and arriving in Belgium (including sand extraction at sea, dredging activities and tugboats) are reported in the category 1A3di(ii) (international inland waterways). The emissions coming from maritime shipping between a Flemish and a foreign harbour (including emissions originating in the Flemish harbour) are reported in the memo item 1A3di(i) 'international maritime navigation'.

Emissions are calculated using emission factors from the Dutch methodology, taking into account IMO Tier II and Tier III NO<sub>x</sub> limits as stated in Marpol Annex VI (for maritime navigation).

The source of emission factors :

- NO<sub>x</sub>, VOC, TSP, CO : Dutch EMS protocol (Oonk, 2003)
- NH<sub>3</sub>, PAH : Dutch study Klein, 2006
- PM<sub>2.5</sub> and PM<sub>10</sub> : % of TSP from Visschedijk et al. (NI)

The Belgian maritime zones comprise the territorial sea (TS) and the Exclusive Economic Zone (EEZ). The former consists of an area extending 12 nautic miles into the North Sea, measured from the base line. The latter comprises that part of the North Sea the contour of which consists of lines connecting following points in the order of numeration:

- |    |                 |   |                 |
|----|-----------------|---|-----------------|
| 1. | 51°16'09" N     | – | 02°23'25" O     |
| 2. | 51°33'28" N     | – | 02°14'18" O     |
| 3. | 51°36'47" N     | – | 02°15'12" O     |
| 4. | 51°48'18" N     | – | 02°28'54" O     |
| 5. | 51°52'34.012" N | – | 02°32'21,599" O |
| 6. | 51°33'06" N     | – | 03°04'53" O     |

A map of the Belgian maritime areas (Figure 3.4 Map of the Belgian maritime areas) is shown below




Figure 3.4 Map of the Belgian maritime areas

In Flanders the emissions originating from inland shipping are also estimated by the EMMOSS model and are reported in the IPCC category 1A3dii (national navigation). Category 1A3dii is a key category of NO<sub>x</sub> emissions in terms of emissions level and trend.

Emission factors for NO<sub>x</sub>, PM, CO, VOS are derived from Oonk et al. (2003), and from 2007 on derived from the CCNR-standards (Central Commission for the navigation of the Rhine). Other are emission factors of the EMEP/EEA handbook (Table 3-38 and Table 3-39).

Table 3-38 Emission factors dependent on year class of the engines (g/kWh)

Motorbouwjaar	NO <sub>x</sub>	PM	CO	NM <sub>VOC</sub>
< 1974	10	0,6	4,5	1,2
1975-1979	13	0,6	3,7	0,8
1980-1984	15	0,6	3,1	0,7
1985-1989	16	0,5	2,6	0,6
1990-1994	14	0,4	2,2	0,5
1995-2001	11	0,3	1,8	0,4
2002-2007	8	0,3	1,5	0,3
2007-2011 <sup>2</sup>	6	0,2	1,3	0,2
2011-2015 <sup>3</sup>	6	0,2	1,3	0,2
2015-2020 <sup>3</sup>	6	0,2	1,3	0,2
2020-2030 <sup>3</sup>	6	0,2	1,3	0,2

Table 3-39 Emission factors based on fuel used (g/kg fuel)

	Emission factor	
SO <sub>2</sub>	$20 \cdot S\%^2$	Base stoichiometric conversion
NH <sub>3</sub>	0,007	EMEP EEA guidebook
Cd	0,00001	EMEP EEA guidebook
Cr	0,00005	EMEP EEA guidebook
Cu	0,0017	EMEP EEA guidebook
Ni	0,00007	EMEP EEA guidebook
Pb	0,01	EMEP EEA guidebook
Zn	0,001	EMEP EEA guidebook

The emissions from inland navigation are estimated in the Walloon and Brussels region by multiplying the sector's fuel consumption by the fuel specific emission factors. The emission factors are those described in the EMEP/EEA Guidebook 2013.

#### 3.4.2.5 Other transportation (pipeline compressors 1A3ei and off-road 1A3eii)

Category 1A3ei includes the emissions from the pipeline compressors. In Flanders CO and NO<sub>x</sub> emissions are provided by the operators of the plants.

In the Walloon region, this category includes also the emissions from the pipeline compressors. Since 2008, a IPPC plant has reported CO and NO<sub>x</sub> emissions and default emission factors have been calculated with these data (Table 3-40). These default emission factors are used for the years before 2008 and for the area part after 2008.

Table 3-40 Emission factors for pipeline compressors in the Walloon region

Pollutant	UNIT	EF
<b>NO<sub>x</sub></b>	g/GJ	193
<b>CO</b>	g/GJ	260

As a result of the In country review in September 2012 of the GES Belgium inventory and to be coherent with this GES inventory, the off-road emissions of the following sectors are now included in the category 1A3eii: ground activities in airports, harbours and trans-shipment activities. Off-road emissions are calculated by the same mathematical model OFFREM (Off-road emission model) (Schrooten et al., 2009) in the three regions.

### **3.5. Other sectors (sector 1A4)**

#### **3.5.1. Source category description (1A4)**

In the category 1A4 the following sources are taken into account in the Belgian atmospheric pollutant inventory: commercial/institutional (1A4a), residential (1A4b) and agriculture/forestry/fishery (1A4c).

For the 3 regions emissions from the off-road sector are included in the categories 1A3e, 1A4b, 1A4c and 1A5b.

#### **3.5.2. Methodological issues**

##### **Commercial/institutional sector (stationary, category 1A4ai)**

Category 1A4ai is a key category of NO<sub>x</sub> emissions in terms of emissions level and trend and of SO<sub>2</sub> emissions in terms of emissions level. The fuel consumption of the stationary combustion in the commercial/institutional sector is based on general statistics of natural gas, supplemented with results from surveys for solid and liquid fuels. The energy use in the commercial/institutional sector is strongly related to the climate (cold winters cause higher energy consumption and hence higher emissions). The relatively warm winter of 2011 is reflected by a lower energy consumption (mostly gaseous and liquid fuels).

The energy consumption of these sectors is published in the regional energy balances.

The following chart (Figure 3.5) shows the trends of the energy consumption in the commercial/institutional sector.




Figure 3.5 Trend of fuel consumption in the commercial/institutional sector.

In Flanders, emissions by heating systems of buildings are calculated on a collective basis. The database consists of emissions due to tertiary combustion (heating by hotels/restaurants, medical services, education, offices and administrative activities, trade, other services and combined heat-power installations (CHP)). Emissions are calculated by multiplying the energy use and emission factors. Data on energy used can be found in the Energy Balance for Flanders. A provisional energy balance is made yearly for year (x-1), whereas a final energy balance is made for year (x-2). The tertiary sector contains energy data on natural gas, fuel, heavy fuel, solid fuels (coal), propane/butane/LPG, electricity, other (mainly waste) and renewable fuels, for some years also lamp petrol. The emission factors to calculate  $\text{NO}_x$  and NMVOC originate from studies,  $\text{SO}_2$  emission calculations are based on the S-content of the fuels (for  $\text{SO}_2$  and  $\text{NO}_x$ , the emission factors are harmonised between the regions), other emission factors ( $\text{CO}$ , particulate matter, heavy metals and  $\text{NH}_3$ ) are taken from the EMEP/EEA Emission Inventory Guidebook 2013.

Emissions and activity data due to combined heat-power installations in joint venture with the energy sector are allocated in NFR sector 1A1a (see also 3.2.1). For the CHP installations in the tertiary sector energy information on natural gas, fuel and other fuels (biogas) is included. A distinction is made between autoproducers and non-autoproducers.

In 2013 emission factors are re-examined as a result of the release of the revised Guidebook. The emission factors are only adapted when expert analysis reveals that better factors are available or when tuning with the other Belgian regions occurs.

In Wallonia, the main data source for this sector is the energy balance delivered yearly by the Energy and Sustainable Building Department. The energy balance describes the quantities of energy imported, produced, transformed and consumed in the Walloon Region in a given year. The energy consumption in the service sector is calculated using the energy data of different sources (regional data on the amount of natural gas and electricity sold in this sector (CWaPE), annual survey carried out by ICEDD for all consumers 'high voltage' (4800 establishments with a respond of 58 %)).

In Brussels, the consumption of the tertiary sector is based on two approaches: one for the 'high voltage customers' and the other for the 'low voltage customers'. For 'high-voltage customers', the energy consumption is calculated using the energy data based on a survey and direct contacts with 'high voltage' consumers and big international public organisms. For 'low-voltage customers', the energy consumption is calculated by the 'top down' method and the consumption of oil-products is estimated from the fuel/natural gas ratio and the Belgian consumptions.

Emission factors used to calculate the emissions of stationary combustion in the commercial sector for the year 2013 in the three regions are given below (Table 3-41 to Table 3-43).

Table 3-41 Emission factors for the sector 1A4a in the Walloon region (EMEP/EEA Guidebook except NO<sub>x</sub> and dust).

	SO <sub>2</sub>	NOx	COVNM	CO	TSP	PM10	PM2,5	BC	NH3					
	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ	g/GJ					
Coal	600	104	89	931	124	117	108	6.9	0.4					
Wood	11	77.5-77.9	300	570	150	143	140	39.2	37					
Diesel oil	48	42.3-43	10	66	5.0	3.9	3.0	1.7	0.1					
Natural gas	0.5	36.1-40.3	0.36	24	0.45	0.45	0.45	0.02	0.6					
Lamp petroleum		60	10	66	5.0	3.9	3.0	1.7	0.1					
LPG		40	0.36	24	0.45	0.45	0.45	0.02	0.6					
	As	Cd	Cu	Cr	Ni	Pb	Sé	Zn	Hg	Dio	HAP	PCB	HCB	
	mg/GJ										ng/GJ	mg/GJ	yg/GJ	yg/GJ
Coal	4	1.8	17.5	13.5	13	134	1.8	200	7.9	203	146.6	170	0.62	
Wood	0.19	13	6	23	2	27	0.5	512	0.56	100	35	0.06	5	
Diesel oil	0.03	0.006	0.22	0.2	0.008	0.08	0.11	29	0.12	1.4	0.02			
Natural gas	0.12	0.00025	8E-05	0.0008	5E-04	0.0015	0.011	0	0.1	0.5	0.003			
Lamp petroleum	0.03	0.006	0.22	0.2	0.008	0.08	0.11	29	0.12	1.4	0.02			
LPG	0.12	0.00025	8E-05	0.0008	5E-04	0.0015	0.011	0	0.1	0.5	0.003			

Table 3-42 Emission factors for the sector 1A4a in the Flemish region.

Fuel	UNIT	SO <sub>2</sub> *	NO <sub>x</sub> *	COV	CO	NH <sub>3</sub>	TSP*	PM10*	PM2,5*	EC
coal	g/GJ	600	104,44 2	200,00	931	-	124	117	108	10.8
wood	g/GJ	6	78,978	500,00	570	37	150	143	140	49
diesel oil	g/GJ	48	40.576	3,00	66	-	4.582	4.582	4.582	1.375
heavy fuel	g/GJ	480	200,00 0	3,00	15.1	-	35.4	25.2	19.3	1.93
natural gas	g/GJ	0.3	27.653	2,00	29	-	0.45	0.45	0.45	.032
LPG	g/GJ	0	40,000	2,00	29	-	0.45	0.45	0.45	.0585
Source		S-content and regional tuning	Study FOD	Study VITO	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	TNO

\*emission factors vary with time. The table shows the emission factors used to calculate emissions 2013

Fuel	UNIT	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn
coal	g/	0.134	0.0018	0.0079	0.004	0.0135	0.0175	0.013	0.001	0.2

	GJ								8	
wood	g/GJ	0.027	0.013	0.00056	0.00019	0.023	0.006	0,002	0,0005	0.512
diesel oil	g/GJ	0.000012	0.000001	0.00012	0.000002	0.0002	0.00013	0.000005	0.000002	0.00042
heavy fuel	g/GJ	0.00456	0.0012	0.00034	0.00398	0.00255	0.00531	0.255	0.00206	0.0878
lamp petrol	g/GJ	0.000012	0.000001	0.00012	0.000002	0.0002	0.00013	0.000005	0.000002	0.00042
natural gas	g/GJ	0.0000015	0.00000025	0.0001	0.00012	0.0000076	0.00000076	0.00000051	0.0000011	0.0000015
LPG	g/GJ	0.0000015	0.00000025	0.0001	0.00012	0.0000076	0.00000076	0.00000051	0.0000011	0.0000015
Source		EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA

Fuel	benzo(a)pyrene (kg/TJ)	benzo(b)fluoranthene (kg/TJ)	benzo(k)fluoranthene (kg/TJ)	hexachlorbenzene (kg/TJ)	indeno(1,2,3-cd)pyrene (kg/TJ)	PCDD/F (mg TEQ/TJ)
natural gas						0,0005
diesel oil	0,000101	5,38E-05	8,19E-05		0,000201	0,0005
heavy fuel	0,002118	0,001108	0,001749		0,004187	0,0025
coal	0,000404	0,000844	0,000618	5,46E-07	0,000167	0,01
LPG						0,0005
wood	0,017	0,014333	0,008833	0,000004	0,010933	0,05
lamp petrol	0,000101	5,38E-05	8,19E-05		0,000201	0,0005
Source	TNO	TNO	TNO	EMEP/CORINAIR	TNO	UNEP

Table 3-43 Emission factors for the sector 1.A.4.a in the Brussels region.

Fuel	UNIT	NOx	NM VOC	SOx	NH3	PM2.5	PM10	TSP	BC (EC)	CO	PCDD/PCDF*
Natural gas	g/GJ	34,5	23	0,67	0,6	0,78	0,78	0,78	0,031	29	0,52
Gas oil	g/GJ	42,29	25	47	0,1	20	20	20	11,2	66	1,4
Wood	g/GJ	91	300	11	37	140	143	150	39,2	570	100
Butane/Propane	g/GJ	40	23	0,67	0,6	0,78	0,78	0,78	0,031	29	0,52
Fuel	UNIT	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Total HAP
Natural gas	mg/GJ	0,011	0,0009	0,54	0,1	0,013	0,0026	0,013	0,058	0,73	0,0058
Gas oil	mg/GJ	0,08	0,006	0,12	0,03	0,2	0,22	0,008	0,11	29	0,0201
Wood	mg/GJ	27	13	0,56	0,19	23	6	2	0,5	512	35
Butane/Propane	mg/GJ	0,011	0,0009	0,54	0,1	0,013	0,0026	0,013	0,058	0,73	0,0058

\* ng-TEQ/GJ

### **Residential sector (stationary, category 1A4bi)**

Category 1A4bi is a key category of NO<sub>x</sub>, NMVOC, SO<sub>x</sub>, CO and PM2,5, PM10, TSP, Cd, Hg, As, Cr, Ni, Zn, PCDD/F and HCB emissions in terms of emissions level and trend and of NH<sub>3</sub>, Pb and Se emissions in terms of emissions level or trend. The fuel consumption of the stationary combustion in the residential sector is based on general statistics of natural gas, supplemented with results from surveys for solid and liquid fuels. The energy use in the households is strongly related to the climate (cold winters cause higher energy consumption and hence higher emissions). The relatively warm winter of 2011 is reflected by a lower energy consumption (mostly gaseous and liquid fuels)

The following chart (Figure 3.6) shows the trends of the energy consumption in the residential sector.




Figure 3.6 Trend of fuel consumption in the residential sector.

In Flanders, emissions by heating systems of buildings are calculated on a collective basis. The database consists of emissions due to residential combustion (heating by households). Emissions are calculated by multiplying the energy use and emission factors.

Data on energy used can be found in the Energy Balance for Flanders. A provisional energy balance is made yearly for year (x-1), whereas a final energy balance is made for year (x-2). For households energy data on electricity use, natural gas, fuel, solid fuels (coal), propane/butane/LPG, renewable fuels (mainly wood) are included. SO<sub>x</sub> emission calculations are based on the S-content of the fuels (for SO<sub>x</sub> and NO<sub>x</sub>, the emission factors are harmonised between the regions), other emission factors are taken from the EMEP/EEA Emission Inventory Guidebook. In 2013 emission factors are re-examined as a result of the release of the revised Guidebook. The emission factors are only adapted when expert analysis reveals that better factors are available or when tuning with the other Belgian regions occurs.

In Wallonia, the main data source for this sector is the energy balance delivered yearly by the Energy and Sustainable Building Department. The energy balance describes the quantities of energy imported, produced, transformed and consumed in the Walloon Region in a given year. The energy consumption of the household sector is calculated on the basis of regional data on the amount of natural gas and electricity sold in this sector (CWaPE), on the basis of national data (liquid fuels and solid fuels), on the basis of the socio-economic survey of 2001 (size, isolation,...) and on the basis of weather data (degree-days).

In Brussels, the information about energy consumption in the household sector is compiled in the regional energy balance. Then the consumption is multiplied for the emission factors described in Table 3-46. The emission factors presented here are the emission factors used in 2013. The emission factors for NO<sub>x</sub> change each year in the residential sector as they follow the evolution of boilers (park and performance) in this sector.

For the 2015 submission, the three regions agreed to use the emission factors for wood combustion from the Guidebook 2013 for estimating the wood emissions for the residential sector.

Emission factors used to calculate the emissions of stationary combustion in the residential sector for the year 2013 in the three regions are given below (Table 3-44 to Table 3-46).

Table 3-44 Emission factors for the sector 1A4bi in the Walloon region for 2013 (EMEP/EEA Guidebook 2013 except NO<sub>x</sub> and dust)

Guidbook 2016 except NO <sub>x</sub> and dust																										
	SO <sub>2</sub>		NO <sub>x</sub>		NMVOC		CO		TSP		PM <sub>10</sub>		PM <sub>2,5</sub>		BC		NH <sub>3</sub>									
	g/GJ		g/GJ		g/GJ		g/GJ		g/GJ		g/GJ		g/GJ		g/GJ		g/GJ									
Coal	600		104		484		4 600		444		404		398		25.472		0.3									
Wood	11		80		600		4 000		800.0		760.0		740.0		74.0		70									
Diesel oil	48		42.4-43		0.69		57		1.9		1.9		1.9		0.2		0.1									
Natural gas	0.3		30-40		1.9		26		1.2		1.2		1.2		0.1		0.6									
LPG	0.3		40		1.9		26		1.2		1.2		1.2		0.1		0.6									
	As		Cd		Cu		Cr		Ni		Pb		Sé		Zn		Hg		Dio		HAP		PCB		HCB	
	mg/GJ														ng/GJ		mg/GJ		yg/GJ		yg/GJ					
Coal	2.5		1.5		22.3		11.2		12.7		130		1		220		5.1		800		800		170		0.62	
Wood	0.19		13		6		23		2		27		0.5		512		0.56		800		345		0.06		5	
Diesel oil	0.002		0.001		0.13		0.2		0.005		0.012		0.002		0.42		0.12		5.9		0.35					
Natural gas	0.12		0.0003		7.6E-05		0.0008		0.00051		0.0015		0.011		0.002		0.1		1.5		0.003					
LPG	0.12		0.0003		7.6E-05		0.0008		0.00051		0.0015		0.011		0.002		0.68		1.5		0.003					

Table 3-45 Emission factors for the sector 1A4bi in the Flemish region.

Fuel	UNIT	SO <sub>2</sub> *	NO <sub>x</sub> *	COV	CO	NH <sub>3</sub>	TSP*	PM <sub>10</sub> *	PM <sub>2,5</sub> *	EC
coal	g/GJ	600	104,442	200	4600	0,3	444	404	398	139.3
Wood	g/GJ	11	80	600	4000	70	800	760	740	74
diesel oil	g/GJ	48	40.576	3	57	0	1.9	1.9	1.9	0.570
natural gas	g/GJ	0.3	27.676	2	26	0	1.2	1.2	1.2	0.084
LPG	g/GJ	0	40	2	26	0	1.2	1.2	1.2	0.156
Source		S-content and regional tuning or EMEP/EEA (natural gas)	Study FOD	Study VITO	EMEP /EEA	EMEP /EEA	EMEP/EEA	EMEP/EEA	EMEP/EEA	TNO

\*emission factors vary with time. The table shows the emission factors used to calculate emissions 2013

Fuel	UN IT	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn
coal	g/G J	0,13	0,0015	0,0051	0,0025	0,0112	0,0223	0,0127	0,001	0,22
wood	g/G J	0,027	0,013	0,0005 6	0,0001 9	0,023	0,006	0,002	0,0005	0,512
diesel oil	g/G J	0.0000 12	0.00000 1	0.0001 2	0.0000 02	0.0002	0.00013	0.00000 5	0.0000 02	0.0004 2
natural gas	g/G J	0.0000 015	0.00000 025	0.0001	0.0001 2	0.00000 076	0.00000 0076	0.00000 051	0.0000 11	0.0000 015
LPG	g/G J	0.0000 015	0.00000 025	0.0001	0.0001 2	0.00000 076	0.00000 0076	0.00000 051	0.0000 11	0.0000 015
Source		EMEP/ EEA	EMEP/ EEA	EMEP/ EEA	EMEP/ EEA	EMEP/ EEA	EMEP/E EA	EMEP/ EEA	EMEP/ EEA	EMEP/ EEA

Fuel	Polluent	Unit	Value	Source
Coal	PCDD/F	µg/GJ	0,070	UNEP Standardized Toolkit; Category 3e1: Power Generation and Heating; Domestizing heating - Fossil fuels; Coal fired stoves
Wood	PCDD/F	µg/GJ	0,800	EMEP2013 tier1 residential plants using biomass
Diesel oil	PCDD/F	µg/GJ	0,010	UNEP Standardized Toolkit; Category 3e2: Power Generation and Heating; Domestizing heating - Fossil fuels; Oil fired stoves
Natural gas	PCDD/F	µg/GJ	0,0015	UNEP Standardized Toolkit; Category 3e3: Power Generation and Heating; Domestizing heating - Fossil fuels; Natural gas fired stoves
LPG	PCDD/F	µg/GJ	0,0015	UNEP Standardized Toolkit; Category 3e3: Power Generation and Heating; Domestizing heating - Fossil fuels; Natural gas fired stoves
Coal	HCB	g/GJ	0,000000 55	EMEP/CORINAIR Guidebook (2005)
Wood	HCB	g/GJ	0,000005	EMEP2013 tier1 residential plants using biomass

Fuel	benzo(a)pyreen (kg/TJ)	benzo(b)fluorantheen (kg/TJ)	benzo(k)fluorantheen (kg/TJ)	indeno(1,2,3- cd)pyreen (kg/TJ)	PCB's (polychloorbiphenyls) (kg/TJ)
natural gas					
diesel oil	0,000101	5,38E-05	8,19E-05	0,000201	
coal	0,007006	0,022293	0,019745	0,006688	
LPG					
Wood*	0,121	0,111	0,042	0,071	6E-08
source	TNO	TNO	TNO	TNO	TNO

\*wood:EMEP/EEA

Table 3-46 Emission factors for the sector 1A4bi in the Brussels region.

Fuel	UNIT	NOx	NM VOC	SOx	NH3	PM2.5	PM10	TSP	BC (EC)	CO	PCDD/PCDF*
Natural gas	g/GJ	29	1,9	0,3	0,6	1,2	1,2	1,2	0,065	26	1,5
Gas oil	g/GJ	42,37	0,69	70	0,1	1,9	1,9	1,9	0,1615	57	5,9
Wood	g/GJ	80	600	11	70	740	760	800	74	4000	800
Coal	g/GJ	104,4	484	900	0,3	398	404	444	25,472	4600	800
Butane/Propane	g/GJ	40	1,9	0,3	0,6	1,2	1,2	1,2	0,065	26	1,5
Fuel	UNIT	Pb	Cd	Hg	As	Cr	Cu	Ni	Se	Zn	Total HAP
Natural gas	mg/GJ	0,002	0,00025	0,1	0,12	0,0008	7,6E-05	0,0005	0,011	0,002	0,00308
Gas oil	mg/GJ	0,012	0,001	0,12	0,002	0,2	0,13	0,005	0,002	0,42	0,35
Wood	mg/GJ	27	13	0,56	0,19	23	6	2	0,5	512	345
Coal	mg/GJ	130	1,5	5,1	2,5	11,2	22,3	12,7	1	220	800
Butane/Propane	mg/GJ	0,002	0,00025	0,1	0,12	0,0008	7,6E-05	0,0005	0,011	0,002	0,00308

\* ng-TEQ/GJ

#### **Agriculture/forestry/fishery (stationary, category 1A4ci)**

Category 1A4ci is a key category of SO<sub>x</sub> emissions in terms of emissions trend.

Agricultural fuel consumption is estimated from statistical information concerning area used, etc., combined with specific energy consumption from literature. The following chart (Figure 3.7) shows the trends of the energy consumption in the agricultural sector.




Figure 3.7 Trend of fuel consumption in the agricultural sector.

The sector 1A4ci Agriculture/Forestry/Fishing (stationary combustion) includes the emissions originating from greenhouse culture, arable farming- intensive livestock breeding, remaining crops – soil-bound agriculture and pasture. The activity data (energy consumption data) of the sectors 1A4ci are taken from the regional energy balances. Emission factors originate from the EMEP/EEA Guidebook, are region specific (NMVOC, TSP, PM10, PM2,5) or derived from the S-content and regional tuning (SO<sub>x</sub>). Table 3-47 give an overview of the emission factors used in 2013 in Flanders.

Table 3-47 Emission factors used for the sector 1A4ci in the Flemish region in 2013.

Stationary combustion	natural gas	diesel oil	heavy fuel	solid fuels	LPG	palm oil and rape oil	biogas	wood	petrol
CO (kg/TJ)	30	66	66	931	30	66	30	570	66
NOx (kg/TJ)	64	50	125	300	40	50	64	91	50
CxHy (kg/TJ)	2	3	3	200	2	2,5	2	300	3
stof (kg/TJ)	0.45	5	60	124	0.45	5	0.45	150	5
PM10 (kg/TJ)	0.45	5	50	117	0.45	5	0.45	143	5
SOx (kg/TJ)	0.3	48	480	648	0.3	95	0.3	11	95
Pb (kg/TJ)	0.0000015	0.00008	0.00008	0,134	0,0000015	0,00008	0,0000015	0,027	0,00008
NH3 (kg/TJ)								37	
As (kg/TJ)	0.00012	0.00003	0,00003	0,004	0,00012	0,00003	0,0012	0.00019	0.00003
Cd (kg/TJ)	0,000000025	0,000006	0,000006	0,0018	0,000000025	0,000006	0,000000025	0,013	0,000006
Cr (kg/TJ)	0,000000076	0,0002	0,0002	0,0135	0,000000076	0,0002	0,000000076	0,023	0,0002
Cu (kg/TJ)	0,0000000076	0,00022	0,00022	0,0175	0,0000000076	0,00022	0,0000000076	0,006	0,00022
Hg (kg/TJ)	0,00010	0,00012	0,00012	0,0079	0,00010	0,00012	0,00010	0,00056	0,00012
Ni (kg/TJ)	0,000000051	0,000008	0,000008	0,013	0,000000051	0,000008	0,000000051	0,002	0,000008
Se (kg/TJ)	0,000011	0,00011	0,00011	0,0018	0,000011	0,00011	0,000011	0,0005	0,00011
Zn (kg/TJ)	0,0000015	0,029	0,029	0,200	0,0000015	0,029	0,0000015	0,512	0,029
PM2,5 (kg/TJ)	0,45	5	40,000	108	0,45	5	0,45	140	5

The Table 3-48 gives an overview of the emission factors used in 2013 in Wallonia.

Table 3-48 Emission factors for the sector 1A4ci in the Walloon region (EMEP/EEA Guidebook 2013)

		<b>Gasoil</b>
<b>SO<sub>2</sub></b>	g/GJ	48
<b>NOx</b>	g/GJ	163
<b>COVNM</b>	g/GJ	10
<b>CO</b>	g/GJ	66
<b>TSP</b>	g/GJ	5.0
<b>PM10</b>	g/GJ	3.9
<b>PM2,5</b>	g/GJ	3.0
<b>BC</b>	g/GJ	1.7
<b>NH3</b>	g/GJ	0.1
<b>As</b>	mg/GJ	0.03
<b>Cd</b>	mg/GJ	0.006
<b>Cu</b>	mg/GJ	0.22
<b>Cr</b>	mg/GJ	0.2
<b>Ni</b>	mg/GJ	0.008
<b>Pb</b>	mg/GJ	0.08
<b>Sé</b>	mg/GJ	0.11
<b>Zn</b>	mg/GJ	29
<b>Hg</b>	mg/GJ	0.12
<b>Dio</b>	ng/GJ	1.4
<b>HAP</b>	mg/GJ	0.0201

### **Off-road sector (category 1A4bii and 1A4cii)**

Category 1A4cii is a key category of NO<sub>x</sub> in terms of emissions level and trend and of NMVOC emissions in terms of emissions level.

The off-road emissions are calculated for the 3 regions by the mathematical model OFFREM (Off-road emission model). Emissions are calculated for machinery used in defence (category 1A5b), harbours, airports and trans-shipment companies (category 1A3eii), in households (category 1A4bii), in agriculture, forestry and green area (category 1A4cii). Exhaust emissions as well as non-exhaust emissions are calculated.

Activity data as input for the model are statistics from harbours and airports, information about households, data on sales of machinery.

For the calculation of energy use and emissions two groups can be divided: off-road machinery and off-road vehicles. Examples of off-road machinery are fork lifts, scissor lifts, lawn mowers. For these machinery the model generates activity data in kWh and methodology of TREMOD is used. Examples of off-road vehicles are luggage carts, quads, sweepers. For these vehicles the model generates activity data in km and aggregated data from MIMOSA model is used (COPERT based model to calculate emissions from road transport).

A complete detailed description about the methodology used can be found in annex 3 of the National Inventory Report (NIR) 2013 where the Quality Management System of the (greenhouse) gas inventory in the Flemish region is described. In the technical procedure of the quality management system VMM/EIL/GP/5.003 'Procedure for the main process: setting up the greenhouse gas emission inventory' this methodology is recorded in annex 7.3.17. with the data acquisition plan for the off-road sector in the Flemish region, which is also used for the emission reporting under CLRTAP.

From the reporting 2015 on, off-road emissions originating from agriculture (combustion emissions from tractors) are taken from OFFREM, as well as off-road emissions in forestry and green area and reported in the category 1A4cii. The agricultural emissions are calculated for arable farming, remaining crops, pasture, intensive livestock and soil-bound agriculture. Emission factors from TREMOD model are used for NO<sub>x</sub>, CO, NMVOC and TSP. NH<sub>3</sub> emission factors are EMEP/EEA Guidebook (Table 3-49).

Table 3-49 Emission factors for the sector 1A4c Agriculture (tractors) in the Flemish region.

			NO <sub>x</sub>	CO	NMVOC	NH <sub>3</sub>	TSP
large farm tractor	<1981	kg/GJ	1,5924137	0,1141090	0,1094474	0,0001826	0,0583325
	1981-1990	kg/GJ	1,1519881	0,1184978	0,0757713	0,0001896	0,0538454
	1991-Stage I	kg/GJ	1,0821256	0,1232377	0,0394011	0,0001972	0,0279996
	Stage I	kg/GJ	0,7342995	0,0739426	0,0236406	0,0001972	0,0139998
	Stage II	kg/GJ	0,5024155	0,0739426	0,0236406	0,0001972	0,0069999
	Stage IIIA	kg/GJ	0,3188406	0,0739426	0,0236406	0,0001972	0,0069999
	Stage IIIB	kg/GJ	0,1739130	0,0739426	0,0133964	0,0001972	0,0014000
medium sized farm tractor	<1981	kg/GJ	0,9057971	0,2200673	0,1407181	0,0001761	0,0874988
	1981-1990	kg/GJ	1,0635230	0,1977321	0,1176151	0,0001839	0,0652976
	1991-Stage I	kg/GJ	1,2598276	0,1691498	0,0927084	0,0001933	0,0274506
	Stage I	kg/GJ	0,7672634	0,0724928	0,0309028	0,0001933	0,0137253
	Stage II	kg/GJ	0,4925642	0,0724928	0,0231771	0,0001933	0,0137253
	Stage IIIA	kg/GJ	0,3125888	0,0724928	0,0231771	0,0001933	0,0137253
	Stage IIIB	kg/GJ	0,2841716	0,0724928	0,0131337	0,0001933	0,0013725

small farm tractor	<1981	kg/GJ	0,6413460	0,2549746	0,1630389	0,0001700	0,1086192
	1981-1990	kg/GJ	0,7553799	0,2375127	0,1432766	0,0001793	0,0763626
	1991-Stage I	kg/GJ	1,0683761	0,2132960	0,1136569	0,0001896	0,0538454
	Stage I	kg/GJ	0,7153475	0,1042781	0,0454628	0,0001896	0,0269227
	Stage II	kg/GJ	0,5109625	0,1042781	0,0303085	0,0001896	0,0134613
	Stage IIIA	kg/GJ	0,3530286	0,1042781	0,0303085	0,0001896	0,0134613
	Stage IIIB	kg/GJ	0,2787068	0,1042781	0,0128811	0,0001896	0,0013461

In Wallonia, emissions from the combustion emissions in the agricultural sector and emissions from farming vehicles are calculated by using the energy use (Energy Balance for Wallonia) and emission factors of the EMEP/EEA guidebook.

The sector 1A4ciii contains the emissions of Agriculture/Forestry/Fishing (national fishing). The activity data (energy consumption data) of the sectors 1A4ciii are taken from the regional energy balances. The emission factors to calculate the emissions for the sector 1A4ciii are given in Table 3-50.

Table 3-50 Emission factors for the sector 1A4ciii, used in Flanders

		diesel oil	
	unit		source
SO <sub>2</sub>	kg/TJ	92	Idem agriculture
Nox	kg/TJ	1200	revised IPCC 1996 Guidelines
NM <sub>VOC</sub>	kg/TJ	200	revised IPCC 1996 Guidelines
CO	kg/TJ	1000	revised IPCC 1996 Guidelines
TSP	kg/TJ	306	VITO
PM <sub>10</sub>	kg/TJ	290,7	VITO
PM <sub>2,5</sub>	kg/TJ	275,400	VITO
Pb	kg/TJ	0,001	

### 3.6. Other (category 1A5a and 1A5b)

In this section the emissions originating from the military transport and off-road emissions of machinery used in defence are included (category 1A5b).

In Wallonia, the Walloon Energy Balance contains the fuel used by military aviation and the emission factors are those described in table 8.8 of the EMEP/EEA guidebook 2009 by using the dutch emission factors (nature of flight: average).

In Flanders, the emissions of military aviation are estimated based on plane movements or fuel use and emission factors. Data on air plane movements of military aviation (airports of Sint-Truiden, Goetsenhoven, Koksijde, Kleine-Brogel and Brasschaat) are reported by the General Staff of the Belgian Airforce, but are not available before 1993 and after 1998. The number of military flights in Melsbroek is reported by Brussels Airport. The Flemish Energy Balance contains the fuel use by military aviation. Based on the evolution of the fuel use, the time series 1999-2013 is calculated.

This section contains also the off-road emissions for machinery used in defence. The emissions are calculated for the 3 regions by the mathematical model OFFREM (Off-road emission model). Exhaust emissions as well as non-exhaust emissions are calculated.

The emissions of category 1A5a are supposed to be included in the sectors 1A1 to 1A4 and 1A5b.

### **3.7. Fugitive emissions from fuels (category 1B1 and 1B2)**

#### **3.7.1. Solid fuel transformation (category 1B1b)**

Emissions during the coke production are caused by the loading of the coal into the ovens, the oven/door leakage during the coking period and by extracting the coke from the ovens.

Activity data (tons of cokes) are delivered by the corresponding industry.

In Wallonia, the emissions factors are the following (ULg 1998)(Table 3-51) :

Table 3-51 Emission factors for the fugitive emissions in Walloon cokeries

	EF	UNIT
SOx	21	g/ Mg PRODUCT
NOx	480	g/ Mg PRODUCT
NMVOC	893	g/ Mg PRODUCT
CO	950	g/ Mg PRODUCT
NH3	138	g/ Mg PRODUCT
TSP	1600	g/ Mg PRODUCT
PM2.5	240	g/ Mg PRODUCT
PM10	560	g/ Mg PRODUCT
As	49	mg/Mg PRODUCT
Cd	123	mg/Mg PRODUCT
Cr	418	mg/Mg PRODUCT
Cu	222	mg/Mg PRODUCT
Hg	30	mg/Mg PRODUCT
Ni	160	mg/Mg PRODUCT
Pb	542	mg/Mg PRODUCT
Se	6	mg/Mg PRODUCT
Zn	542	mg/Mg PRODUCT
DIOXINS	300	ng/Mg PRODUCT
PAH	10000	mg/Mg PRODUCT

In Brussels (plant closed in 1993), the emission factors come from the Guidebook 2013; they are 0.8 g SO<sub>2</sub>/t and 0.9g NO<sub>x</sub>/t. In Flanders no fugitive SO<sub>2</sub> and NO<sub>x</sub> emissions are estimated.

#### **3.7.2. Fugitive emissions from oil (category 1B2a)**

This category includes fugitive emissions from storage and handling in the refinery sector and refinery processes (1B2aiv) as well as emissions originating from petrol stations (1B2av).

Category 1B2aiv is a key category of NMVOC and Pb emissions in terms of emissions level and trend and of Cd, Hg, As and Ni in terms of emissions level.

Category 1B2av is a key category of NMVOC in terms of emissions trend.

Petroleum refineries are all situated in Flanders. Estimation of the emissions from the sector petroleum refining is generally provided by the companies based on monitoring results or emission

factors. Whereas no distinction between fugitive and combustion emissions is possible, emissions of sector 1B2aiv are allocated in 1A1b.

In Wallonia, the fugitive emissions during the distribution of oil products are coming from the transport and storage of gasoline and also from the service stations.

Concerning the service stations, the activity data is the amount of gasoline in the road transport sector in the Walloon energy balance and the emission factor is 2,88 kg/tonne gasoline (Econotec 1998). For the transport and storage, the activity data was estimated via an inquiry in 1996 and recalculated with the annual consumption each year. The emission factor is 0,15 kg/tonne coming from the following legislation : 23 mai 1996 - Arrêté du Gouvernement wallon portant modification du Règlement général pour la protection du travail, en ce qui concerne les dépôts de liquides inflammables, visant à limiter les émissions de composés organiques volatils lors du stockage de l'essence et de sa distribution des terminaux aux stations-service.

In Flanders, for the calculation of NMVOC emissions from gasoline distribution at service stations activity data (amount delivered gasoline) originate from the Belgian Petroleum Federation ([www.petrolfed.be](http://www.petrolfed.be)). Gasoline is distributed for 95% at public service stations and 5% at private, small, stations. The assumption is made that all public service stations are equipped with stage II vapor recovery systems and private stations with stage I vapor recovery systems. The emission factors used are 0.510 g NMVOC/L for stage II systems and 1.3 g NMVOC/L for stations equipped with stage I systems. The factors originate from the BREF 'Best Available Techniques for service stations' (Meulepas P., 1999).

### 3.7.3. Natural gas (category 1B2b)

Category 1B2b is a key category of NMVOC emissions in terms of emissions level.

In the category 1B2b, the fugitive emissions from all transmission, distribution and transport activities of natural gas in Belgium are allocated.

The activity data reported in the category 1B2b is the annual total natural gas amount consumed in Belgium. These activity data originate from SYNERGRID, the federation of the grid operators of gas and electricity in Belgium.

All transmission, distribution and transport activities of gas in Belgium are allocated in this category 1B2b.

The methodology to calculate the emissions of NMVOC originating from the gas distribution (category 1B2biv) is completely harmonised for all the regions in Belgium since the submission in 2004. All information is reported by SYNERGRID, the federation of the grid operators of gas and electricity in Belgium. These emissions are determined on the basis of the length of gas distribution pipelines. The lengths of the main pipelines (exclusive additional, service pipelines which are pipelines going to households) per public utility board are available. The number of additional service pipelines in Flanders is estimated at 1 820 000 for the year 2013. In Wallonia, the number of additional pipelines is estimated at 195 000 for the year 2008. The length per additional pipeline is 5 m in the Flemish and the Walloon region. In Brussels, the number of pipelines is estimated at 186 500 for the year 2006 and is relatively stable for the following years (186 565 in 2010). The average length per pipeline is 3 m because of the urban environment. Depending on the material of the pipeline different emission factors are used. These emission factors are based on measurements carried out. In particular 869, 7865, 869 and 95 m<sup>3</sup>/y/km for respectively steel, pig iron, fibre cement and synthetic material. The density of NMVOC is 1,42 kg/m<sup>3</sup>. The NMVOC content of natural gas distributed is 8 %.

For each material the length of the pipelines is multiplied with the corresponding emission factor. This results in the total natural gas emission in m<sup>3</sup> per year. Multiplying this figure by the NMVOC content

and the density of NMVOC, the diffuse NMVOC emission originating from gas distribution in Belgium is obtained.

Emissions of NMVOC (category 1B2biii, transmission) originating from the storage and transport of natural gas in Belgium are calculated and added to the inventory since the 2006 submission.

These emissions are estimated on the basis of measurements and calculations (taken into account pressure, distance, volume) carried out. All necessary interventions in case of problems are known and the amounts of gas blown-off are registered as accurate as possible. All information is obtained from Fluxys, the independent operator of the gas network in Belgium.

### **3.8. Recalculations and planned improvements**

#### **Recalculations**

- Until the last submission, Wallonia used the database Collector to manage the inventory data's (one collector database by year). But this year, a new air emission software (WAPI) was developed. This software allows the seeing of all data's of a plant or an area source on the complete time series and avoids mistakes during recalculations (jump, zero,...). Some mistakes were corrected as bad values for the Ni and Hg emission factors of heavy fuels in some industrial sectors in 1990, PAH emission factors for the wood boilers in 1A1a, errors of units for some emission factors or direct emissions,...). This new software is also used to report some sectors of the GHG inventory.
- The emission factors of the EMEP/EEA emission inventory guidebook 2013 were used in the three regions. This resulted in new emissions for a number of sectors for the years 1990-2013.  
Actualization of the emission factors according to the guidebook EMEP/EEA 2013
  - o in the Walloon inventory, the new EF are used in almost all of the sectors : 1A1a, 1A2, 1A3c, 1A3dii, 1A4ai, 1A4bi and 1A4c
  - o in Brussels inventory, the new EF are used for sectors: 1A1a, 1A2gviii, 1A3c, 1A3dii, 1A4ai, and 1A4bi.
  - o in the Flemish inventory 1A1a, 1A2, 1A3c, 1A3dii, 1A4ai, 1A4bi and 1A4c
- In the sector 1A3ei (pipeline compressors), recalculation of the NO<sub>x</sub> EF for the year 1990 to 2003 with the average EF on the period 2004-2013.
- For some plants in the sectors 1A2, plant specific emissions were available for the year 2005 but weren't yet integrated in the inventory. The emissions were calculated via emission factors. The plant specific emissions were integrated in this submission.
- In the sector 1A2gvii, the fuel consumption in the quarries was misallocated during the years 2002 to 2009. The fuel consumption was allocated in 1A2f and the emissions factors were the emission factors of stationary combustion. This sector has been recalculated with the emission factors of mobile combustion.
- In the Walloon region, for the sector 1A4ai and the sector 1A4bi, there was no estimation of the SO<sub>2</sub> emissions coming from the combustion of natural gas and LPG. The SO<sub>x</sub> EF of gaseous fuels were integrated in this submission.
- From the reporting year 2014 on redistribution of the off-road emissions from category 1A4aii to categories 1A3e (harbours, airports and transshipment companies) and 1A5b (defence) according to the suggestion of the review team of the GES inventory.
- In Brussels, AD for turbojets was reviewed and corrected. Three turbojets are reported between 1990 to 2008 (closure of one of it). From 2009 to 2013, two turbojets are reported.
- Improvement of reporting HAP; with the new EF, some NE for specific HAP were replaced by emissions.

- Replacement of NE with the emissions for sector 1A1c using the EF of the EMEP/EEA 2013 guidebook for: NH<sub>3</sub>, PM<sub>2.5</sub>, PM<sub>10</sub>, TSP, heavy metals, and POPs with the exception of HCB and PCBs (no EF in the guidebook).
- Replacement of NE for NH<sub>3</sub> for stationary sources using the EF used in Wallonia.
- In Flanders the methodology to calculate off-road emissions originating from agriculture is taken from OFFREM instead of the Tier 1 calculation of emissions by AD x emission factor that was used until reporting year 2014.
- Inclusion of the reporting of BC in the 3 region following the amended Gothenburg Protocol.
- The following recalculations have been implemented for the 3 regions for the road transport :
  - Recalculation of all historic years (1990-2013) with
    - New version of COPERT (v11.1)
    - The inclusion of the COPERT option “Lube oil”
    - The inclusion of scaling factors for the calculation of “fuel sold”
  - A new fully harmonised and unique processing & reporting tool (processing of the COPERT results)

### **Improvements**

- The emission factors of the EMEP/EEA emission inventory guidebook 2013 (tier 2) will be used for wood burning in Flanders and Wallonia.
- For some plants in Wallonia, the emission factors are not consistent throughout the time series. Indeed, from 2005, companies must report their emissions and these emissions are included in the inventory but in previous years, emission factors were sometimes used. For the next submission, emission factors will be calculated on the basis of company data (2005-2014) and used on the entire time series 1990-2004.
- Recalculation of the aviation emissions (time series 1990-2013) with a new model in Flanders
- In Flemish Region the EMMOSS model used to calculate the emissions of sea navigation will be revised for dredging, tugs and extraction of sand.
- The following improvements are planned in the near future for the 3 regions for the road transport:
  - Recalculation using
    - COPERT 4 v11.2 (version 11.1 has known errors regarding the emission factors of Euro VI HDV).
    - Updated mobility figures for 2013 (at the time of the inventory calculation, the mobility for 2013 was not yet published, so the mobility figures of 2012 were used)

### **3.9. QA/QC**

All emissions delivered by the plants are validated and verified by a team of people experienced in emission inventories. In addition, each year a trend analysis is carried out for all emissions per industrial plant and sector. If any inconsistencies or problems are detected by the team, the industry involved is contacted. In exceptional cases the inspection services are contacted.

## Chapter 4. Industrial processes (NFR sector 2)

### 4.1. Source category description

The structure of the industrial sector has undergone profound changes over recent decades. The importance of the (heavy) industrial activities gradually decreases in favour of the service sector, transport and trade. The economic core nowadays in Flanders is situated around the harbours, in Brussels the services become more important and in the Walloon region most industry is situated around some cities. The mining industries have disappeared with the closure of the last coal-mines. The metallurgy and textile sectors have been relatively stable, after several waves of closures and restructuring. The economic crisis hit hard from 2008 on with several closures and restructurings. 2011 was a dark year with the closure of two integrated iron and steel plants in the Walloon region. The two other key sectors of industrial activity are the chemical industry and the food processing industry.

In this sector of industrial processes the emissions of industrial activities which are not related to the combustion of fossil fuels are included. The main source of information on the industrial emissions is obtained from the annual industrial reports. To have a total picture of all emissions by industrial activities, also activities with emissions below the threshold are estimated in a collective way, but this forms a minor fraction of the process emissions.

The emissions of NMVOC in Flanders are estimated by using the results of a study started by Ghent University in 1998 and continued by the Flemish Environment Agency (VMM). In Wallonia, the calculation is based on a methodology established by Econotec. In the Brussels region, the emissions are calculated by using different sources: average emission factors, surveys and information collected from the sector. A recent study (2010) has compiled all information available for the category 'Decorative coating application' and 'Domestic Solvent Use'. The results gave a better overview of these categories and a better estimation of activity data and emission factors.

Tables with detailed NMVOC emissions for 2005, 2010 and 2013 and the Tier methods used are provided for the 3 regions in Annex 2.

Belgium does not report activity data in the NRF tables because part of the activity data is confidential. These source categories consist of several sources and the different activity data are sometimes expressed in different units so it is not possible to show aggregated activity data for these categories.

#### **Allocation of emissions**

The industrial processes in Belgium are covered by

- (1) categories 2A1 (cement production), 2A2 (lime production), category 2A5 (quarrying and mining of minerals other than coal, construction and demolition and storage, handling and transport of mineral products) and 2A6 (other mineral products),
- (2) categories 2B1 (ammonia production), 2B2 (nitric acid production), 2B6 (titanium dioxide production) and category 2B10a (other chemical industry), including 2B10b (storage, handling and transport of chemical products),
- (3) categories 2C1 (metal production i.e. iron and steel industry), 2C7c (other metal production) and 2C7d (storage, handling and transport of metal products),

- (4) categories 2D3 (domestic solvent use, road paving with asphalt, coating applications, degreasing, dry cleaning, chemical products, printing and other solvent use),
- (5) Category 2G (other product use),
- (6) category 2H (pulp and paper and food and drink),
- (7) Category 2I (wood processing),
- (8) category 2K (consumption of POPs and heavy metals),
- (9) category 2L (other production, consumption, storage, transportation or handling of bulk products).

## **4.2. Methodological issues**

The main process emissions of SO<sub>2</sub>, NO<sub>x</sub>, NH<sub>3</sub> and CO are calculated in Belgium by using production figures, mainly directly originating from the industrial plant, combined with emission factors presented in reference works like CITEPA, EMEP/EEA handbook, IPCC Guidelines or other specific bibliographies or calculated via measurements carried out by the industrial companies.

In Flanders, there is a different level of data handling in some years (1990-1993, 1995, 1996, 1998, 2000, 2001, 2008-2012) compared to the other years (1994, 1997, 1999, from 2001 to 2007). In the former years emissions are available on installation level (NFR code), whereas in the latter years the emissions are available on a less detailed level (facility/sector level). A partition key based on the most recent detailed data (e.g. for emission data of 1999, the partition used in 1998 is applied for 1999) is used to attribute the emissions to the appropriate NFR code for the year where less detailed emission data are available.

### **4.2.1. Mineral products (category 2A)**

The mineral industry is one of the most important sectors of industrial process emissions in Belgium.

#### **4.2.1.1 Cement production (2A1)**

This source is a key category of SO<sub>x</sub>, Hg, As, Se, HCB and PCB emissions in terms of emissions level and trend and a key category of NO<sub>x</sub>, PM<sub>10</sub>, TSP and Ni emissions in terms of emission trend.

In Belgium, cement production (5 plants) only takes place in Wallonia.

The activity data is the clinker production collected directly from individual plants.

The emissions of all pollutants are estimated by plant-specific emissions (monitoring and calculation by the plant). The emissions are the sum of combustion and process emissions.

Since 2002, the emissions have varied each year and has been calculated directly by the plant for the PRTR purposes.

An average emission factor by plant and by pollutant has been estimated in 2002 and is applied on the whole time-series 1990-2001.

The evolution of the activity data, the NO<sub>x</sub>, SO<sub>x</sub>, PM10 emissions and the implied emission factors are presented in the Table 4-1.

Table 4-1 Cement production in Wallonia.

	1990	1995	2000	2005	2010	2011	2012	2013
Clinker production (kt)	5292	6055	6089	5555	4740	5060	4869	4694
IEF clinker (kg NO <sub>x</sub> /t)	2,78	2,74	2,87	2,57	2,24	2,31	2.16	1.88
NO <sub>x</sub> emissions (kt)	14,7	16,6	17,5	14,3	10,6	11,7	10.5	8.83
IEF clinker (kg SO <sub>x</sub> /t)	0,81	0,78	0,80	0,94	0,99	0,93	0.8	0.68
SO <sub>x</sub> emissions (kt)	4,3	4,7	4,9	5,2	4,7	4,7	3.9	3.2
IEF clinker (kg PM10/t)			0,20	0,06	0,02	0,02	00004	0.0058
PM10 emissions (kt)			1,2	0,33	0,1	0,1	0.02	0.027

#### 4.2.1.2 Lime production (2A2)

This source is a key category of CO emissions in terms of emissions level and trend and a key category of NO<sub>x</sub>, TSP and Hg emissions in terms of emission trend.

Production of lime also occurs only in the Walloon region of Belgium.

The emissions of lime production (category 2A2) are estimated by using plant-specific emission data for all pollutants excepted NH<sub>3</sub>. The NH<sub>3</sub> emission factor is 5,1 g/t (National Pollutant Inventory in Australia). The emissions of this category are the sum of combustion and process emissions. Since 2002 the emissions have varied each year and have been calculated directly by the plant for the PRTR purposes.

An average emission factor by plant and by pollutant has been estimated in 2002 and is applied on the time-series 1990-2001.

The activity data is the lime and dolomite lime production and is collected directly from individual plants. A part of the lime production is coming from the kraft pulping process.

The evolution of the activity data, the PM10 emissions and the implied emission factors is presented in the Table 4-2.

Table 4-2 Lime and dolomite lime production in Wallonia.

	2000	2005	2006	2007	2008	2009	2010	2011	2012	2013
Lime + dolomite lime (kt)	2640	2601	2774	2677	2586	1783	2116	2234	2091	2034

IEF (kg PM10/t)	0,96	0,62	0,80	0,54	0,44	0,38	0,34	0,03	0,01	0.0081
PM10 emissions (kt)	0,36	0,24	0,29	0,20	0,17	0,21	0,16	0,01	0.026	0.01648

#### 4.2.1.3 Quarrying and mining of minerals other than coal (2A5a)

This source is a key category of TSP emissions in terms of emission level and trend and PM2,5 and PM10 emissions in terms of emission level.

The emissions of this category are the sum of the emissions from the quarrying of minerals and the emissions from storage of minerals in the Walloon region.

Estimation of the emissions from storage of minerals was provided by a study on dust (Econotec 2001).

Emissions from the quarrying of minerals are the sum of plant specific emissions. These plants have had to report to PRTR since 2007. From 2000 to 2006, the estimation of the emissions was also provided by the study on dust.

The evolution of the PM10 emissions is presented in the Table 4-3.

Table 4-3 PM10 emissions in 2A7a

	PM10 (tonnes) (2000-2006)	PM10 (tonnes) (2007-)
Storage of mineral products	1957	1957
Quarrying	284	Plant specific emissions

Emissions (kt)	2000-2006	2007	2008	2009	2010	2011	2012	2013
2A7a Quarrying and mining of minerals other than coal	2,23989	2,24019	2,3364	2,2127	216,336	2,238	2,290	2,376

#### 4.2.1.4 Other mineral products (2A6)

This source is a key category of SOx emissions in terms of emission level and trend.

The category includes the emissions of the clay processing industry (bricks, expanded clay, tiles and glazed stoneware pipes), plaster, fibre cement, fluid concrete and asphalt stirring installations. The process emissions of the glass industry (flat and bowed glass, glass fiber, glass wool,...) are included in sector 1A2f.

The emissions are calculated with plant specific emission factors, based on information reported in the environmental annual reports submitted by the operator of the plants or - if this information is not available - on literature data (Schrooten & Van Rompaey, 2002). Emissions of PM10 and PM2,5 are calculated as a fraction of TSP.

## 4.2.2. Chemical industry (category 2B)

### 4.2.2.1 Ammonia production (2B1)

Nowadays there is ammonia production in 2 companies in Belgium.

In Flanders the emissions of SO<sub>2</sub>, NO<sub>x</sub>, CO and NH<sub>3</sub> originating from the production of ammonia are obtained by monitoring results or calculation with plant specific factors.

In the Walloon region, the producer of ammonia provides the annual NO<sub>x</sub> emissions based on their production and on monitoring.

The following chart (fig. 4.1) shows the trend of the ammonia production :




Figure 4.1 Trend of ammonia production.

### 4.2.2.2 Nitric acid production (2B2)

Despite the closure of 2 plants in the Flemish region, in 1995 and in 2000 respectively, the production of nitric acid still increases. NO<sub>x</sub> emissions are also provided by the plant involved and based on measurements. Since 2000 only one plant is involved in this sector. In Flanders the emissions of SO<sub>2</sub>, NH<sub>3</sub> and CO originating from the production of nitric acid are obtained by monitoring results, emissions of NO<sub>x</sub> are calculated with plant specific factors.

The producer of nitric acid in the Walloon region provides the NO<sub>x</sub> emissions based on their production and on monitoring. There are three installations on the plant. There is only one installation with an abatement technology (SCR) installed in 1996 which lead also in a strong increase of the production in 1996.

The following chart (fig. 4.2) shows the trend of the nitric acid production :




Figure 4.2 Trend of nitric acid production.

#### 4.2.2.3 Other chemical industry (2B10a)

This source is a key category of Nox and NMVOC in terms of emission level and trend and of, SOx, TSP, Cd, Hg, Ni and Zn emissions in terms of emission level or trend.

This category involves all the chemical industry in Belgium which produce an environmental report. In the Walloon region, these are in particular the IPPC plants. In Flanders, in addition to the emissions of the chemical plants, also the emissions of the naphtha cracking installation in one refinery is included in this sector. Also the emissions of the category 2B10b (Storage, handling and transport of chemical products) are included.

The emissions under 2B10a Other chemical industry are mostly taken from reports from the industry.

Industrial plants have to report their emissions of air pollutants from the moment they exceed a defined threshold (in tonne/year) via their yearly environmental reporting obligations. The industry also has the obligation to report the methods used to estimate these emissions.

In the Flemish region an important source for the emissions of the chemical industry is the yearly reporting obligation by the industrial companies via the integrated environmental reports. Nearly all emissions are reported this way. More than 90% of the Flemish NMVOC emission is collected this way for the chemical industry.

The other smaller part of the emissions is estimated based on a survey performed by Ecolas authorized by the Environmental Department of the Flemish Government (Bogaert et al, 2004).

### 4.2.3. Metal production (category 2C)

#### 4.2.3.1 Iron and steel production (2C1)

This source is a key category of NO<sub>x</sub>, SO<sub>x</sub>, PM<sub>10</sub>, TSP, Pb, Cd, Hg, As, Cr, Zn, PCDD/F and PCB emissions in terms of emission level and trend and of PM<sub>2.5</sub>, CO, Ni, Cu and PAH in terms of emission level or trend.

In Flanders, the process emissions from iron and steel production are based on monitoring results provided by the companies. There is one integrated steel plant, one plant that produces stainless steel and one that handles molybdenum to be used in the production of stainless steel. All process emissions from sinter production, blast furnaces, rolling mills, steel production and electric arc furnaces are included.

In Flanders, the HCB emissions are calculated based on activity data and emission factors. The activity data are reported by the industrial companies via the integrated environmental reports. The emission factors are listed in Table 4-4.

Table 4-4 Emission factors of HCB for the sector 2C1 in the Flemish region

	Unit	Value	Reference
Ferro - coke	ng/tonne	596	Liu et al (2009)
Ferro - sinter	µg/tonne	32	EMEP/CORINAIR Guidebook (2005)

In Flanders, this activity is not significant for PCB-emissions.

In the Walloon region, iron is produced through the reduction of iron oxides (ore) with metallurgical coke (as the reducing agent) in a blast furnace to produce pig iron. Steel is made from pig iron and/or scrap steel using electric arc or basic oxygen.

The emission estimates in this sub-sector include also emissions from the production of steel in basic oxygen type furnaces but not emissions from the combustion of the fuel.

In Wallonia, in 2013, there were only five electric arc furnaces. The last integrated iron and steel plant closed in 2011. The process emissions from iron and steel production are based on monitoring results provided by the companies.

The emissions from electric arc furnaces include all the emissions (combustion and process).

#### 4.2.3.2 Other metal production (2C7c)

This category is a key category of Hg, As and Zn emissions in terms of emission level and trend and of Pb, Cd and Ni emissions in terms of emission level and includes emissions from the following activities :

- Surface treatment of metals (galvanizing, electroplating,...)
- In Flanders emissions from non-ferro activities are included in this sector.

The process emissions are based on monitoring results or calculations provided by the companies.

#### 4.2.3.3 Storage, handling and transport of metal products (2C7d)

The emissions from handling of metal products in Brussels are based on monitoring provided by the companies.

#### 4.2.4. Solvent and product use

##### 4.2.4.1 Domestic Solvent Use (category 2D3a)

This source is a key category of NMVOC and PAH emissions in terms of emission level and trend.

A recent study (2010) in the Brussels region has compiled all information available for the category 'Decorative coating application' and 'Domestic Solvent Use' ('NMVOC emissions through domestic solvent use and the use of paints in the Brussels Capital Region', Arcadis, 2010). Thanks to this study, the NMVOC emissions of paint application for construction and building and domestic use have been completely revised in 2010.

The activity data is the population. The emission factors for the different product groups (office products, leather and furniture care, cosmetics and personal care, cleaning products, car products, adhesives/DIY – consumer, insecticides & plant protection products) have been determined by the 2010 study of Arcadis for the Brussels region for the year 2008. The emission factors have been slightly adapted for Flanders and Wallonia. For the Flemish, Walloon and Brussels region, the global emission factors are respectively 1,324, 1,412 and 1.219 kg/person for 2008.

According to the study, VOC-contents in household products have not been severely regulated over the past years. There is no legislation that significantly influenced the VOC-contents in cosmetics, cleaning products or other important VOC-containing household products. Evolution is therefore largely depending on activity data and minor VOC-specific changes. Bearing in mind the recent update of the emission registration methodology (and historical recalculations) in the Netherlands, the evolution for the Netherlands has been transferred to Belgium (1990-2008). A similar evolution of activity data can be assumed as it's a neighbouring country and culture and climate closely relate to each other. For the next years (starting 2009-2013), the emission factors can be assumed to remain constant.

##### 4.2.4.2 Road paving with asphalt (2D3b)

NMVOC emissions due to asphalt production in Flanders are allocated in sector 2L.

These emissions are not estimated in the Walloon inventory.

##### 4.2.4.3 Coating Application: Construction, Building and Domestic Use, car repairing, wood, manufacture of automobiles, other industrial and non-industrial application (category 2D3d)

This source is a key category of NMVOC emissions in terms of emission level and trend.

##### 4.2.4.3.1 Construction, building and domestic use

A recent study (2010) in the Brussels region has compiled all information available for the category 'Decorative coating application' and 'Domestic Solvent Use'. Thanks to this study, the NMVOC emissions of paint application for construction and building and domestic use have been completely revised in 2010 and this for the 3 regions in Belgium.

Information is obtained from IVP (Industry of paints, varnishes and inks) on the sales of decorative paint in Belgium, for both water based and solvent based paints. It is assumed that the IVP data represent 85% of the Belgian market. These activity data are confidential.

The key to allocate the Belgian data to each region is calculated using the number of residential and non-residential buildings and the volume of these buildings for construction and building and using the number of households and the expenses for decorative paint per household for the domestic use of paint.

The solvent content of water based and solvent based paints is obtained from CEPE (the European Council of the Paint, Printing Inks and Artists' Colours Industry). The allocation key between Construction and Building and Domestic Use is obtained from RAINS (Regional Air Pollution Information and Simulation model, developed by IIASA).

#### 4.2.4.3.2 Car Repairing

Since the year 2003, information is obtained from DuPont Refinish Belgium on volumes of paints and thinners sold to the car refinishing industry in Belgium (CRB data). It is assumed that the CRB data represent 85% of the Belgian market. The total volume sold to the car refinishing industry in Belgium is confidential.

The key to allocate the Belgian data to each region is calculated on the basis of the number of car refinishing facilities in 2003: 60% for Flanders, 31% for Wallonia, 9% for Brussels.

The solvent content of the different products are available from DuPont Refinish Belgium for the years 2003 and 2007. The solvent content between 2003 and 2007 is assumed to be equivalent to 2003 and the solvent content after 2007 is assumed to be equivalent to 2007.

For Brussels, an emission factor per company has been established. The AD is the number of companies in the region<sup>4</sup>.

#### 4.2.4.3.3 Wood

In the Flemish region an important source for estimating these emissions is the yearly reporting obligation by the industrial companies via the integrated environmental reports. Together with a correction factor the total emission is calculated (De Roo et al., 2009).

In Wallonia, the activity data is calculated on the basis of the paint sales for the wood and wooden products industry in Belgium in 1996 (IVP data). It is assumed that the paint sales for this sector have followed the same evolution as the economic activity since 1996 and that IVP represents 85% of the

---

4

[http://economie.fgov.be/fr/statistiques/chiffres/economie/entreprises/vie\\_entreprises/jaarevolutie/](http://economie.fgov.be/fr/statistiques/chiffres/economie/entreprises/vie_entreprises/jaarevolutie/)

Belgian market. The number of workers in the wood industry is then used as allocation key to calculate the Walloon sales.

The proportion of water based and solvent based paints as well as the solvent content of these paints come from IVP (2001 & 1996): 30% of water based paints, 5% of solvent in water based paint and 40% of solvent in solvent based paint. As the efficiency of the abatement techniques is not known, it is assumed that no abatement technique exists.

#### 4.2.4.3.4 Manufacture of automobiles

In the Flemish and Brussels regions an important source for estimating these emissions is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

In Wallonia, there is no activity for this sector.

#### 4.2.4.3.5 Other Industrial Application

In the Flemish region an important source for estimating the emissions from other metal coating is the yearly reporting obligation by the industrial companies via the integrated environmental reports. Together with a correction factor the total emission is calculated (De Roo et al., 2009).

In Wallonia, part of the emissions of other industrial coating is the yearly reporting obligation by the industrial companies via the integrated environmental reports. The rest of the emissions is estimated. The activity data comes from IVP (Industry of paints, varnishes and inks). An estimation for the sales of paint for industrial applications in Belgium is assumed. According to IVP, the sales of paint have decreased by 20% between 2009 and 2013. The number of workers in the metal fabrication industry is then used as allocation key to calculate the Walloon sales.

The solvent content in the paints comes from IVP. An average of 40% of solvent has been assumed.

In Brussels the source for estimating these emissions is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

#### 4.2.4.3.6 Other Non-Industrial Application

The emissions of road marking are included here. The activity data (paint consumption data) was obtained from UBATc (Belgium's authority offering technical approval of construction materials, products, systems and installers) in 2011: 6000 t of paint (200 t of water based – 5800 t of solvent based).

It is assumed that the water based paints do not contain solvent. The solvent content of the solvent based paints comes from Ökopol (the Institute for Environmental Strategies): 25%.

The NMVOC emissions of road marking for Belgium are 1450 t in 2013 (870 t for Flanders and 580 t for Wallonia).

The emissions of PM<sub>2,5</sub>, PM<sub>10</sub> and TSP, due to smoking of tobacco in Flanders, used to be reported under 7.A, but it was decided to allocate these emissions in the more appropriate sector 2.G, according to the EMEP/EEA Guidebook 2013.

Emissions are calculated by multiplying the tobacco consumption in Flanders (calculated from the Belgian tobacco consumption, taking into account the number of households and the average

spending, [www.statbel.fgov.be](http://www.statbel.fgov.be)) and an emission factor of 40 kg/ton for TSP, PM10 and PM2,5 (TNO, 2001). The methodology is described in detail in a study performed by VITO under the authority of VMM (Schrooten et al., 2002)

#### 4.2.4.3.7 Degreasing (category 2D3e)

The sales figures of methylene chloride, trichloroethylene and perchloroethylene in UEBL (Economic Union of Belgium and Luxembourg) are obtained each year from ECSA (European Chlorinated Solvent Association). The allocation key is assumed to be 97% for Belgium. The split of applications (pharmaceutical industry, paint stripping, adhesives, metal degreasing, dry cleaning...) is also given by ECSA for Benelux (Belgium, Netherlands, Luxembourg) for the 3 chlorinated solvents.

The following allocation keys is used in Flanders:

- monetary value of sales figures for metal degreasing (De Roo et al, 2009);

The following allocation keys is used in Wallonia:

- Workers in the metal fabrication industry for metal degreasing (adjusted annually);

In de Flemish region the methodology for calculating the NMVOC emission of metal degreasing was optimized in a study conducted by the University of Ghent commissioned by VMM [De Roo et al., 2009]. The consumption of chlorinated solvent for metal degreasing is calculated on the basis of data received from ECSA . The consumption of non-chlorinated solvent for metal degreasing is calculated by making assumptions on the share of cleaning products (2011: non- chlorinated solvents 55%; water-based products 30-35%; chlorinated products 10-15%). The consumption figures of solvent are confidential.

The NMVOC emission factor for the activity without the application of an abatement technology is 0,72 t/t. For the different abatement technologies (closed cold cleaner, closed activated carbon filter closed bag system) the degree of implementation, the technical efficiency and the applicability are estimated. This is done for the use of chlorinated and non-chlorinated solvents. (De Roo et al., 2009).

The NMVOC emission for metal degreasing is calculated using the following formula (D'Haene et al., 2002):

$$E_{i,j} = \sum_{t=1}^n \left( A_{i,j} * EF_{i,j} * \gamma_{i,j,t} * (1 - \eta_{i,j,t} * \alpha_{i,j,t}) \right)$$

with	$E_{i,j}$	NMVOC emission for activity i and year j
	$A_{i,j}$	total activity figure for activitiy i (t solvent/year)
	t	abatement technology
	$EF_{i,j}$	NMVOC emission factor of activity i without application of an abatement technology
	$\gamma_{i,j,t}$	degree of implementation of the abatement technology for the activity (-)
	$\eta_{i,j,t}$	technical efficiency of the abatement technology t (-)

$\alpha_{i,j,t}$  applicability of the technology  $t$  = the part of the emission on which the technology can be applied

In Wallonia, part of the emissions of other industrial coating is the yearly reporting obligation by the industrial companies via the integrated environmental reports. The rest of the emissions is estimated. The consumption of chlorinated solvent for metal degreasing is calculated on the basis of data received from ECSA. The consumption of non-chlorinated solvent for metal degreasing is calculated by making assumptions on the type of existing machines (closed machines using chlorinated solvent, opened machines using chlorinated solvent and opened machines using non-chlorinated solvent) and on the solvent recovery of the various types of machines. The ratio between non-chlorinated solvent and chlorinated solvent is then equal 2,76. The consumption figures of solvent are confidential. It is assumed that 90% of the solvent is lost to air and 10% to other media (water, soil).

In Brussels the source for estimating these emissions is the yearly reporting obligation by the industrial companies via the integrated environmental reports. The reports are available from 2003, the years before are considered constant and equal to the first available year.

#### 4.2.4.3.8 Dry Cleaning (category 2D3f)

The sales figures of methylene chloride, trichloroethylene and perchloroethylene in UEBL (Economic Union of Belgium and Luxembourg) are obtained each year from ECSA (European Chlorinated Solvent Association). The allocation key is assumed to be 97% for Belgium. The split of applications (pharmaceutical industry, paint stripping, adhesives, metal degreasing, dry cleaning...) is also given by ECSA for Benelux (Belgium, Netherland, Luxembourg) for the 3 chlorinated solvents.

The following allocation keys is used in Flanders:

- numbers of dry cleaning companies for dry cleaning (Federation of Belgian textile care; adjusted annually)

The following allocation keys is used in Wallonia:

- Population for dry cleaning (adjusted annually);

In the Flemish region the consumption of chlorinated solvent (PER or perchloroethylene) for dry cleaning is calculated on the basis of data received from ECSA. The consumption of hydrocarbon for dry cleaning is calculated by assuming that hydrocarbons are used in 12% of the dry cleaning machines and that 50% less hydrocarbon is used per kilogram of textiles. The amounts of PER-containing waste and hydrocarbon-containing waste collected from dry cleaning activities in Flanders and the share of PER and hydrocarbon in the waste are obtained from SITA Recyper (Belgian waste management, subsidiary of Suez Environnement). These amounts of products are recycled and not emitted into the air.

The total emission of NMVOC is obtained by deducting the quantities of PER and hydrocarbon in the waste from the consumption of PER and hydrocarbon.

In Wallonia, the consumption of chlorinated solvent (perchloroethylene) for dry cleaning is calculated on the basis of data received from ECSA. The consumption of non-chlorinated solvent for dry cleaning is calculated by assuming that the chlorinated solvents represent 90% of the total consumption. The consumption figures of solvent are confidential. It is assumed that 90% of the solvent is lost to air and 10% to other media (water, soil).

In Brussels, dry cleaning emissions are calculated on the basis of the emission factor of 5.31 g NMVOC/hab determined in 2002, combined with the evolution of the total population.

#### 4.2.4.3.9 Other Industrial Cleaning (category 2D3e)

In Wallonia, the consumption of chlorinated solvent for other industrial cleaning is calculated on the basis of data received from ECSA. The consumption of non-chlorinated solvent is not determined for this sector. The consumption figures of solvent are confidential.

The following allocation keys is used in Wallonia:

- Workers in industry for the other applications (adjusted annually).

It is assumed that 90% of the solvent is lost to air and 10% to other media (water, soil).

#### 4.2.4.4 Chemical Products, Manufacture and Processing (NFR 2D3g)

The category 2D3g is a key category of NMVOC emissions in terms of emission level.

##### 4.2.4.4.1 Polyester Processing

In the Flemish region an important source for the emissions of polyurethane processing is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

In Wallonia, the activity data used to come from Reinforplast (Association of Belgian Manufacturers of Reinforced Plastics/Composites). No statistics of production are available. In 1996, Reinforplast estimated the Belgian production based on information coming from the fiberglass suppliers). A small half of the producers were located in Wallonia but most of the big producers were located in Flanders. In terms of production, this represented 75% for Flanders and 25% for Wallonia.

In 2001 contact was made with Reinforplast, an estimation was made on the Walloon production based on an assumption of the Belgian production and assuming 65% in Flanders and 35% in Wallonia..

In 2010 contact was made with Federplast, the Association of Belgian Manufacturers of Articles in Plastics and Elastomers within Agoria (Belgian Federation for the Technology Industry) and Essenscia (Belgian Federation for Chemistry and Life Sciences Industries). No production figures are available even at European level. There are approximately 400 composites manufacturers in Belgium; half of them are located in Wallonia but all of relatively small size. At European level, the sector is growing but in Belgium it decreases. In the past, 75% of the production was attributed to Flanders but, in 2010, this proportion has decreased to 60% because many big producers have disappeared in Flanders.

In 1996, according to the fiberglass suppliers, the proportion of the different application techniques was: 42% for contact, 12% for filament winding, 35% for projection and 11% for other techniques. The styrene content in the resin depends on the process and can vary between 30 and 50%. A styrene content of 40% was assumed. For each application technique, the following styrene emissions (in % of the styrene used) were assumed: 3.2% for contact (1% in case of LSE resin), 4% for filament winding (2% for LSE resin), 8,3% for projection (3% in case of LSE resin) and 1,3% for other techniques (0,6% for LSE resin). In 1996, the proportion of low styrene emission resin was approximately 20% but this proportion has increased since then and is estimated to 40% in 2010. It is assumed that no abatement techniques are applied.

Emissions from the cleaning agents must be added to the styrene emissions. It is assumed that those emissions represent 40% of the total emissions for the composite production.

#### 4.2.4.4.2 Polyvinyl Chloride Processing (PVC)

For the Flemish region, the NMVOC-emissions are included in other categories.

In Wallonia, the activity data for this sector is the consumption of plastic for the manufacture of electric cables. In 1996, this consumption was coming from the CRIF (Centre de Recherche scientifiques et techniques de l'Industrie des Fabrications métalliques – became SIRRIS in 2007). Only part of the plastic consumption must be attributed to flexible PVC but there is a lack of information so it is considered that 100% of the plastic used is PVC.

In 2012 contact was made with SIRRIS (Collective Centre of the Belgian Technology Industry) to actualize the activity data. Unfortunately, no current global activity data is available. The plastic consumption in 2010 is assumed to be identical to 1996. This assumption is conservative because the plastic activities have decreased since 1996.

The proportion of plasticizers (phthalates as DOP and DEHP) in the resin can vary from 20% to 60% depending on the applications. A proportion of 40% of plasticizers is assumed. The emissions of plasticizers are assumed to be 2,5% of their consumption.

#### 4.2.4.4.3 Polyurethane Processing

In the Flemish region an important source for the emissions of polyurethane processing is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

In Wallonia, the activity data for this sector is the production of polyurethane foam. In 1996, the PUR production in Wallonia was estimated on the basis of the following information/assumptions:

- Belgian production of cellular products (INS 1993) ;
- No Belgian production figures for PUR exists, an assumption was made;
- Other plastics can be made cellular (PP, PE), an assumption for the Belgian production was made
- 15% of PUR is produced in Wallonia (based on the number of producers in 1996).

In 2012 contact was made with SIRRIS (Collective Centre of the Belgian Technology Industry) to actualize the activity data. Unfortunately, no current global activity data is available. The PUR production in 2010 is assumed to be identical to 1996. This assumption is conservative because the plastic activities have decreased since 1996.

The emission factor is 15 kg VOC/t PUR foam (Cahier sectoriel 'Technologies et Environnement', volume « Les thermoplastiques », Ministère de la Région wallonne, DGTRE, 1996).

#### 4.2.4.4.4 Polystyrene foam processing

In the Flemish region an important source for the emissions of polystyrene foam processing is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

In Wallonia, the activity data for this sector is the production of expanded polystyrene. In 1996 the EPS production in Wallonia was estimated on the basis of the following information/assumptions:

- Belgian production of expanded polystyrene = 20 kT (CRIF 1996);
- 40% of EPS is produced in Wallonia (based on the number of producers in 1996).

In 2012 contact was made with SIRRIS (Collective Centre of the Belgian Technology Industry) to actualize the activity data. Unfortunately no current global activity data is available. The PUR production in 2010 is assumed to be identical to 1996. This assumption is conservative because the plastic activities have decreased since 1996.

The emission factor is 33,3 kg VOC/t PUR foam (Cahier sectoriel 'Technologies et Environnement', volume « Les thermoplastiques », Ministère de la Région wallonne, DGTRE, 1996).

#### 4.2.4.4.5 Rubber processing

In the Flemish region an important source for the emissions of the rubber processing is the yearly reporting obligation by the industrial companies via the integrated environmental reports. More than 80% of the Flemish NMVOC emissions is collected this way for the rubber processing activities.

The other smaller part of the emissions is calculated based on:

- the number of tires produced in Belgium (the Federal Public Service for Economy, General Directorate for Statistics and Information on Economy)
- emission factor 100 g/tyre (D'Haene et al., 2002)
- the key to allocate the Belgian data to the Flemish region is calculated on the basis of the number of rubber processing companies (91% in 2013).

In Wallonia, this activity does not exist anymore. From 1990 to 2001, there was only one tyre manufacturer. The NMVOC emissions of this manufacturer have decreased in 1996 due to a modification in the process, In 2001, the company has closed.

#### 4.2.4.4.6 Pharmaceutical Products Manufacturing

In the Flemish region the emissions of the pharmaceutical products manufacturing includes the emissions of the synthesis and the formulation. For the synthesis an important source for the emissions of the pharmaceutical products manufacturers is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

The other smaller part of the emissions caused by the formulation is based on a survey performed by Ecolas authorized by the Environmental Department of the Flemish Government (Bogaert et al, 2004).

In Wallonia, the emissions are directly obtained from the pharmaceutical products manufacturers. The NMVOC emissions for Wallonia include the emissions of the cleaning agents.

#### 4.2.4.4.7 Coating Manufacture: Paint

In the Flemish region an important source for the emissions of the coating manufacturing is the yearly reporting obligation by the industrial companies via the integrated environmental reports. Since 2007 even 100% of the Flemish NMVOC emissions is collected this way for the coating manufacturing.

For the period 1990-2006, the other smaller part of the emissions is estimated based on the total solvent content in produced coatings in Flanders minus the solvent content of the Flemish companies with a yearly environmental report. An estimation is necessary for those coating manufacturers who have no obligation to report their emissions.

The activity data is the total Flemish paint production. These figures are confidential.

The estimation is based on production figures of decorative and industrial coatings (source IVP, Industry of paints, varnishes and inks). The part of the production allocated to Flanders is 79,4%.

The average solvent content in the paint is calculated on the basis of the solvent content in the coatings: 10% in water based decorative and industrial coatings; 40% in solvent based decorative coatings; 50% in solvent based industrial coatings (source IVP)

An emission factor of 4,4% of the solvent consumption is assumed (IVP).

In Wallonia, the activity data is the Walloon paint production. These figures are confidential. This data is calculated on the basis of the following data:

- Belgian sales of decorative paint (adjusted each year on the basis of IVP data ;
- Assumption on the proportion of the decorative paint exportations (contact with IVP, 2009): 90% sold in Belgium – 10% exported ;
- Belgian sales for the car repairing sector (adjusted each year on the basis of CRB data;
- Assumption on the proportion of car refinish paint exportations (contact with IVP, 2009): 50% sold – 50% exported;
- Assumption on the Belgian sales of paint for other industrial applications (contact with IVP, 2009 and 2013;
- Assumption on the Belgian production of paint for other industrial applications (contact with IVP, 2009);
- Assumption on the part of the production that must be allocated to Wallonia: 20%.

The average solvent content in the paint is calculated on the basis of the solvent content in the decorative paints (adjusted each year - 9% in 2010), car refinish paints (adjusted each year - 35% in 2010) and industrial paints (40% - estimation of IVP in 2013). The average solvent content in the paint is 30% in 2010. An emission factor of 4,4% of the solvent consumption is assumed (IVP).

#### 4.2.4.4.8 Inks Manufacturing

In the Flemish region an important source for the emissions of the inks manufacturing is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

In Wallonia, before 2002, IVP data were used to estimate the NMVOC emissions. Since 2002, the data are obtained directly from the inks producers. Most of the producers are located in Flanders. There are few producers in Wallonia. The producers calculate their emissions on the basis of a solvent management plan. The activity data is the solvent consumption. The implied emission factor depends on the type of ink produced and the use of an abatement technique. (It can vary from 1,5 kg NMVOC/T solvent used to 50 kg NMVOC/T solvent used).

#### 4.2.4.4.9 Glues Manufacturing

For the Flemish region the methodology has been optimized in a study performed by the University of Ghent authorized by the VMM (De Roo et al., 2009). The activity data for Flanders are confidential and obtained from the Federal Public Service for Economy (General Directorate for Statistics and Information on Economy). The share of solvent based glues is 7% of the total production figure of glues. The solvent content of the glues is 60%. The emission factor is 1,25%.

The emission of one company is not included in the activity figure and is extracted from the integrated environmental report. The production of urea formaldehyde (UF) based glues is also not included in the activity figure. In Flanders two companies produce UF glues. An emission figure for each

company is taken into account, based on the integrated environmental report or a survey performed by the VITO (Lodewijks et al., 2003).

In Wallonia, this activity is not significant. The NMVOC emissions of the few producers are reported under category 2B10a.

#### 4.2.4.4.10 Adhesive and Magnetic Tapes, Film and Photographs Manufacturing

In Wallonia, the NMVOC emissions are obtained directly from the only adhesives producer on the basis of a solvent management plan .

#### 4.2.4.4.11 Leather tanning

For the Flemish region this activity is not significant. The NMVOC emissions are not estimated.

In Wallonia, the NMVOC emissions are obtained directly from the 2 tanneries. There is no abatement technique. The emissions are equal to the solvent consumptions.

#### 4.2.4.4.12 Other Chemical Product Manufacturing or Processing

For the Flemish region no other NMVOC emissions are allocated here.

In Wallonia, most of the NMVOC emissions of other chemical product manufacturing or processing are reported under category 2.B.10.a. The emissions of only one producer are allocated here. The NMVOC emissions are calculated on the basis of a solvent management plan.

#### 4.2.4.5 Printing (category 2D3h)

This source is a key category of NMVOC emissions in terms of emission level and trend. In the Flemish region an important source for the emissions of the printing industry is the yearly reporting obligation by the industrial companies via the integrated environmental reports. More than 80% of the Flemish NMVOC emissions is collected this way for the printing industry.

The other smaller part of the emissions is estimated. An estimation is necessary for those sheet-fed offset companies who have no obligation to report their emissions. The estimation is based on a survey carried out by FETRA (the Belgian federation of paper- and board manufacturing industries) and Febelgra (the Belgian professional representative federation of the graphic industry).

In Wallonia, part of the emissions of the printing industry is the yearly reporting obligation by the industrial companies via the integrated environmental reports. The rest of the emissions is estimated. The activity data is the Walloon ink consumption. The figures of inks sales in Belgium and Luxembourg are obtained from IVP. It is assumed that 97% of the sales can be attributed to Belgium. The part to be attributed to Wallonia is then calculated on the basis of the number of workers in the printing industry.

The proportion of each printing techniques used to come from IVP but since 2007 these data could not be actualized. The average solvent content of the ink for each printing technique were obtained by IVP in 2000 and have been partially actualized in 2009 on the basis of the Guidance on VOC Substitution and Reduction for Activities Covered by the VOC Solvents Emissions Directive (March 2009, Final Report, European Commission – DG Environment). On the basis of these data, the

Walloon solvent consumption can be calculated. The abatement efficiency for each printing technique also comes from the Guidance on VOC (see reference above). The emission factors with and without abatement are obtained from an EGTEI document (100% of solvent emitted without abatement – 5% with abatement). On the basis of these data, the Walloon emissions of the solvents in inks can be calculated.

In Brussels, the emissions from printing are estimated on the basis of NMVOC balances (big establishments) and estimations from average emissions factors (small businesses).

#### 4.2.4.6 Application of Glues and Adhesives (category 2G/2D3i)

In the Flemish region (2D3i) the following activities are included:

- bonding (gluing) of wood: the emissions of the chipboard companies are extracted from the integrated environmental reports.
- bonding (gluing) of synthetic material: an important source for estimating the emissions is the yearly reporting obligation by the industrial companies via the integrated environmental reports. Together with a correction factor the total emission is calculated (De Roo et al., 2009).

In Wallonia, (2G), the activity data is the glues and adhesives sales. This data is obtained from a study of DETIC (Belgian-Luxembourg Association of producers and distributors of soaps, cosmetics, detergents, cleaning products, hygiene and toiletries, glues, and related products) in 2002. As most of the sales are attributed to the construction sector, the part to be attributed to Wallonia is calculated on the basis of the population figures. According to DETIC, their members represent 70% of the Belgian market for glues and adhesives. On the basis of these data, the Walloon consumption of solvent based glues and adhesives is estimated excluding domestic use) in 2002. Unfortunately, this data is not available annually so the same figure is used from 2002 to 2010.

In addition to providing the sales of solvent based glues and adhesives for different applications, the DETIC study also provided the average solvent content of the glues and adhesives for each application. According to DETIC, the solvents in the solvent based glues and adhesives represent 90% of the total solvents (in both solvent and water based glues and adhesives). On the basis of these data, the Walloon consumption of solvent in glues and adhesives has been estimated for 2002 (excluding domestic use). As the data cannot be adjusted annually, the same figure is used from 2002 to 2013.

It is assumed that the emissions equal the consumptions (emission factor of 1 kg/kg).

In order to achieve alignment, the three regions will allocate these NMVOC-emissions in sector 2D3i in the next submission.

#### 4.2.4.7 Preservation of Wood (category 2G/2D3i)

In the Flemish region (2D3i) the emissions are caused by the use of creosote and solvent based products. Creosote B is gradually replaced by creosote C and solvent based products are gradually replaced by water based products. The emissions caused by the use of creosote are collected by a yearly survey. In 2013 there is only one user of creosote in Flanders with negligible emissions. The emissions caused by the use of solvent based products are extracted from the Flemish BAT (Best Available Technology) study Wood manufacturing industry (Polders et al., 2011).

In Wallonia, (2G), to estimate the emissions from 1990 to 1999, assumptions have been made on the consumptions of wood impregnation products (ECONOTEC, 2000). A VOC content of 27% has been

assumed. This corresponds to the VOC content of creosote B at 40°C. It was assumed that the emissions equal the consumptions (emission factor of 1 kg/kg). Since 2000, as there is a lack of global information on the volume of impregnated wood and the products consumption, contact has been established with the main producers to estimate their emissions on the basis of the product consumption, the VOC content of the different products (depending on the condition of use), the process and the abatement techniques used. Creosote B is gradually replaced by creosote C and solvent based products are gradually replaced by water based products, so the global NMVOC emissions tend to decrease over time.

In order to achieve alignment, the three regions will allocate these NMVOC-emissions in sector 2D3i in the next submission.

#### 4.2.4.8 Fat, edible and no-edible oil extraction (category 2D3i)

In the Flemish region an important source for the emissions of oil extraction is the yearly reporting obligation by the industrial companies via the integrated environmental reports.

In Wallonia, this activity is not significant. The NMVOC emissions are not estimated. The emissions of one producer are reported under category 2D3g.

### 4.2.5. Pulp and paper (2H1)

This category includes process emissions from the following activities :

- Paper pulp plant (kraft process) (NMVOC emissions, no relevant NMVOC emissions in Flanders)
- Graphic sector
- Publishers/press

The process emissions are based on monitoring results provided by the companies.

### 4.2.6. Food and drink (2H2)

This source is a key category of NMVOC emissions in terms of emission level and trend.

This category includes emissions from the following activities :

- Bread production
- Production of beer and other drinks (including milk)
- Abattoirs
- Oil production for consumption
- Production of starch
- Industrial fish smoking (PM)
- Meat cooking and barbecue (PM)
- Production of all other food

In Flanders, the process emissions from food and drink production of NO<sub>x</sub> and CO are based on monitoring results provided by the companies. Dust and NMVOC emissions are calculated based on activity data and emissions factors, given below in table 4.5.

In Wallonia and Brussels, the emissions are calculated with the activity data and the emission factors given in Table 4-5.

Table 4-5 AD and EFs used in 2H2

Type of products	region	Activity data	Emission factor	Reference
Bread	Flanders	111 g/hab.day	4500 g NMVOC/t	AD: Statbel EF: Emep guidebook 2009 Study Van Rompaey (1999)
Bread	Wallonia	125 g/hab.day	4500 g NMVOC/t	AD : Professional union of bakers and Statbel EF : Emep guidebook 2009
Beer	Flanders	74% x Belgium production	0,035 kg NMVOC/hl beer	AD : Beerparadise EF : Emep guidebook 2009
Beer	Wallonia	26 % x Belgium production	0,035 kg NMVOC/hl beer	AD : Beerparadise EF : Emep guidebook 2009
Bread/cookies	Brussels	112 g/hab.day	4500 g NMVOC/t	AD : Statbel EF : Emep guidebook 2013
Fish smoking	Flanders	Prodcom statistics	0.080 kg TSP/ton	Study Schrooten & Van Rompaey (2002)
Meat cooking	Flanders	76 kg/hab.year	1.30 kg TSP/ton	Study Schrooten & Van Rompaey (2002)
Barbecue (meat cooking)	Flanders	130 g/hab.year	40 kg TSP/ton	Study Schrooten & Van Rompaey (2002)
Barbecue (charcoal emissions)	Flanders	165 g/hab.year	2.40 kg TSP/ton	Study Schrooten & Van Rompaey (2002)

#### 4.2.7. Consumption of POPs and heavy metals (category 2K)

##### 4.2.7.1 The use of PCB transformers and capacitors

Directive 96/59/EC on the disposal of PCBs and PCTs aims at disposing completely of PCBs and equipment containing PCBs as soon as possible. This Directive sets the requirements for an environmentally sound disposal of PCBs. Member States have to make an inventory of big equipment containing PCBs, have to adopt a plan for disposal of inventoried equipment, and outlines for collection and disposal of non-inventoried equipment (small electrical equipment very often present in

household appliances manufactured before the ban on marketing of PCBs). The PCB Directive further mandates that Member States had to dispose of big equipment (equipment with PCB volumes of more than 5 litres) by the end of 2010 at the latest.

In 2000 the OVAM (Public Waste Agency of Flanders) started a PCB disposal plan for Flanders with a stepwise destruction (based on the year of manufacture) of PCB-containing transformers/capacitors containing more than 1 litre of liquid with more than 0.05% PCBs.

The activity data are obtained from the OVAM:

- the total amount of destroyed and reported transformers and capacitors;
- the amount of liquid volume classified by concentration of PCBs in the liquids.

The emission factors are taken from the EMEP/EEA Emission Inventory Guidebook. Based on the total amount of liquid volumes from the reported transformers and capacitors minus the amount of liquid volumes of the destroyed transformers and capacitors the remaining liquid volume can be calculated. Based on the known PCB content and the emission factors (Table 4-6), the PCB-emissions can be calculated.

Table 4-6 Emission factors of PCB for sector 2K in the Flemish region

	Unit	Value	Reference
PCB transformer	kg/tonne PCB	0.06	EMEP guidebook 2009
PCB capacitor	kg/tonne PCB	1,6	EMEP guidebook 2009

#### 4.2.8. Other production, consumption, storage, transportation or handling of bulk products (category 2L)

This is a key category of HCB emissions in terms of emission level and trend.

In Flanders HCB-emissions are caused by the production of tetrachloromethane, trichloroethylene and tetrachloroethylene. The activity data are obtained from ECSA (European Chlorinated Solvent Association) and are confidential. No activity data are obtained for tetrachloromethane. The emission factors are taken from the EMEP/EEA Emission Inventory Guidebook (Table 4-7).

Table 4-7 Emission factors for the sector 2L in the Flemish region

	Emission factor	Reference
Production of solvents: tetrachloromethane	10 g/ton	EMEP/CORINAIR Guidebook (2005) Sources of HCB
Production of solvents: trichloroethylene	4 g/ton	EMEP/CORINAIR Guidebook (2005) Sources of HCB
Production of solvents: tetrachloroethylene	6 g/ton	EMEP/CORINAIR Guidebook (2005) Sources of HCB

For particulate matter and heavy metals, process emissions originating from the wood, textile, rubber and plastic handling, automobile, electrotechnical industry are allocated in this sector. These emissions are reported by the facilities in the annual industrial reports.

### **4.3. Recalculations**

In the Flemish region the following recalculations were made to optimize the inventory:

- NMVOC: The sectors 2D3g and 2D3h have been updated with the latest environmental reports available. AD for the sector 2D3g have been updated with new available statistics.
- PAH: the sector 2C1 has been updated with the latest environmental reports available.

In Brussels region, the following recalculations have been taken place:

- The AD for bread and cookies production was corrected between 1990-2001 in Brussels database.
- There was a mistake on population data for sectors 2D3a, 2D3d and 2D3f which was corrected.
- The sector 2D3h has been updated with the latest environmental reports available.
- Sector 2D3e is being reported for the first time, information comes from annual environmental reports. The data is available from 2003. The emissions for the period 1990 – 2002 are kept constant and equal to the first available data.
- AD for car repairing (2D3d) and printing (2D3h) has been updated with new available statistics from the year 2012.

In Wallonia, the following recalculations have been performed:

- 2H2 : modification of the NMVOC EF for bread (125 g/hab/year)
- 2D3d Coating applications:
  - o Construction and buildings and domestic use: Emissions of the years 2011 and 2012 have been slightly modified because of an error in the AD figures.
  - o Other industrial paint application: Emissions of 1990 have been modified to reflect the change in other sectors. Emissions from 2007 to 2012 have been modified to take into account the emissions determined on the basis of the yearly integrated environmental reports and the decrease of the global paint consumption between 2009 and 2013.
- 2D3e Degreasing:
  - o Metal degreasing: Emissions from 2010 to 2012 have been modified to take into account the emissions determined on the basis of the yearly integrated environmental reports.
- 2D3g Chemical products:
  - o Paints manufacturing: Emissions from 2010 to 2013 have been modified to reflect the decrease of the global paint consumption between 2009 and 2013.
  - o Pharmaceutical products manufacturing: Emission of 2012 have been slightly actualized on the basis of new data provided by one plant.
- 2D3h Printing:

- Printing industry: Emissions from 1990 to 2012 have been revised in order to ensure a coherence in the methodology for the all time series and to take into account the emissions determined on the basis of the yearly integrated environmental reports.
- 2D3i Other solvent use:
  - Preservation of wood: Emissions from 2010 to 2012 have been actualized on the basis of new activity data provided by one plant.

#### **4.4. QA/QC**

All emissions delivered by the plants are validated and verified by a team of people experienced in emission inventories. In addition, each year a trend analysis is carried out for all emissions per industrial plant and sector. If any inconsistencies or problems are detected by the team, the industry involved is contacted. Numerous contacts take place with the plant operators as well as with the federations involved. In exceptional cases the inspection services are contacted.

#### **4.5. Planned improvements**

In the Flemish region, the following improvements are planned:

- Fine tuning the emissions of polyurethane processing;
- Fine tuning the emissions of polystyrene foam processing;
- Estimation of the emissions of production of solvents: tetrachloromethane.

In Wallonia, the following improvements are planned:

- For some plants, the emission factors are not consistent throughout the time series. Indeed, from 2005, companies must report their emissions and these emissions are included in the inventory but in previous years, emission factors were sometimes used. For the next submission, emission factors will be calculated on the basis of company data (2005-2014) and used on the entire time series 1990-2004.
- The emission factors for PM10 and PM2.5 are not consistent for the time series as since 2005, the lime plants have performed PM10 analyses and have made an estimation of their PM2.5 emissions. Following information's coming from these plants, the size of the dust is high and there is very little fine dust. Before 2005, the proportion between PM10 and TSP was the proportion written in the Emep Guidebook. A recalculation is planned to harmonise the proportion TSP/PM10/PM2.5 with plant data for the entire period.
- Estimation of the NMVOC emissions from the sector road paving with asphalt.
- Revision of the emissions for Wood paint application ;
- Revision of the emissions for non-chlorinated solvents for Metal degreasing, Dry cleaning and Other industrial cleaning;
- Revision of the emissions for Polyester processing, Polyvinylchloride processing, Polyurethane processing, Polystyrene processing;
- Estimation of the missing emissions (NE) for Textile finishing, Glass wool enduction, Mineral wool enduction and Domestic use of pharmaceutical products ;
- Revision of the emissions from key sources in order to move from Tier 1 to Tier 2 methodology when necessary

## Chapter 5. Agriculture (NFR sector 3)

### 5.1. Overview

#### 5.1.1. Allocation of emissions

The agricultural sector includes the emissions originating from animal manure management (NFR sector 3B), the use of synthetic N-fertiliser (NFR sector 3Da1), animal manure applied to soils (3Da2a), urine & dung deposited by grazing animals (NFR sector 3Da3) and from manure processing (NFR sector 3Dd). More detailed information on emissions due to fuel use in the agricultural sector is included in Chapter 3 Energy (3.5). The emissions reported in NFR sector 3 are based on calculations using specific regional information. The categories 3B1a and 3B1b (cattle dairy and non-dairy), 3B3 (swine), 3B4gi and 3B4gii (laying hens and broilers), 3Da1 are key categories for NH<sub>3</sub>, either in terms of emission level, trend or both level and trend. The category 3B3 (swine) is a key category for PM10 in terms of emission level. For TSP, the categories 3B1b, 3B3 and 3B4gii (non-dairy cattle, swine and broilers) are key categories in terms of emission level.

#### 5.1.2. Description of the sector

The land used for agriculture in 2013 in Belgium extends to 1 338 556 hectares. In 2013, the number of agricultural and horticultural businesses amounted to 37 761. This number had dropped by 39% since 2000. The disappearing of small businesses being a general trend in the sector. Additionally in Flanders, this partly can be explained due to the subsidized cut down of the number of cattle. This was in 2001 and 2002 only the case for swine. In 2003 however an extension to bovine and poultry occurred. Nevertheless the land area used for agricultural purposes remained more or less the same during this period. In 2013 Wallonia has 54% of the land used for agriculture, but 66% of agricultural businesses are situated in Flanders. The land area used for farming is on average 25 ha per farm in the Flemish region and 56 ha per farm in the Walloon region.

#### 5.1.3. Climate:

With an average temperature of 10.1°C in 2013 (<http://www.meteo.be/meteo/view/fr/13812254-2013.htm>), Belgium as a whole has a 'cool' climate.

#### 5.1.4. Data sources

The main activity data are the livestock figures, N-excreted and amount of synthetic fertilizer use. 'Statistics Belgium' (Statbel) publishes data on livestock figures yearly in its agricultural census. As the main statistical authority in Belgium, 'Statistics Belgium' is in charge of collecting, processing and disseminating relevant, reliable and commented statistical and economic information. Until 2008, the agricultural census reached 100% of the farms. Since 2008 (with exception of 2010) this inquiry has changed slightly. 75% (before 2008 this was 100%) of all agricultural businesses (including the biggest farms) have to fill in a form each year about the situation on the farm on the 1st of May of that year.

The other 25% is estimated. To come to this 75%/25% ratio, the farms are divided in two groups: 50% contain the biggest farms, the other 50% the smaller farms. The 50% biggest farms have to fill in the form each year. From the other 50% smaller farms, the half has to fill in the form in year x and the other half is estimated. The next year (x+1) the part of small farms that is not contacted in the year x,

is obliged to fill in the form. At this way every two years 100% of the farms are questioned. To be compliant with the European legislation, in the survey 2010 once again 100% of the farms are questioned.

Further details on the agricultural census methodology and QA/QC issues can be found on the Statbel website: <http://economie.fgov.be/en/statistics/surveys-methodology/>.

#### 5.1.4.1 Livestock

The livestock numbers are the primary activity data used in the calculation of NH<sub>3</sub> emissions. The numbers originate from the agricultural census which is available on:

[http://statbel.fgov.be/fr/modules/publications/statistiques/economie/downloads/agriculture\\_-\\_chiffres\\_d\\_agricole\\_de\\_2013.jsp#.VOr1pJ2G\\_wx](http://statbel.fgov.be/fr/modules/publications/statistiques/economie/downloads/agriculture_-_chiffres_d_agricole_de_2013.jsp#.VOr1pJ2G_wx)

These data are available for and used by the two regions, Wallonia and Flanders. Table 5-1 gives an overview of the origin of livestock number in the two regions for the different time periods.

Table 5-1: Origin of the livestock numbers in the two regions

Livestock numbers	Flanders	Wallonia
1990-1999	STATBEL	STATBEL
2000-2013	Manure Bank (VLM)	STATBEL

In Flanders, however, from 2000 on, input data as animal number, N-production e.o. are obtained by the Manure Bank of the Flemish Land Agency (VLM; <http://www.vlm.be/landtuinbouwers/mestbank/aangifte/Pages/default.aspx>). This information is available on the level of the stable as necessary for the NH<sub>3</sub>-model. In 2009, in Flanders, a new model for the calculation of the NH<sub>3</sub> emissions was developed. This model calculates the NH<sub>3</sub> emission in different emission stadia taking into account the manure flow. This is done on the level of the stable. Therefore data (animal number, manure transport, N-excretion) were necessary on this detailed level. These data are inventoried by the Manure Bank from the Flemish Land Agency (VLM). The VLM, a Flemish government agency is, among other things, responsible for the execution of the Flemish Manure Policy. Statbel can provide data on animal number, only on the level of municipality. This is not detailed enough for the NH<sub>3</sub>-model. On the other hand, data from the Manure Bank are only available from 2000 on. To be consistent between different models used (NH<sub>3</sub>, N<sub>2</sub>O, CH<sub>4</sub>) Flanders decided to use the VLM data source for animal number and N-excretion for all models and from 2000 on. From 1990-1999 Flanders uses the Statbel numbers, which also means that NH<sub>3</sub> emissions in this period can only be calculated on the level of the municipality.

It is of course true that the animal number between Statbel and the manure bank is not exact the same. Statbel collects data on the 1<sup>st</sup> of May, which means that farmers give the animal number present at the farm at the 1<sup>st</sup> of May. For the manure bank farmers give the average animal population of the past year. This difference explains differences in animal number between the two data sources. The differences between the data sets do not exceed 10%, which is the uncertainty level for the animal population data from STATBEL.

#### 5.1.4.2 N-excretion factors

For the N-excretion factors of swine and poultry in Flanders, a farmer can choose to use the standard excretion factors (no special effort to reduce N and/or P production) . Or they can choose (or in some cases are obliged) to use the other systems (regressive balance, animal feed covenant, a complete fodder (input-output) balance). These data are obtained by the Manure Bank of the Flemish Land Agency. The N-excretion factors of cattle, sheep, goats, horses, mules and rabbits used in 2011 are described in the manure decree of December 2006 (or MAP3): [http://www.vlm.be/SiteCollectionDocuments/Publicaties/mestbank/bemestingsnormen\\_2013.pdf](http://www.vlm.be/SiteCollectionDocuments/Publicaties/mestbank/bemestingsnormen_2013.pdf).

Unfortunately no translation in English is available.

For dairy cows, in MAP3, these N-excretion factors depend on the average milk production per cow. Till 2006 the N-excretion factors of the manure action plan (MAP2bis) is used.

In Wallonia nitrogen excretion factors were first determined for the implementation of the CE Nitrates Directive 91/676\_ but were representing the nitrogen *after* deduction of the atmospheric losses, so new factors were calculated on this basis for the purposes of estimating atmospheric emissions. A check for update has been done with the latest legislation (PGDA 2014, <http://www.nitrawal.be/resources/shared/telechargements/feuilletpgdallv2.pdf> ) and no significant difference has been observed. So the Nex factors are kept unchanged.

Table 5-2 gives an overview of the livestock numbers and N-excretion factors (weighted average) used in both regions in 2013.

Table 5-2 Animal number and weighted average of nitrogen excretion factors for each animal category in Flanders and in Wallonia (2013)

Category	Population		Weighted Average Nex (kg N/head.yr)	
	<i>Flanders</i>	<i>Wallonia</i>	<i>Flanders</i>	<i>Wallonia</i>
Dairy Cattle	249521	202 080	116,1	120,5
Brood cows	164307	300 138	65,0	97,8
Other cattle	873331	674 322	43,3	47,0
Fattening Pigs	4180260	216 501	11,3	16,1
Swine	2069509	110 285	6,2	12,1
Sheep	56913	48 375	8,1	6,9
Goats	24974	10 665	8,7	7,6
Horses	39036	13 341	48,5	75,0
Rabbits and fur animals	63775	NE	2.8	-
Laying Hens	6680025	1 014 214	0,8	0,8
Broilers	18177660	3 058 233	0,6	0,4
Other Poultry	3344109	275 918	0.5	0,6

The Table 5-3 gives the allocation of animals to animal waste management system in Wallonia.

Table 5-3 Allocation of animals to AWMS for each category in Wallonia (2013)

	<b>Solid storage</b>	<b>liquid storage</b>
Bovines under 6 months	87%	13%
Bovines between 6 months and 1 year: male	90%	10%
Bovines between 6 months and 1 year: female	87%	13%
Bovines more than 1 year for fattening: male	87%	13%
Bovines more than 1 year for reproduction: male	77%	23%
Bovines more than 1 year: female	77%	23%
Dairy cows	56%	44%
Brood cows	91%	9%
Swine (included piglet & fattening pigs)	25%	75%
Sows	42%	58%
Breeding males	43%	57%
Lambs	100%	0%
Sheep	100%	0%
Goat	100%	0%
Horses	100%	0%
Broilers	89%	11%
Laying hens	6%	94%
Other poultry	26%	74%

The allocation of animals to animal waste management system (AWMS) in Wallonia (see Table 5-5) comes from Statbel, the agricultural census of 1992 and 1996, where those data were collected by animal type. Those data are not collected on a yearly basis by Statbel given their slow pace of change. However, an update of the 1996 data would likely be useful in the near future. So far we have no information about Statbel planning regarding this update. Experts from the sector have been contacted and they confirm that these figures are still valuable in the absence of new detailed information.

The allocation of animals to AWMS (Table 5-4) originate in Flanders from the Department of Agriculture and Fisheries.

Table 5-4 Allocation of animals to AWMS for each category in Flanders (2013)

	<b>Solid storage (%)</b>	<b>Liquid storage (%)</b>
<b>Bovine</b>		
Slaughter calves	0	100
Bovines under 1 year	94	6
Bovines under 1yr for replacement	91	9
Bovines from 1 to 2 year	92	8
Bovines from 1 to 2 yr for replacement	34	66
Bovines more than 2 year	78	22
Dairy cows	30	70
Brood cows	91	9

<b>Swine</b>		
Piglet from 7 to 20 kg	1	99
Fattening pigs from 20 to 110 kg	1	99
Fattening pigs more than 110 kg	2	98
Boars	31	69
Sows including piglets less than 7 kg	2	98
<b>Sheep</b>	100	0
<b>Goats</b>	100	0
<b>Horses</b>	100	0
<b>Rabbits and fur animals</b>	100	0
<b>Poultry</b>	With litter	Without litter
Broilers (for breeding)	100	0
Broilers (for fattening)	100	0
Laying hens (for breeding)	36	64
Laying hens	33	67
Ostriches	100	0
Turkeys	100	0
Other poultry	100	0

## 5.2. *Animal husbandary and manure management (category 3B)*

### 5.2.1. NH<sub>3</sub>

The NH<sub>3</sub> emission estimation from livestock is based on the amount of gross nitrogen excreted by each animal category, estimated through local production factors (see 5.1.4). The calculation takes into account the different stable types, the number of days in pasture, the different manure management systems, the manure application on land. The models used in the two regions differ and are individually described hereunder.

In Flanders, for the entire time series, the EMANV-model (Foqué & Demeyer, 2009) is used. As described in 5.1.4 this model calculates the NH<sub>3</sub> emission in different emission stadia taking into account the manure flow. From 1990 to 1999 the NH<sub>3</sub> emission is calculated on the level of the municipality, using livestock numbers from Statbel. From 2000 on this is done on the level of the stable, using detailed input data, as animal number, stable type and N-production, on the level of the farm of the Flemish Land Agency.

There is a significant decrease in NH<sub>3</sub> emissions from 4B1a, 4B1b and 4B8 between 1999 and 2000. This decrease is mainly due to the implementation of the successive Manure Action Plans in Flanders.

In 1991 there was the first Manure Decree. One of the items of this Decree was the reduction of the period (months) in which manure can be applied to land. This had a minor impact on the NH<sub>3</sub> emissions. The Manure Action Plan2bis in 2000 focussed on the reduction of the manure surplus. Among others this was done by reducing the amount manure that can be applied to land and furthermore the obligation to apply manure in an emission poor way. Also the regulation concerning manure processing had an impact on the NH<sub>3</sub> emission. Since 2003 all new stables for swine and

poultry have to be constructed in an emission poor way. The licensed stables are inventoried by the manure bank of the Flemish Land Agency. Based on the amount of ammonia emission-poor stables, the stable emission factors for swine and poultry are yearly adjusted.

In Flanders, the use of sewage sludge on agricultural soils is forbidden. This is described in the manure decree (article 13, paragraph 8:

<http://navigator.emis.vito.be/milnav-consult/plainWettekstServlet?wettekstId=17942&lang=nl>).

Unfortunately no translation in English is available.

In Wallonia, the emissions are calculated using a model developed by a consultant agency Siterem updated this year to take into account new data: 1) the use of injectors for the application of slurry (activity data coming from national statistics & abatement factors coming from the 2013 EMEP Guidebook), 2) new emission factor for the storage of solid manure (coming from a Walloon study<sup>5</sup>) and 3) a new emission factor for the application of mineral fertilisers derived from the 2013 EMEP Guidebook.

Different data are used as input in the model which then calculates NH<sub>3</sub> emissions. Four emissions sources were pointed: the excreta of animals 1) during grazing, 2) deposited in buildings and collected as either liquid slurry or solid manure, 3) applied to land (animal manure) and 4) the use of mineral fertilisers.

Manure application to land counts for a little less than 40% of the NH<sub>3</sub> agricultural emissions while the three other sources represent each one roughly 20% in 2013.

Emissions from the manure application has dropped in 2003 by roughly 10% because it is the first year with data on injectors and the decrease is going on with the multiplication of the use of injectors (from 14% of the equipment sold in 2003 to 30% in 2013, *[communication from experts from Agra-Ost]*). Emissions from the manure storage and from grazing are following the trends of the cheptel evolution.

The emission factor for the storage of cattle solid manure has been reviewed in the light of a study conducted during 6 years in Walloon cattle stables<sup>4</sup>. The aim of the study was to follow the production and the storage of cattle manure and to assess the environment impact. This study has allowed to divide by 3 the emission factor linked to cattle solid manure storage (for the other animals, the emission factors has not been changed).

## 5.2.2. Particulate matter

In Wallonia, the dust emissions for the category 3B are calculated by using the Tier 1 emission factors coming from the 2013 EMEP Guidebook (table 3.3 p.17 *Chap. 3.B Manure management*) and the activity data from national statistics.

In Flanders, the TSP emissions are calculated for the category 3B by using emission factors as described in Table 5-5 (Sleeuwaert & Polders, 2006). PM10 and PM2,5 are calculated as a percentage of TSP. The IEF of TSP, PM10 and PM2,5 for the fattening pigs is derived from a study recently performed by the Institute for Agricultural and Fisheries Research (Van Ransbeeck, 2013). The subdivision per NFR subcategory (dairy/non-dairy, laying hens/broilers/turkeys/other poultry) is performed by taking into account the animal numbers given in Table 5-5.

---

<sup>5</sup> Mathot *et al.* (2011) *Filière de gestion des effluents d'élevages bovins : Impact environnemental de la production et du stockage*. UCL, CRAW

Table 5-5 Emission factors for the calculation of TSP emissions and fraction of PM10 and PM2,5 of livestock breeding in Flanders

		TSP	%PM10	%PM2,5
poultry	g/year/animal	49.4	49	14
cattle	g/year/animal	394	44	11
Fattening pigs	g/year/animal	227	44	3
Other swine	g/year/animal	446	44	10

### 5.2.3. NO<sub>x</sub>

In Wallonia and Flanders, the NO<sub>x</sub> emissions for the category 3B are calculated by using the Tier 1 emission factors coming from the 2013 EMEP Guidebook (table 3.2 p.15 *Chap. 3.B Manure management*) and the activity data from national statistics (Wallonia) and the Flemish Land Agency (Flanders).

### 5.2.4. NMVOC

In Wallonia and Flanders, the NMVOC emissions for the category 3B are calculated by using the Tier 1 emission factors coming from the 2013 EMEP Guidebook (table 3-3 p.16 *Chap. 3.B Manure management*) and the activity data from national statistics (Wallonia) and the Flemish Land Agency (Flanders).

## 5.3. Direct soil emission (category 3D)

### 5.3.1. Synthetic fertilizer use (category 3Da1)

#### 5.3.1.1 NH<sub>3</sub>

In Flanders, the NH<sub>3</sub> emission from fertilizer use is calculated using the same EMAN-model in which the area of a certain agricultural zone is multiplied with the respective fertiliser use (kg N/ha) and emission coefficient. Depending of the type of mineral fertiliser, a different emission coefficient is used. The relative amount of different types of mineral fertiliser used in Belgium originate from the *International Fertilizer Industry Association (IFA)*. The emission factor (%) in table 5.5 is the average of all emission coefficients.

For the amount of fertilizer use the *Department Agriculture and Fisheries* and the *Institute for Agricultural and Fisheries Research* conducts surveys on a representative sample of the different types of agricultural businesses and produces yearly weighted average values on the fertiliser use, taking into account the manure pressure (Campens & Lauwers, 2002).

In Wallonia, the NH<sub>3</sub> emissions are calculated by multiplying the quantity of fertilizer use (Wallonian statistics) by an updated emission factor, derived from the 2013 EMEP Guidebook (Table 3-2 p 14 *Chapter 3D Crop production & agricultural soils*) and the same relative amount of different types of

mineral fertiliser than in Flanders. With this update, the emission factor has nearly doubled (from 2.3% to 4.2%).

The data on the use of mineral fertiliser come from the Agricultural Economy Analysis Department of the region. The use of mineral fertilizer is decreasing since 1990, The amount of synthetic fertilizer use and emission factors per region is shown in Table 5-6.

Table 5-6 The amount (kg N) of synthetic fertilizer use (per ha) (2013) and the emission factor used in each Region

	Synthetic fertilizer use (kg N /ha)	Emission factor (%)
<b>Flanders</b>		
<b>Dunes</b>	139	3.99
<b>Kempen area</b>	98	3.99
<b>Sandy area</b>	107	3.99
<b>Sand Loam area</b>	105	3.99
<b>Loamy area</b>	106	3.99
<b>Polders</b>	139	3.99
<b>Luikse Meadow area</b>	98	3.99
<b>Wallonia</b>	101.38	4.2

#### 5.3.1.2 NO<sub>x</sub>

In Wallonia and Flanders, NO<sub>x</sub> emissions are calculated following the Tier 1 Methodology of 2013 EMEP Guidebook (Table 3-1 p 11 of the EMEP Guidebook). The quantity of fertiliser is equal to the sum of mineral and organic fertiliser (animal manure).

#### 5.3.1.3 NMVOC & PM

In Wallonia, NMVOC & PM emissions are calculated following the Tier 1 Methodology of 2013 EMEP Guidebook (Table 3-1 p 11 of the EMEP Guidebook). The activity data are the number of ha of cropped area (national statistics). In Flanders only the NMVOC emissions are calculated analogously to Wallonia.

### 5.3.2. Manure processing (category 3Dd)

As described above in chapter 5.1, Flanders has a severe manure surplus. Therefore successive manure action plans (MAP) are implemented. Among other things, the MAP describes the amount of manure that a farmer can apply to his agricultural soils. Briefly, this depends on the proportion of the amount manure produced to the available agricultural soils of that farmer. The manure surplus (the part that may not be applied to the soil) must be either exported or processed. This amount (net export and amount processed) is inventoried by the Manure Bank of the VLM. NH<sub>3</sub> -emissions from

manure processing in Flanders in 2013 account for 1% of the total NH<sub>3</sub>-emission in Flanders. The emissions for manure processing are calculated using the same model as used for the calculation of ammonia emission from livestock and synthetic fertilizer: the EMAN-model. The Flemish coordination centre for manure processing (VCM, [http://www.vcm-mestverwerking.be/organisation/detail\\_en.phtml?id=1](http://www.vcm-mestverwerking.be/organisation/detail_en.phtml?id=1)) inventories and publishes yearly the operational manure processing companies and the techniques used. Based on this information, the amount and type of processed manure and the corresponding emission coefficient, the NH<sub>3</sub> emission from manure processing can be calculated.

NH<sub>3</sub> emissions from manure processing in Flanders are taken into account from 2000 on. Before 2000 manure processing was rare. The amount of processed manure from 2000 on increases significantly. However, the NH<sub>3</sub> emission stabilizes since 2008. This is due to the techniques used. Since 2008 more manure is processed in a biological treatment. This technique has a significant lower emission coefficient (0.05%) than e.g. biothermal drying (5.63%). The amount of manure processed by biothermal drying has stabilized.

### 5.3.3. Urine and dung deposited by grazing animals (category 3Da3)

In both regions, the ammonia emission from grazing is estimated, taking into account the number of days in pasture, the nitrogen excreted by each animal category, and the EEA emission factor of 8%. In Table 5-7 an overview is given of the different factors used in both regions and for the different grazing animal categories.

Table 5-7 The days in pasture (%/yr), nitrogen excreted on pasture (ton) and the emission factor used for each grazing category in 2013

	Days in pasture (% of year)	Nitrogen excreted on pasture (ton)	Emission factor (%)
<b>Flanders</b>			
bovine	± 50	13053	8
Sheep	80	361	8
Horses	50	1144	8
<b>Wallonia</b>			
Bovine	50	40130	8
sheep	50	153	8
horses	50	502	8

## 5.4. The use in pesticides (4G)

In Flanders HCH-emissions are calculated for the past. There is a ban on the use of HCH in Belgium from 2002. The activity data are taken from a study performed by VITO/TNO under the authority of VMM (Sleeuwaert, 2012).

## **5.5. Recalculations**

In Flanders, during this submission February 2015, NH<sub>3</sub> emissions from 2012 are updated. In the submission of February 2014, NH<sub>3</sub>-emissions 2012 were reported as the same as NH<sub>3</sub> emissions from 2011. This was due to lack of activity data on the time of reporting.

In Wallonia, as mentioned before, recalculations occurred in the category 3B. Thanks to new information of a Walloon study (*Mathot & al, 2011*), the emission factor for the storage of solid manure of cattle has been reviewed for the entire time series. This results in a decrease of the emission of  $\pm 66\%$ .

Also in Wallonia, in category 3D, information about the application of slurry and the use of precise equipment (injectors), a reduction of emissions has been applied since 2003. In 2013, this measure results in a reduction of  $\pm 12\%$  of the emissions from manure application.

Finally, in Wallonia, the emission factor for the use of mineral fertilizer has been updated with the value of 2013 EMEP guidebook (Table 3-2, p 14 *Chapter 3.D*). This time, the result is roughly a doubling of the emissions of the category 3Da1.

## **5.6. Planned improvements**

In Flanders, an actualisation of the EMAN model is planned.

In Wallonia, no improvements are planned in the near future.

## Chapter 6. Waste (NFR sector 5)

Waste sector emissions are classified into 4 categories as described in Table 6-1.

Table 6-1. Main emissions of the 4 waste categories in Belgium

Waste categories	Main emissions
Solid waste disposal sites	NM VOC, PM <sub>2.5</sub> , PM <sub>10</sub> , TSP
Waste water treatment	NH <sub>3</sub>
Waste incineration & cremation	Hg, Dioxins & furans, HCB
Other	NH <sub>3</sub> , PM <sub>2.5</sub> , PM <sub>10</sub> , TSP, BC

### 6.1. Solid waste disposal on land (category 5A)

The NM VOC emissions from land disposal of solid waste are calculated in Flanders and in Wallonia.

The methodology applied by the Flemish region to estimate NM VOC emissions is the same methodology as the one used to calculate the CH<sub>4</sub>-emissions (see description in NIR of April 2015, Chapter 8.2.). Just one factor is different: biogas contains 2420 ppm NM VOC (D'Haene et al., 2002).

No waste disposal sites are located in the Brussels Capital Region in Belgium.

In Wallonia, NM VOC emissions calculations are based on Tier 1 methodology of the 2009 EMEP/EEA Guidebook. The volume of landfill gas resulted from the IPCC model used for GHG inventory and from the recovery data of the sites managers. The methodology is the one described in the 2006 IPCC Guidelines. The emission factor used for NM VOC is 5.65 g NM VOC/m<sup>3</sup> landfill gas, coming from the 2009 EMEP/EEA Guidebook. the 2013 EMEP/EEA Guidebook, the emission factor is expressed in g NM VOC/Mg waste. However, the composition of solid waste disposed is changing every year and by the way, the content in NM VOC is supposed to change too. We must also take into account the volume of biogas recovered. So it is preferred to use the net volume of biogas (calculated following the IPCC methodology) and the emission factor of 5.65 g NM VOC/m<sup>3</sup> landfill gas, which corresponds to the 1.56 kg/Mg waste (2013 EMEP/EEA Guidebook) with the hypothesis of a default methane content of 50% (value close to the Walloon situation).

PM emissions (PM<sub>2.5</sub>, PM<sub>10</sub> & TSP) are calculated following the Tier 1 methodology of the 2013 EMEP/EEA Guidebook.

### 6.2. Waste incineration (category 5C)

The waste incineration category (category 5C) includes incineration of municipal and industrial waste, incineration of hospital waste and incineration of corpses (crematoriums). The emissions of the waste incineration plants with energy recovery are allocated to the category 1A1a.

The category 5C1a and 5C1b are key categories of PCDD/F respectively Ni and Hg emissions in terms of emission trend. The category 5C2 is a key category of PCDD/F emissions in terms of emission level and trend.

## Waste incinerators

In Wallonia, following a legal decree in 1998, the air emissions from municipal waste incineration were measured in 1998 by ISSEP and the results were validated by a Steering Committee. Since 2004, the amount of incinerated waste (in ton) and the annual emissions are reported annually by the operators in a software dedicated to environmental reporting, called REGINE, in the context of PRTR. The annual emissions are calculated on the basis of stack measurement (when they are available) or emission factors (when stack measurement are not performed annually).

The only hospital waste incinerator has closed since 2005.

Since 2000, a continuous measurement of dioxins has been put in place for the 4 incinerators: <http://environnement.wallonie.be/data/air/dioxines/menu/menu.htm>

In the early 1990s, about 45% of the waste was still incinerated without energy recovery. Since 2006, the 4 municipal waste incineration plants are fully equipped to produce electricity. The emissions with energy recovery are allocated in the energy sector, category 1A1a. A small part of the emissions from municipal waste incineration is still allocated in the waste sector, category 5C, when waste is incinerated without energy recovery because of occasional problems in the energy recovery systems. In 2010, this represents 2% of the incinerated waste. In 2013, this represents 20% of the incinerated waste. In 2013, the fraction of waste that has been incinerated without energy recovery is higher than the previous years because the turbine of 2 of the 4 waste incineration plants in Wallonia had to be stopped during more than 6 months for repair.

In Flanders, the plants are obliged to report their emissions yearly in an emission report. These data are integrated in the emission inventory. All (intermunicipal) waste incineration plants produce electricity since 2006. The emissions are allocated in the category 1A1a when energy is recycled or in 5C1bi (industrial waste incineration). It was not possible yet to split up the emissions between industrial and municipal waste incineration, hence the emissions due to municipal waste incineration without energy recycling are also allocated in category 5C1bi instead of 5C1a. A methodology will be developed to distinguish between the emissions of industrial and municipal waste incineration.

In Flanders the PCDD/F emissions of 1990 and 1995 (industrial and domestic waste) are based on the results of a study performed by VITO under the authority of VMM (Polders et al., 2003). Since 2000 the emissions of domestic waste incineration are reported in the yearly environmental reports. Since 2000 the emissions of industrial waste incineration are calculated by using activity data and emission factors. The activity data are the amount of waste obtained from OVAM (Public Waste Agency of Flanders). The emission factors are taken from the UNEP Standardized Toolkit for PCDD/F (Table 6-2).

The HCB emissions are calculated by using activity data and emission factors. The activity data are the amount of waste obtained from OVAM (Public Waste Agency of Flanders). The emission factors are taken from the EMEP/CORINAIR Guidebook for HCB (Table 6-3).

Table 6-2 Emission factors of PCDD/F for the sector 1A1a Incineration of waste in the Flemish region

Fuel	Unit	Value	Reference
Industrial waste	µg TEQ/tonne	0.5	UNEP Standardized Toolkit; Category 1a4: Waste incineration; Municipal solid waste incineration; High tech. combustion, sophisticated APCS
Hazardous waste	µg TEQ/tonne	0,75	UNEP Standardized Toolkit; Category 1b4: Waste incineration; Hazardous waste incineration; High

			tech. combustion, sophisticated APCS
Clinical waste	µg TEQ/tonne	1	UNEP Standardized Toolkit; Category 1c4: Waste incineration; Medical/hospital waste incineration; High tech, continuous, sophisticated APCS
Sewage sludge	µg TEQ/tonne	0,4	UNEP Standardized Toolkit; Category 1e3: Waste incineration; Sewage sludge incineration; State-of-the-art, full APCS

Table 6-3 Emission factors of HCB for the sector 1A1a Incineration of waste in the Flemish region

Fuel	Unit	Value	Reference
Industrial waste	g/tonne	0.0001	EMEP/CORINAIR Guidebook (2005)
Hazardous waste	g/tonne	0.01	EMEP/CORINAIR Guidebook (2005)
Clinical waste	g/tonne	0.019	EMEP/CORINAIR Guidebook (2005)
Sewage sludge	g/tonne	0.002	EMEP/CORINAIR Guidebook (2009)
Domestic waste	g/tonne	0.001	EMEP/CORINAIR Guidebook (2005)

Emissions due to clinical waste incineration (category 5C1biii) are included in category 5C1bi (industrial waste incineration).

In Brussels, the last waste incinerator without recuperation of energy has closed in 1998.

#### Emissions by cremation

For Flanders: the activity data is 38977 cremations in 2013 (derived from the yearly STATBEL<sup>6</sup>. statistics of crematoria), for dioxins an emission factor of 0,069 µg TEQ/cremation is used (Results of measurements made by the Flemish government – Department Leefmilieu/Afdeling Milieu-inspectie). The calculation of particulate matter (TSP, PM10, PM2,5) is done with an emission factor of 0.005 kg/cremation and for Hg an emission factor of 0.049 g/cremation is used.

For Wallonia: for dioxins, the emission factor from the Netherlands of 4µg TEQ/corpse, reported in EMEP (Emission Inventory Guidebook December 2006 B991-9) is applied. A specific emission factor is used for Hg emissions. This emission factor is equal to 2 g/corpse and is coming from measurements and analysis made in a Walloon crematorium.

Emissions from the other pollutants are estimated using the emission factors of the EMEP/EEA guidebooks 2013. The number of corpses is coming from the national Belgian statistics available on the website of STATBEL.

For Brussels, the emission factor for dioxins is 27 ng TEQ/cremation as stated in the EMEP/EEA 2013 guidebook. The number of cremations comes also from national statistics available on the website of STATBEL, for 2013, it was 5651.

<sup>6</sup> <http://statbel.fgov.be/fr/statistiques/chiffres/population/autres/cremations/>

### Open combustion of waste (small scale waste burning) (category 5C2)

Only Flanders estimates emissions of combustion in open barrels of particulate matter, dioxins and PAH's. These emissions are allocated in sector 5C2. To make the calculation, it is assumed that 5% of the average amount of municipal waste is burnt in open barrels (Van Rompaey et al., 2001). The amount of municipal waste per household can be found on the website of the Public Waste Agency of Flanders ([www.ovam.be](http://www.ovam.be)). The number of households can be found on [www.statbel.fgov.be](http://www.statbel.fgov.be). The emission factor of TSP is taken from a study performed by VITO under the authority of AMINAL/Aminabel (Wevers, 2002). The emission factors of dioxins and PAH's are taken from a study performed by VITO/TNO under the authority of VMM (Sleeuwaert, 2012). Emission factors for PM10 and PM2,5 are taken from TNO (2001). Other combustion emissions are not estimated.

In Wallonia, these emissions are not estimated.

### **6.3. Wastewater treatment (category 5D)**

In Belgium, emissions originating from septic tanks are estimated by multiplying the emission factor (an NH<sub>3</sub> emission factor of 750 g/person is used) by the number of inhabitants not connected to a municipal wastewater treatment plant. In Flanders, the number of inhabitants not connected to a municipal wastewater treatment plant is calculated by subtracting the number of inhabitants connected to a municipal wastewater treatment plant (data obtained from the Team Data Management Sewerage Infrastructure of the Flemish Environment Agency) from the total number of inhabitants in Flanders (<http://bestat.economie.fgov.be/BeStat/BeStatMultidimensionalAnalysis?loadDefaultId=440>).

### **6.4. Other**

NH<sub>3</sub> emissions from compost production, allocated in category 5B1 are estimated in Wallonia using regional activity data combined with a default emission factor of 0,24 kg NH<sub>3</sub>/ton compost (EMEP/EEA Guidebook 2013).

The emissions of PM2,5 and PM10, due to smoking of tobacco in Flanders, used to be reported under 7A, but it was decided to allocate these emissions in the more appropriate sector 3D3, according to the EMEP/EEA Guidebook 2013 (see also Chapter 4 Industrial Processes).

In Wallonia and Flanders, the other pollutants emissions come from the annual environment report of waste companies (other than incinerators). Each year, companies have to fulfil an integrated environmental survey in the context of PRTR. The data in the air emissions section are used to compile the Walloon emissions.

### **6.5. Recalculation**

In Flanders the emissions of dioxins and HCB are re-allocated between 1A1a and 5C1. The emissions of the waste incineration plants with energy recovery are allocated to the category 1A1a.

EF of the EMEP/EEA 2013 guidebook are applied for sectors 5C1a, 5C1biii and 5C1bv in Brussels.

Brussels has replaced the NE for NH<sub>3</sub> for sectors 5C1a and 5C1biii, using the EF from the EMEP/EEA 2013 guidebook. Replacement of NE for Se, PAH's, HCB and PCBs with the EF of the EMEP/EEA 2013 guidebook for sectors 5.C.1.b.iii and 5.C.1.b.v in Brussels.

## **6.6. QA/QC**

All emissions delivered by the plants are validated and verified by a team of people experienced in emission inventories. In addition, each year a trend analysis is carried out for all emissions per industrial plant and sector. If any inconsistencies or problems are detected by the team, the industry involved is contacted. Numerous contacts take place with the plant operators as well as with the federations involved. In exceptional cases the inspection services are contacted.

## **6.7. *Planned improvements***

In the Flemish region, the following improvements are planned:

- It was not possible yet to split up the emissions between industrial/hazardous/sewage sludge and municipal waste incineration, hence the emissions due to municipal waste incineration without energy recycling are also allocated in category 5C1bi instead of 5C1a. A methodology will be developed to distinguish between the emissions of municipal and industrial waste incineration, and as far as possible between industrial, hazardous and sewage sludge incineration.

## **Chapter 7. Land use and land use change**

No emissions are reported for LRTAP or NEC. The sector is not included in the NFR tables.

## Chapter 8. Other and natural emissions

### 8.1. Biogenic emissions

#### *Flanders*

NMVOC emissions of different forest types and of grassland are reported under IPCC code 11C. These biogenic emissions are substantial and can account up to nearly 15 % of the total reported NMVOC emissions.

The methodology to calculate the collective estimation of the biogenic NMVOC emissions is described in Van Hyfte & Van Langenhove (2000) and based on a model by Guenther et al. (1993).

The basic formula used to calculate the biogenic emissions is:

$$E = \sum_{y=1}^z N_y (D_y \cdot \varepsilon_y \cdot \gamma_y)$$

with:  $N_y$ : the total area taken by ground cover y ( $\text{m}^2/\text{year}$ )

$z$ : the number of species of ground cover y

$D_y$ : leaf density ( $\text{kg dry matter}/\text{m}^2$ )

$\varepsilon_y$ : VOC emission factor for ground cover y at 30°C and light intensity of 1000  $\mu\text{mol}/\text{m}^2\text{s}$  ( $\mu\text{g}/\text{m}^2\text{hour}$ )

$\gamma_y$ : correction factor for real leaf temperature and light intensity

The ground cover in Flanders is defined by the wood mapping performed by the Flemish Region, based on visual reading of coloured infrared aerial views taken during the period 1981-1992 and ground monitoring. After handling the information this results in a ground accuracy to within 1 are. The ground cover is corrected based on the LUC matrix. The LUC matrix is determined by the Gembloux University (Gembloux Agro Bio Tech), a study conducted specifically for the LULUCF reporting in Belgium (Bauwens et al., 2011).

The emission factors give the emissions in  $\mu\text{g}$  per hour in terms of the leaf density ( $\text{g dry matter}/\text{m}^2$  ground cover). Emission factors are taken from literature and are specified for different compounds of NMVOC (isoprene, monoterpenes and other VOC) and for different kinds of ground cover. An overview of the emission factors used is given in Table 8-1.

Table 8-1 Emission factors for isoprene, monoterpenes and other VOC for different species of ground cover in Flanders based on Simpson et al. (1999)

Ground cover	Isoprene (ng/m <sup>2</sup> /s)	Monoterpenes (ng/m <sup>2</sup> /s)	Other VOC (ng/m <sup>2</sup> /s)
<i>Broadleaf trees</i>			
Beech	8.89	57.78	192.78
Oak/American oak	5333.33	17.78	192.78
Poplar	5333.33	0.00	192.78
Other	5333.33	17.78	242.22
<i>Mixed broadleaf</i>			
Beech	20.56	152.22	285.28
Oak	5298.61	112.50	285.28
Poplar	5298.61	95.00	285.28
Other	5298.61	112.50	317.50
<i>Mixed conifers</i>			
Larch	1349.44	91.11	197.22
Scots pine	2492.22	168.06	305.56
Black pine	2492.22	321.67	605.56
Spruce	4658.33	302.78	572.78
Douglas	3349.44	225.83	572.78
Other (default)	2492.22	321.67	258.89
<i>Conifers</i>			
Larch	8.33	125.00	192.78
Scots pine	19.44	291.67	359.72
Black pine	19.44	583.33	359.72
Spruce	388.89	583.33	770.83
Douglas	27.78	416.67	770.83
Other	19.44	583.33	287.78
<i>Grassland</i>	0.00	11.11	23.33

The leaf density of a tree species expresses the amount of dry matter (g) of a tree in terms of the ground area, taken by this species. The leaf density can vary significantly in the course of the seasons. Since several factors can influence the leaf density, the calculations are made with average leaf densities (already taken in account in table 9.1).

Since the leaf temperature and the light intensity are the most important factors that influence VOC emissions, a correction factor (specified for isoprene and terpene emissions) is taken from literature.

## **Wallonia**

### **Methodology**

The methodology used by the AWAC is based on Simpson and Guenther (EMEP/CORINAIR atmospheric Emission Inventory Guidebook, 1999). The mass emission time of a plant species occupying a given area is given by the relation:

$$\text{Hourly mass emission (g / h)} = S * B * C * FE (T^{\circ}, \text{Light})$$

S = Surface in m<sup>2</sup>

EF = emission factor standard of the species (g / gh)

B = foliar biomass of the species (g / m<sup>2</sup>)

C (T °, Light): VOC emissions are highly dependent on temperature and sometimes light, depending on the considered VOCs. This is taken into account by the correction factor dimensionless. This factor can be calculated on an hourly basis, but the calculation has been done here on a monthly basis, which here constitutes a good compromise between the accuracy of the estimate and the availability of data (data on PAR, photosynthetically active radiation from 400 to 700 nm, are not available on an adequate scale for the Walloon Region). This simplification increases the error of the order of 20%, which is far less than the uncertainties in the emission factors.

Isoprene emissions depend on both temperature and light intensity. The correction factor is then:

$$C = CL * CT$$

CL = Number of days per month \* Number of hours of daylight the month (depending on latitude)

$$CT = \text{Exp} ((95000 * (T-T_s)) / (8.314 * T * T_s)) / (1 + \text{exp} ((230000 * (T-314)) / (8.314 * T * T_s)))$$

T = temperature in Kelvin foliar experimental (measured)

T<sub>s</sub> = temperature reference leaf (very generally 303 K or 30 ° C) at which the emission factor is determined

The other figures are empirical coefficients and the ideal gas constant.

For monoterpenes and other VOC, emission depends only on the temperature and the relationship becomes:

$$C = CL * CT$$

CL = Number of days per month \* 24 (hours)

$$CT = \text{Exp} (0.09 * (T-T_s))$$

T = temperature in Kelvin foliar experimental (measured)

T<sub>s</sub> = temperature reference leaf (very generally 303 K or 30 ° C) at which the emission factor is determined.

### **Forest area**

The area of forest is taken from the forest inventories. The first Walloon forest inventory was conducted between 1979 and 1984 (central year is 1981). The current permanent systematic sampling of the permanent forest inventory was conducted between 1994 and 2008 (central year is 2001) and covers each year 10 % of the approximately 11000 sampling points (Lecomte & Rondeux, 1994). The third cycle of the forest inventory started in 2009 and first results were made available by the end of 2011 (central year is 2010).

### **Biomass**

Regarding leaf biomass, Simpson and Guenther (1995) strongly recommend the use of local data if they are available. For the main Walloon forest species (oak, beech, spruce, Douglas fir, pine), we therefore sought densities measured in Belgium, including those compiled by Duvigneaud et al in the 70's, or densities measured in neighboring regions (North of France and the Netherlands). For other species, the values used in France (Luchetta et al., 2000) were included (Table 8-2).

Table 8-2 Leaf biomass for the main Walloon forest species

Species (latin name)	Leaf biomass (kg/ha)	Country of measure	Source
<i>Acer pseudoplatanus</i>	3300		in Luchetta et al, 2000
<i>Alnus glutinosa</i>	2800	B	in Luchetta et al, 2000
<i>Betula pendula</i>	3200	B	Duvigneaud et al ,1977
<i>Carpinus betulus</i>	3500	F	in Luchetta et al, 2000
<i>Castanea sativa</i>	3600	F	in Luchetta et al, 2000
<i>Fagus sylvatica</i>	3118	B, F, NL	Duvigneaud et al 1977 ; Gloaguen et al, 1982 ; Bartelink 1997
<i>Fraxinus excelsior</i>	2700	DK	in Luchetta et al, 2000
<i>Larix decidua</i>	3300		in Luchetta et al, 2000
<i>Picea abies</i>	16390	B, F	Duvigneaud et al ,1977 ; Teller, 1983 ; Guns, 1990 ; Belkacem et al 1992 ; Ranger et al, 1981 ;
<i>Pinus nigra laricio</i>	8133	B, F	Neiryck et al 1998 ; Bonneau, 1995
<i>Pinus nigra nigra</i>	9400	F	in Luchetta et al, 2000
<i>Pinus sylvestris</i>	8000	F	in Luchetta et al, 2000
<i>Populus sp</i>	3300		in Luchetta et al, 2000
<i>Prunus avium</i>	3300		in Luchetta et al, 2000
<i>Pseudotsuga menziesii</i>	12633	B, F	Duvigneaud et al ,1977 ; Ponette et al, 2000, Ranger et al, 1996
<i>Quercus rubra</i>	3200		in Luchetta et al, 2000
<i>Quercus sp (robur + petrae)</i>	3290	B, F	Duvigneaud et al ,1977 ; Gloaguen et al, 1982

### Emission factors

No emission factor determined in Belgium has been found in the literature. Emission factors are essentially the compilation made by Luchetta et al. (2000) for France. The consistency of these emission factors with those taken in the compilation of Hewitt (2001), which includes the emission factors of more than 1200 species, has been systematically verified. Factors proposed by Hewitt (2001) were used for three species: red oak (not treated with Luchetta), chestnut (the figure seems Luchetta underestimated), beech (Luchetta used for monoterpenes a factor of 21.7, based on a measurement made in France, which strongly deviates values quoted in 6 other references) (Table 8-3).

Table 8-3 Emission factors for a number of species

Species	Emission factor isoprene ( $\mu\text{g/g}\cdot\text{h}$ )	Emission factor monoterpene ( $\mu\text{g/g}\cdot\text{h}$ )	Emission factor Other VOC ( $\mu\text{g/g}\cdot\text{h}$ )	Vegetation period	
<i>Acer pseudoplatanus</i>	0	0	1,5	1 May	30 October
<i>Alnus glutinosa</i>	0,1	3,4	1,5	1 May	30 November
<i>Betula pendula</i>	0,01	2,9	1,5	15 March	15 October
<i>Carpinus betulus</i>	0	0,1	1,5	15 April	15 October
<i>Castanea sativa</i>	0	13,66	1,5	15 April	15 October
<i>Fagus sylvatica</i>	0,1	0,47	1,5	15 April	30 October
<i>Fraxinus excelsior</i>	0,1	0	1,5	1 June	30 October
<i>Larix decidua</i>	0,1	8,2	1,5	15 March	15 October
<i>Picea abies</i>	1,1	2,1	1,5		
<i>Pinus nigra laricio</i>	13,2	0	1,5		
<i>Pinus nigra nigra</i>	13,2	0	1,5		
<i>Pinus sylvestris</i>	0,1	7,9	1,5		
<i>Populus sp</i>	51	4,6	1,5	1 May	30 September
<i>Prunus avium</i>	0	0,3	1,5	1 May	30 October
<i>Pseudotsuga menziesii</i>	0,45	14,8	1,5		
<i>Quercus rubra</i>	37,9	1,8	1,5	1 May	30 October
<i>Quercus sp</i>	57,3	0,46	1,5	1 May	15 November

### Correction factors

The average monthly temperatures of IRM were coded for each of the stations. The provincial averages was then calculated. For light, monthly data proposed by Guenther, depending only on the latitude, were used, based on an average latitude of 50 ° N for the Region. These two parameters were used to calculate correction factors CT and CL on a monthly basis at the level of provinces and

districts.

***Vegetation period***

Dates of budburst and leaf fall are listed in 'Ecological Species File' published by the DGRNE (MRW-Walloon Region Ministry, 1999). When calculating emissions from deciduous factor 0,  $\frac{1}{2}$ , or 1 is included in the equation as the leaves are absent or present during 15 days present during all the month.

## Chapter 9. Recalculations and improvements

### 9.1. Recalculations and improvements in the energy sector

- Until the last submission, Wallonia used the database Collector to manage the inventory data's (one collector database by year). But this year, a new air emission software (WAPI) was developed. This software allows the seeing of all data's of a plant or an area source on the complete time series and avoids mistakes during recalculations (jump, zero,...). Some mistakes were corrected as bad values for the Ni and Hg emission factors of heavy fuels in some industrial sectors in 1990, PAH emission factors for the wood boilers in 1A1a, errors of units for some emission factors or direct emissions,...). This new software is also use to report some sectors of the GHG inventory.
- The emission factors of the EMEP/EEA emission inventory guidebook 2013 were used in the three regions. This resulted in new emissions for a number of sectors for the years 1990-2013.  
Actualization of the emission factors according to the guidebook EMEP/EEA 2013
  - o in the Walloon inventory, the new EF are used in almost all of the sectors : 1A1a, 1A2, 1A3c, 1A3dii, 1A4ai, 1A4bi and 1A4c
  - o in Brussels inventory, the new EF are used for sectors: 1A1a, 1A2gviii, 1A3c, 1A3dii, 1A4ai, and 1A4bi.
  - o in the Flemish inventory 1A1a, 1A2, 1A3c, 1A3dii, 1A4ai, 1A4bi and 1A4c
- In the sector 1A3ei (pipeline compressors) in the Walloon region, recalculation of the NO<sub>x</sub> EF for the year 1990 to 2003 with the average EF on the period 2004-2013.
- For some plants in the sectors 1A2 in the Walloon region, plant specific emissions were available for the year 2005 but weren't yet integrated in the inventory. The emissions were calculated via emission factors. The plant specific emissions were integrated in this submission.
- In the sector 1A2gvii in the Walloon region, the fuel consumption in the quarries was misallocated during the years 2002 to 2009. The fuel consumption was allocated in 1A2f and the emissions factors were the emission factors of stationary combustion. This sector has been recalculated with the emission factors of mobile combustion.
- In the Walloon region, for the sector 1A4ai and the sector 1A4bi, there was no estimation of the SO<sub>2</sub> emissions coming from the combustion of natural gas and LPG. The SO<sub>x</sub> EF of gaseous fuels were integrated in this submission.
- Recalculations in the sectors 1A4ci and 1A4cii are due to changes in the energy balances from 2007 on.
- The emission inventory of Persistent Organic Pollutants (POPs) is optimized by VITO under the authority of VMM (Sleeuwaert et al., 2012).
- during the 2014 submission the Flemish region updated the gasoil energy activity data from the residential sector. The fuel switch between gasoil and natural gas is now better estimated. This resulted in new emissions for the years 2002-2012.
- during the 2014 submission the Flemish region updated the wood consumption from the residential sector on the base of the Eurostat survey. Before the energy activity data were underestimated.
- The emission factors of the EMEP/EEA emission inventory guidebook 2013 (tier 2) will be used for wood burning in Flanders..
- In Brussels, AD for turbojets was reviewed and corrected. Three turbojets are reported between 1990 to 2008 (closure of one of it). From 2009 to 2013 two turbojets are reported.
- Improvement of reporting HAP; with the new EF, some NE for specific HAP were replaced by emissions.
- Replacement of NE with the emissions for sector 1A1c using the EF of the EMEP/EEA 2013 guidebook for: NH<sub>3</sub>, PM<sub>2.5</sub>, PM<sub>10</sub>, TSP, heavy metals, and POPs with the exception of HCB and PCBs (no EF in the guidebook).
- Replacement of NE for NH<sub>3</sub> for stationary sources using the EF used in Wallonia.
- Inclusion of the reporting of BC in the 3 region following the amended Gothenburg Protocol.

The following recalculations and improvements have been implemented for the 3 regions:

- Recalculation of all historic years (1990-2013) with
  - o New version of COPERT (v11.1)
  - o The inclusion of the COPERT option “Lube oil”
  - o The inclusion of scaling factors for the calculation of “fuel sold”
- A new fully harmonised and unique processing & reporting tool (processing of the COPERT results)

The improvements planned are:

- The emission factors of the EMEP/EEA emission inventory guidebook 2013 (tier 2) will be used for wood burning in Flanders and Wallonia.
- For some plants in Wallonia, the emission factors are not consistent throughout the time series. Indeed, from 2005, companies must report their emissions and these emissions are included in the inventory but in previous years, emission factors were sometimes used. For the next submission, emission factors will be calculated on the basis of company data (2005-2014) and used on the entire time series 1990-2004.
- In Flemish Region the EMMOSS model used to calculate the emissions of sea navigation will be revised for dredging, tugs and extraction of sand.
- For the road transport sector, recalculations are planned in the near future using :
  - o COPERT 4 v11.2 (version 11.1 has known errors regarding the emission factors of Euro VI HDV).
  - o Updated mobility figures for 2013 (at the time of the inventory calculation, the mobility for 2013 was not yet published, so the mobility figures of 2012 were used)

## ***9.2. Recalculations and improvements in the sector of industrial processes and products use***

- In Flanders the NMVOC emission of the other chemical industry (2B5a) is optimized by using more detailed information from the yearly reported integrated environmental reports. As a result the emissions of the polyurethane processing and polystyrene foam processing are now allocated under category 3C.
- In Flanders the emission inventory of Persistent Organic Pollutants (POPs) is optimized by VITO under the authority of VMM (Sleeuwaert et al., 2012).

In the Flemish region the following recalculations have been performed :

- 3C Chemical products:
  - o Polyester processing: The sector is allocated in 3C for the first time. The methodology has been modified. More detailed information from the yearly reported integrated environmental reports is used.
  - o Polyurethane processing: The sector is allocated in 3C for the first time. The methodology has been modified. More detailed information from the yearly reported integrated environmental reports is used.

- Polystyrene foam processing: The sector is allocated in 3C for the first time. The methodology has been modified. More detailed information from the yearly reported integrated environmental reports is used.
- Glues manufacturing: the AD of STATBEL from 2000 to 2011 have been modified due to an estimation of missing and incorrect figures.
- 3D Other Use of Solvents and Related Activities:
  - Printing industry: the emissions from 2009 to 2011 have been slightly modified due to corrections of figures in a couple of yearly integrated environmental reports.

In Wallonia, the following recalculations have been performed:

- 2H2 : modification of the NMVOC EF for bread (125 g/hab/year)
- 2D3d Coating applications:
  - Construction and buildings and domestic use: Emissions of the years 2011 and 2012 have been slightly modified because of an error in the AD figures.
  - Other industrial paint application: Emissions of 1990 have been modified to reflect the change in other sectors. Emissions from 2007 to 2012 have been modified to take into account the emissions determined on the basis of the yearly integrated environmental reports and the decrease of the global paint consumption between 2009 and 2013.
- 2D3e Degreasing:
  - Metal degreasing: Emissions from 2010 to 2012 have been modified to take into account the emissions determined on the basis of the yearly integrated environmental reports.
- 2D3g Chemical products:
  - Paints manufacturing: Emissions from 2010 to 2013 have been modified to reflect the decrease of the global paint consumption between 2009 and 2013.
  - Pharmaceutical products manufacturing: Emission of 2012 have been slightly actualized on the basis of new data provided by one plant.
- 2D3h Printing:
  - Printing industry: Emissions from 1990 to 2012 have been revised in order to ensure a coherence in the methodology for the all time series and to take into account the emissions determined on the basis of the yearly integrated environmental reports.
- 2D3i Other solvent use:
  - Preservation of wood: Emissions from 2010 to 2012 have been actualized on the basis of new activity data provided by one plant.

In the Flemish region, the following improvements are planned:

- Fine tuning the emissions of polyurethane processing;
- Fine tuning the emissions of polystyrene foam processing ;
- Estimation of the emissions of production of solvents: tetrachloromethane.

In Wallonia, the following improvements are planned:

- For some plants, the emission factors are not consistent throughout the time series. Indeed, from 2005, companies must report their emissions and these emissions are included in the inventory but in previous years, emission factors were sometimes used. For the next submission, emission factors will be calculated on the basis of company data (2005-2014) and used on the entire time series 1990-2004.
- The emission factors for PM10 and PM2.5 are not consistent for the time series as since 2005, the lime plants have performed PM10 analyses and have made an estimation of their PM2.5 emissions. Following information's coming from these plants, the size of the dust is high and there is very little fine dust. Before 2005, the proportion between PM10 and TSP was the proportion written in the EmeP Guidebook. A recalculation is planned to harmonise the proportion TSP/PM10/PM2.5 with plant data for the entire period.
- Estimation of the NMVOC emissions from the sector road paving with asphalt.
- Revision of the emissions for Wood paint application ;
- Revision of the emissions for non-chlorinated solvents for Metal degreasing, Dry cleaning and Other industrial cleaning;
- Revision of the emissions for Polyester processing, Polyvinylchloride processing, Polyurethane processing, Polystyrene processing;
- Estimation of the missing emissions (NE) for Textile finishing, Glass wool enduction, Mineral wool enduction and Domestic use of pharmaceutical products ;
- Revision of the emissions from key sources in order to move from Tier 1 to Tier 2 methodology when necessary

In Brussels region, the following recalculations have been taken place:

- The AD for bread and cookies production was corrected between 1990-2001 in Brussels database.
- There was a mistake on population data for sectors 2D3a, 2D3d and 2D3f which was corrected.
- The sector 2D3h has been updated with the latest environmental reports available.
- Sector 2D3e is being reported for the first time, information comes from annual environmental reports. The data is available from 2003. The emissions for the period 1990 – 2002 are kept constant and equal to the first available data.
- AD for car repairing (2D3d) and printing (2D3h) has been updated with new available statistics from the year 2012.

### **9.3. Recalculations and improvements in the agricultural sector**

In Flanders, during this submission February 2015, NH<sub>3</sub> emissions from 2012 are updated. In the submission of February 2014, NH<sub>3</sub>-emissions 2012 were reported as the same as NH<sub>3</sub> emissions from 2011. This was due to lack of activity data on the time of reporting.

Also in Flanders a new IEF of TSP, PM10 and PM2.5 for fattening pigs is used for the entire time series. This results in a decrease of the TSP and PM10 emission of  $\pm 30\%$  and for PM2.5 of  $\pm 50\%$ .

In Wallonia, recalculations occurred in the category 3B. Thanks to new information of a Walloon study (*Mathot & al, 2011*), the emission factor for the storage of solid manure of cattle has been reviewed for the entire time series. This results in a decrease of the emission of  $\pm 66\%$ .

Also in Wallonia, in category 3D, information about the application of slurry and the use of precise equipment (injectors), a reduction of emissions has been applied since 2003. In 2013, this measure results in a reduction of  $\pm 12\%$  of the emissions from manure application.

Finally, in Wallonia, the emission factor for the use of mineral fertilizer has been updated with the value of 2013 EMEP guidebook (Table 3-2, p 14 *Chapter 3.D*). This time, the result is roughly a doubling of the emissions of the category 3Da1.

In Flanders, an actualisation of the EMAN model is planned.

In Wallonia, no improvements are planned in the near future.

#### **9.4. Recalculations and improvements in the waste sector**

- The EF of the EMEP/EEA 2013 guidebook are applied for sectors 5C1a, 5C1biii and 5C1bv in Brussels.
- Brussels has replaced the NE for NH<sub>3</sub> for sectors 5C1a and 5C1biii, using the EF from the EMEP/EEA 2013 guidebook.
- Replacement of NE for Se, HAPs, HCB and PCBs with the EF of the EMEP/EEA 2013 guidebook for sectors 5C1biii and 5C1bv in Brussels.
- In Flanders the emissions of dioxins and HCB are re-allocated between 1A1a and 5C1. The emissions of the waste incineration plants with energy recovery are allocated to the category 1A1a.
- In the Flemish region, the following improvements are planned: It was not possible yet to split up the emissions between industrial/hazardous/sewage sludge and municipal waste incineration, hence the emissions due to municipal waste incineration without energy recycling are also allocated in category 5C1bi instead of 5C1a. A methodology will be developed to distinguish between the emissions of municipal and industrial waste incineration, and as far as possible between industrial, hazardous and sewage sludge incineration.
- In Wallonia, no improvements are planned in the near future.

## Chapter 10. Projections

Belgian emission projections are the sum of:

- Emission projections for the stationary sources in each of the three regions;
- Emission projections for mobile sources in Belgium

### 10.1. Mobile sources

Emission projections used for reporting in 2014 have been used. See IIR 2014 for details.

### 10.2. Stationary sources

#### 10.2.1. Flanders: Flemish energy and greenhouse gas simulation model

A new Flemish simulation model has been developed in 2014 to construct short term projections for Flanders. The simulation model is a projection model for energy demand, greenhouse gas emissions and emissions of air pollutants (SO<sub>2</sub>, NO<sub>x</sub>, PM and VOC) that covers most of the relevant emission sectors (energy sector, industry, waste, agriculture, residential and commercial buildings).

This simulation model works as a “bottom-up” type, i.e. explaining energy consumptions and emissions from activity variables expressed as far as possible in physical units, and the main determining factors of the evolution of energy demand and emissions.

The model, which includes a database on the energy consumption, emission factors, activity data and reduction effects of climate & energy and air quality policy measures, can be used in particular for:

- the construction of a reference scenario (business as usual), representing the expected future evolution in the absence of any new emission reduction policy based on expected economic and demographic evolutions;
- constructing emission reduction scenarios, based on the implementation of a combination of reduction measures;
- assessing the impact of existing or draft legislations on energy consumption and emission levels.

The model starts from reference year data:

- energy demand per industrial sector;
- emissions per industrial sector;
- large combustion plants and all electricity producing plants are included at installation level (energy consumption, electricity production and emissions);
- detailed information on the evolution of the installed power for electricity generation (including electricity import);
- a representation of the structure of the residential heating (type and age) and of residences (idem for the heating of tertiary buildings).
- Share of the emissions, per sector, that comes from processes (and thus is not related to fuel consumption).
- For the agricultural emissions (dust, greenhouse gasses and ammonia emissions coming from stables and from manure), the starting point is the number of animals (detailed per animal category and per type of stable) and the amount of manure that is spread out.

For the residential sector, projections are driven by assumptions on degree days in the future, the share of new residences and the lifetime of existing installations. Policies on energy efficiency and on ecodesign are taken into account.

For industry, major assumptions are the evolution of industrial activity and energy efficiency (yearly growth rate per sector), the share of CHP per sector and the lifetime of installations (since new

installations mostly can respect lower emission levels than the existing ones). This leads to a projection on energy consumption and electricity.

Electricity demand from all sectors (including transport) is the main driver for the electricity part of the model. The model searches for the most cost optimal mix of electricity generating installations (including import) to produce the necessary electricity, taking into account different time slices (electricity demand is not equal in winter and in summer, neither during night or day), based on production efficiencies and fuel cost. The model has the possibility to install additional production capacity (CCGT or gas turbine).

For all energy consuming sectors, energy consumption is translated into emission projections through emission factors (per fuel) that reflect policy (either current policy or additional measures). For industry and electricity production, current emission factors are compared to the emission factors based on policy and the lowest of both is used (installations that already comply with future emission standards don't need to realize additional reductions). For the residential sector, the emission factors take into account the use of different types of boilers and stoves.

For the agricultural sector, the predicted number of animals is multiplied with animal specific emission factors (both for the greenhouse gasses as for ammonia and dust). These emission factors are lower for the new low emission stables. Also the amount of manure that is spread out is multiplied with specific emission factors. At the time of reporting, final results were not yet available, so projections are likely to change soon.

## 10.2.2. Walloon region

### *SO<sub>2</sub> and NO<sub>x</sub>*

For Wallonia, projections of SO<sub>2</sub> and NO<sub>x</sub> emissions from stationary sources (combustion and processes) up to 2030 are similar to those reported in the previous issue. They have been established using the EPM model developed by ECONOTEC. This model is also a techno-economic bottom-up simulation model.

Projections are established on the basis of energy projections, in combination with the consideration of several abatement measures taken or in project in the Walloon industry, concerning the relevant air pollutants.

The energy projections are those reported in 2013 under Article 3(2)b of Decision No 280/2004/EC in the framework of the so-called "with measures scenario", ie a reference situation considering the impact of existing and implemented measures aiming at the reduction of greenhouse gases.

Newcomer in the present projections: NO<sub>x</sub> emissions from agriculture (for activity variables, see the contribution on NH<sub>3</sub> hereunder).

### *NM VOC*

Projections of NMVOC emissions have been established on the basis of the following assumptions:

- Energy sectors (1A1, 1A2, 1A4, 1A5): As the combustion of wood in the residential sector is the main contributor to the NMVOC emissions, the NMVOC emissions in 2030 have been calculated on the basis of the projection for wood consumption for the residential sector. The other emissions have been considered identical to 2013.
- Industrial processes (2A1, 2A2, 2B10a, 2C1, 2C7c, 2H1, 2H2): The complete shutdown of all blast furnaces has been taken into account. The emissions of the food and beverage sector

have been assumed to follow the evolution of population between 2013 and 2030. The other emissions have been considered identical to 2013.

- Solvent and other product use (2D, 2G): Most of the facilities falling under the Solvent Directive have already implemented the VOC reductions. When the activity data is related to population, the emissions have been assumed to follow the evolution of population between 2013 and 2030. For other sectors, the emissions have been considered identical to 2013.

New entry also: NMVOC emissions from agriculture & waste.

### *Agriculture*

The projections for agricultural emissions (NH<sub>3</sub>, NO<sub>x</sub>, NMVOC) are calculated with the same methodology described in chapter 6. The equations, the parameters and the emissions factors are the same. The activity data (heads of animals, crop areas and fertiliser use) are coming from the GAINS model.

### *Waste*

Concerning the projections of emissions from waste sector (NO<sub>x</sub>, NMVOC, SO<sub>2</sub> & NH<sub>3</sub>), the hypothesis followed is conservative and the emissions from the last submission are kept constant.

## 10.2.3. Brussels capital region

The Brussels Institute for Environmental Management has developed its own projection model for energy demand and atmospheric emissions from stationary sources (residential, tertiary, industry and heat and electricity sector). The horizon of the projections is 2030, after this year extrapolation for each sector has been done in order to estimate emissions for 2035.

As bottom-up type model, changes in consumption of the several energy carriers used in the Brussels-Capital Region (natural gas, light oil, propane/butane, coal, electricity, wood, solar and heat pump) and their associated emissions (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, NO<sub>x</sub>, CO, NMVOC, SO<sub>x</sub>, NH<sub>3</sub>, PM) are determined by the evolution of parameters that define the consumption of each sector.

For example, the residential sector is defined by the following main parameters :

- population and average household size (those 2 parameters define the net requirement for new dwellings)
- climate (expressed in degree-days, this parameter is of great importance for the Brussels-Capital Region, as it reflects the need for heating of buildings which represents about 70% of regional GHG direct emissions)
- renovation rate, improvement of energy efficiency expected in case of renovation and new constructions. This improvement depends on the typology of building stock composed of 140 type-dwellings : apartment or house, 5 age range of the building concerned, the 7 energy carrier used for heating, and the installed heating system (central or decentralised).

The model has been calibrated for each sector with the regional annual energy balances from 2000 to 2012. The modelled energy consumptions have then been converted into atmospheric emissions through emission factors, the ones used to establish the emission inventories.

The model also takes into account the GHG direct emissions that are not related to energy consumption: i.e. the fugitive methane emissions of natural gas delivery, the use of N<sub>2</sub>O for anesthesia and aerosols, the emissions from the decomposition of organic matter (composting plant, water purification plant).

This model is a dynamic one. It allows new future available data to be integrated (for instance future energy balances) as well as new assumptions reflecting new studies and new phenomena (in the fields of regulation, technological change, through awareness campaigns, incentives, or the evolution of energy costs, ...).

## **Chapter 11.        Reporting of gridded emissions and LPS**

No gridded emissions or LPS emissions were reported in the 2015 reporting cycle. The last submission of gridded and LPS data was made in 2012 on emissions of 2010.

## Chapter 12. Adjustments

*Please refer to the 'adjustment report 2015', submitted together with this IIR for more detailed information on the calculation of the adjustments. In this IIR, only the conclusions of the adjustment report are included.*

### 12.1. Adjustments - summary

Belgium signed and ratified the 1999 Protocol to Abate Acidification, Eutrophication and Ground-level ozone (Gothenburg Protocol) and Belgium as EU Member State adopted the National Emission Ceiling Directive (2001/81/EC) in 2001. Hereby, Belgium committed itself to reduce its emissions of NO<sub>x</sub>, SO<sub>2</sub>, NMVOC and NH<sub>3</sub> to the agreed national emission ceilings by 2010 and to respect these ceilings from 2010 onwards.

Table 1 summarizes the emission totals, based on **fuel used** for compliance under the NECD and the Gothenburg Protocol emission ceilings. Belgium exceeds its national emissions ceiling for NO<sub>x</sub> by 32% in 2010 against the NECD and by 28% against the Gothenburg Protocol ceiling. The provisional estimates for 2013 give an exceedance of 13% and 10% against NEC en Gothenburg respectively. For NMVOC, both emission ceilings were exceeded in 2010. Since 2011, the Gothenburg emission ceiling is met and since 2013 NMVOC emissions are below the NEC ceiling. The NEC and Gothenburg emission ceilings for the other pollutants were met in 2010 or earlier.

**Table 12-1: National non-adjusted total emissions (fuel used) and national emission ceilings.**

National Total (fuel used)*	NO <sub>x</sub>	NMVOC	SO <sub>x</sub>	NH <sub>3</sub>
<b>2010</b>	231.79	153.72	60.55	64.98
<b>2011</b>	216.92	141.49	53.17	64.17
<b>2012</b>	206.76	139.68	47.45	63.29
<b>2013</b>	199.49	136.68	45.55	62.16
<b>NEC Emission ceiling 2010</b>	<b>176</b>	<b>139</b>	<b>99</b>	<b>74</b>
<b>% above NEC ceiling in 2013</b>	13%	-2%	-54%	-16%
<b>Gothenburg Emission ceiling 2010</b>	<b>181</b>	<b>144</b>	<b>106</b>	<b>74</b>
<b>% above Gothenburg ceiling in 2013</b>	10%	-5%	-57%	-16%

*\*NOTE: The national totals (FU) reported under NECD (24/12/2014) slightly differ from the national totals reported under CLRTAP (resubmission 14/03/2015) due to little corrections made in the inventory. These changes do not affect the conclusions on compliance or non-compliance for the years 2010-2013.*

The non-compliance for NO<sub>x</sub> emissions up to 2013, as well as for NMVOC emissions in 2010 (Gothenburg Protocol) up to 2012 (NECD) are due to changes in the emission inventory, not foreseen at the time the emission ceilings were set. These changes include the partial failure of certain EURO vehicle emission standards, especially for diesel vehicles, as well as the inclusion of new source categories.

Based on EB decision 2012/4 allowing the provisional application of article 3, paragraph 11 quinquies of the amended Gothenburg Protocol, Belgium applies for making use of the adjustment procedure as described in Decision 2012/3 and according the Guidance in the consolidated version of the adapted Decision 2012/12 and additional Guidances for its emission inventory for NO<sub>x</sub> for 2010 to 2013 and NMVOC for 2010 to 2012 in order to prove its compliance with the 2010 NEC ceilings.

For **NO<sub>x</sub>**, the adjustment application to the emission inventory is the result of two (aggregated) adjustments, both of them in accordance with one of the circumstances as described in Decision 2012/12, article 6:

1. **Road transport (1A3bi-iv):** Significant change in emission factors (article 6(b))
2. **Agriculture (3B, 3Da1 and 3Da2a):** new source categories (article 6(a))

For **NMVOC**, the adjustment application to the emission inventory is due to the inclusion of the emissions from agricultural soils and manure management, two source categories that were not taken into account at the time the emission ceilings were set, and therefore in accordance with circumstance a) in Decision 2012/12:

**Agriculture (3B and 3De):** new source categories (article 6(a)).

**Table 12-2** summarizes the individual adjustments as well as the adjusted national total emissions. For compliance purposes, Belgium is allowed to use national total emissions based on **fuel used**<sup>7</sup> (EB2013, (4)). **With application of the adjustments, Belgium is in compliance with its NEC emission ceilings from 2010 on for all NEC pollutants. A fortiori, the Gothenburg Protocol emission ceilings are met.**

**Table 12-2: Total Emissions and adjustments for NO<sub>x</sub> and NMVOC.**

NO <sub>x</sub>	2010	2011	2012	2013	NEC Emission ceiling 2010	Gothenburg emission ceiling 2010
<b>National Total (fuel used)</b>	231.79	216.92	206.76	199.49	176	181
<b>Adjustment Road transport (1A3bi-iv)</b>	-47.98	-47.77	-47.17	-46.54		
<b>Adjustment Agriculture - Manure management (3B)</b>	-0.39	-0.38	-0.38	-0.37		
<b>Adjustment Agriculture - Inorganic N-fertilizers (3Da1)</b>	-5.94	-5.85	-5.64	-5.73		
<b>Adjustment Agriculture -Animal manure applied to soils (3Da2a)</b>	-7.60	-7.29	-7.07	-6.95		
<b>Adjusted national total for compliance*</b>	169.89	155.62	146.50	139.89		

<sup>7</sup> ECE/EB.AIR/125. Guidelines for Reporting Emissions and Projections Data under the Convention on Long-range Transboundary Air Pollution, Chapter V, A.23.

NMVOC	2010	2011	2012	2013	NEC Emission ceiling 2010	Gothenborg emission ceiling 2010
National Total (fuel used)	153.72	141.49	139.68	136.68	139	144
Adjustment Agriculture - Manure management (3B)	-36.59	-35.90	-35.60	NA		
Adjustment Agriculture - Cultivated crops (3De)	-1.22	-1.20	-1.19	NA		
Adjusted national total for compliance*	115.91	104.40	102.90	NA		

*\*IMPORTANT NOTE: adjusted national totals for compliance reported under NECD (24/12/2014) were lower due to the inclusion of an adjustment for the off road sector. After the NEC submission Belgium decided not to apply for an adjustment of the off road sector. Please refer to the adjustments calculated in the LRTAP resubmission 14/3/2015 and described in this report for the right adjustment under NECD and LRTAP.*

## References

- Arcadis (2010). NMVOC emissions through domestic solvent use and the use of paints in the Brussels Capital Region.
- Bartelink, H.H. (1997). Allometric relationships for biomass and leaf area of beech (*Fagus sylvatica* L) . *Annales des Sciences Forestières*, 54 pp 39-50.
- Bauwens, S. (2010). Inventaire sur l'affectation des terres et du changement d'affectation des terres et la foresterie (LULUCF) de la Belgique, Rapport intermédiaire 2 février 2010 Gemboux Agro Biotech.
- Bauwens, S. (2011). Inventaire sur l'affectation des terres et du changement d'affectation des terres et la foresterie (LULUCF) de la Belgique, Rapport final Mai 2011 Gemboux Agro Biotech.
- Bauwens, S. (2011). Optimalisatie emissie-inventaris lucht : sector landgebruik en verandering in landgebruik, Final Report, December 2011. Gemboux Agro Biotech.
- Beerparadise. URL via [www.beerparadise.be](http://www.beerparadise.be)
- Belkacem, S., Nys, C. & Gelhaye, D. (1992). Effets d'une fertilisation et d'un amendement sur l'immobilisation d'éléments dans la biomasse d'un peuplement adulte d'épicéa commun (*Picea abies* L Karst). *Annales des Sciences Forestières*, 49, pp 235-252.
- Bogaert, S., Devoldere, K., Van Hyfte, A., Van Biervliet, K. & Le Roy D. (2004). Evaluatie van het reductiepotentieel voor diverse pollutiemissies naar het compartiment lucht in een aantal homogene subsectoren van de chemische industrie in Vlaanderen, deel III. Aminoal – Aminabel. 442p.
- Bonneau (1995). La fertilisation en forêt.
- Campens, V. & Lauwers, L. (2002). Kunstmestgebruik en gewasproductie als determinanten van de nutriëntenemissie, uitgevoerd door de VMM en CLE.
- D'Haene, V., Van Hyfte, A. & Van Langenhove, H. (2002). Emissies van vluchtige organische stoffen in Vlaanderen: verfijning van de inventarisatie en van het relationeel verband met troposferische ozon. Universiteit Gent, Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen, Vakgroep Organische Chemie. Studie uitgevoerd in opdracht van de Vlaamse Milieumaatschappij, onderzoeksopdracht nr. VMM.AMO.2000, 346 p.
- D'Haene, V., Van Hyfte, A. & Van Langenhove, H. (2002). Emissies van vluchtige organische stoffen in Vlaanderen: verfijning van de inventarisatie en van het relationeel verband met troposferische ozon. Study performed by Ghent University under the authority of the Flemish Environment Agency, VMM.AMO.2000, 346 p.
- De Roo, K., Philips, G. & Van Durme, J. (2009). Optimalisatie emissie-inventaris vluchtige organische stoffen van sectoren 'coating', 'droogkuis' en 'reinigen en ontvetten'. Studie uitgevoerd door de Universiteit Gent in samenwerking met PRG Odournet in opdracht van de VMM. Ref. G08/VMM/11, 133p.
- Denier van der Gon, H. & Hulskotte, J. (2010). Methodologies for estimating shipping emissions in the Netherlands; a documentation of currently used emission factors and related activity data. Study conducted under the auspices of the Netherlands Research Program on Particulate Matter (BOP), 56 p. URL: <http://www.rivm.nl/bibliotheek/rapporten/500099012.pdf>
- DNV, CCI, TNO (2014). Uncertainty analysis of emission inventories of NEC/LRTAP air pollutants. Final report, 115p.

Duvigneaud, P., Denaeyer, S. & Kestemont, P. (1977). Productivité primaire de forêts belges de types variés. Productivité biologique en Belgique, Travaux de la section belge du Programme Biologique International. Editions Duculot, Paris-Gembloux.

ECE/EB.AIR/125 (2013). Guidelines for Estimating and Reporting Emission Data under CLRTAP.

Available via

[http://www.ceip.at/fileadmin/inhalte/emep/2014\\_Guidelines/ece.eb.air.125\\_ADVANCE\\_VERSION\\_reporting\\_guidelines\\_2013.pdf](http://www.ceip.at/fileadmin/inhalte/emep/2014_Guidelines/ece.eb.air.125_ADVANCE_VERSION_reporting_guidelines_2013.pdf)

Econotec (1997), Inventaire 1996 des émissions atmosphériques en région wallonne-Groupe 2, 5 et 6 de la nomenclature Snap.

Econtec (2010). Study on dust : analyse prévisionnelle des émissions de poussières en région wallonne à l'horizon 2010, juillet 2010.

EPA (1996), AP42-Hot mix asphalt plant.

European Environment Agency (2013). Emep/EEA air pollutant emission inventory guidebook – 2013.

EXcoser : Quantification des émissions d'HAP dans l'environnement en Belgique - Rapport final, Excoser SA pour la DGRNE, 30/5/95.

Essen (1997), Identification of relevant industrial sources of dioxins and furans in Europe.

Foqué D. & Demeyer P. (2009). Optimalisering en actualisering van de emissie-inventaris ammoniak landbouw. Instituut voor Landbouw- en Visserijonderzoek. Studie uitgevoerd in opdracht van de Vlaamse Milieumaatschappij. Mededeling ILVO nr 69, 147 p.

FPS Health. Brochure 'CCIM – één unieke stem op de internationale milieuscène'. Available via URL <http://www.health.belgium.be/eportal/Environment/Inspectionandenvironmentalrigh/international/CCPIE/index.htm?fodnlang=nl>

Gloaguen, J.-C. & Touffet, J. (1982). Production de litière dans une chênaie-hêtraie atlantique. Relations avec les caractères climatiques. Revue Forestière française, XXXIV, pp108-118.

Guns, A. (1990). Effets à moyen terme d'un apport de fertilisants sur une jeune plantation d'épicéa commun (*Picea abies* L Karst). Travail de fin d'études. Section interfacultaire d'agronomie, Université Libre de Bruxelles.

Hewitt C.N., Street, R.A. & Scholefield P.A. (1997). Isoprene and monoterpene-emitting species survey 1997. <http://www.es.lans.ac.uk/es/people/pg/pas/download.html>

Klein, J., Hoen, A., Hulskotte, J., Van Duynhoven, N., Smit, R., Hensema, A. & Broeckhuizen, O. (2006). Methoden voor de berekening van de emissies door mobiele bronnen in Nederland.

Van Mierlo, J., Boureima, F., Messagie, M., Sergeant, N., Govaerts, L., Denys, T., Michiels, H., Vernaillen, S., Schrooten, L., Beckx, C., Macharis, C., Turcksin, L., Bernardini, A., Hecq, W., Klopfert, F., Englert, M., De Caemel, B. & De Vos, M. (2009). CLEAN VEHICLE RESEARCH: LCA AND POLICY MEASURES 'CLEVER' - Final Report. Brussels : Belgian Science Policy 2009 – 119 p. (Research Programme Science for a Sustainable Development).

Lodewijks, P., Van Rompaey, H. & Sleeuwaert, F. (2003). VOS-emissies naar de lucht bij de productie en het industrieel gebruik van coatings, inkt en lijm in Vlaanderen. Studie uitgevoerd door de VITO in opdracht van AMINAL ref. nr. 2003/IMS/R155, 359 p.

Lodewijks, P., Polders, C. & Van Rompaey, H. (2005). Evaluatie van de inschatting van NMVOS emissies door verbrandingsprocessen in Vlaanderen. Study performed by VITO under the authority of Aminal, 2005/IMS/R/323, 132 p.

Luchetta, L., Simon, V. & Torres, L. (2000). Emission des principaux composés organiques volatils biogéniques en France. *Pollution Atmosphérique*, n° 167.

Meulepas P., Vercaemst P., *Beste Beschikbare Technieken voor de benzinestations*, VITO, 1999/PPE/P/103, 233pg

MNZ (1995), Quantification des émissions d'hydrocarbures aromatiques polycycliques (HAP) dans l'environnement en Belgique.

Neiryneck, J., Maddelein, D., de Keersmaecker, L., Lust, N. & Muys, B. (1998). Biomass and nutrient cycling of a highly productive Corsican pine stand on former heathland in northern Belgium. *Annales des Sciences Forestières*, 55, pp 389-405.

NIR (2014). Belgium's greenhouse gas inventory (1990-2012). Available via [http://unfccc.int/files/national\\_reports/annex\\_i\\_ghg\\_inventories/national\\_inventories\\_submissions/application/zip/bel-2014-nir-10apr.zip](http://unfccc.int/files/national_reports/annex_i_ghg_inventories/national_inventories_submissions/application/zip/bel-2014-nir-10apr.zip)

NIS (2010). Belgium's National Inventory System for the estimation of anthropogenic greenhouse gas emissions by sources and removals by sinks under Article 5, paragraph 1, of the Kyoto Protocol, 45 p.

OVAM (2008). Tarieven en capaciteiten voor storten en verbranden - Actualisatie tot 2006, evolutie en prognose. D/2007/5024/79. 69 p.

Parcom (1992), Emission factors manual.

Polders, C., Hooyberghs, E., Vanassche, S. & Huybrechts, D. (2011). *Beste Beschikbare Technieken (BBT) voor de houtverwerkende nijverheid*. Gent, Academia Press, xiv + 635 p.

Polders C., Wevers M. & Van Rompaey H. (2003). Haalbaarheidsonderzoek en aanzet tot de opmaak van een emissie-inventaris van dioxines. Studie uitgevoerd door de VITO in opdracht van VMM, ref. 2003/IMS/R/006, 107 p.

Ponette, Q. & Ranger, J. (2000). Biomasses et minéralomasses aériennes de cinq peuplements de douglas du réseau Rénécofor : quantification et implications sylvicoles. *Revue Forestière française*, LII-2, pp115-133.

Ranger, J., Cuirin, G., Bouchon, J., Colin, M., Gelhaye, D. & Mohamed Ahamed, D. (1992). Biomasse et minéralomasse d'une plantation d'épicéa commun (*Picea abies* L Karst) de forte production dans les Vosges. *Annales des Sciences Forestières*, 43, pp 651-668.

Ranger, J., Marques, R., Colin-Belgrand, M., Flammang, N. & Gelhaye, D. (1996). La dynamique d'incorporation d'éléments nutritifs dans un peuplement de douglas. Conséquences pour la gestion sylvicole. *Revue Forestière française*, XLVIII-3, pp217-230.

Renders, N., Duerinck, J., Altdorfer, F. & Baillot, Y. (2010). Potentiële emissiereducties van de verwarmingssector tegen 2030. Study performed by VITO and Econotec under the authority of the Federal Public Service Health, food chain safety and environment, 2010/TEM/R, 256 p.

Schrooten, L. & Van Rompaey, H. (2002). Ontwikkeling van een methodologie voor een emissie-inventaris van PM10 en PM2,5 en opstellen van een emissie-inventaris voor 1995 en 2000. Study performed by VITO under the authority of VMM, 2002/IMS/R/200, 246 p.

Schrooten, L., Jespers, K., Baetens, K., Van Esch, L., Gijsbers, M., Van Linden, V. & Demeyer, P. (2009). OFFREM. Model voor emissies door niet voor de weg bestemde mobiele machines. Study performed by ILVO and VITO under the authority of Environment, Nature and Energy Department of the Flemish Government (2009/TEM/R). 133 p.

Schrooten, L., Van Rompaey, H., Berghmans, P., Vanderreydt, I. & Bleux, N. (2002). Deel A: Emissie-inventaris fijn stof Vlaanderen voor 1995 en 2000. Deel B: Monitoring fijn stof emissies voor industriële bronnen. Study performed by VITO under the authority of VMM. 2002/IMS/R/200.

Sleeuwaert, F., Polders C. & Van Rompaey, H. (2006). Optimalisatie en actualisatie van de emissie-inventaris fijn stof in het kader van internationale ontwikkelingen. Study performed by VITO under the authority of VMM, 2006/IMS/R/391, 140 p.

Sleeuwaert F., Van Esch L. & Engelen G. (2012). Ontwikkelen en optimalisatie van een emissie-inventaris Persistente Organische Polluenten (POP's). Study performed by VITO in cooperation with TNO (Coenen P., Visschedijk A., Van der Gon H. & Hulskotte J.) under the authority of VMM. 2012/MRG/R/389, 163 p.

Sleeuwaert, F., Van Esch, L., Jespers, K., Van Rompaey H. & Engelen, G. (2010). Actualisering en optimalisering van de inschatting van de verbrandingsemissies door de collectief geregistreerde bedrijven. Study performed by VITO under the authority of VMM, 2010/MRG/R/352, 88 p.

Teller, A. (1983). Biomasse, productivité et évaluation des déchets dans un peuplement d'épicéa à Strainchamps (Ardennes) Travail de fin d'études. Section interfacultaire d'agronomie, Université Libre de Bruxelles.

ULG (1998). Inventaire des émissions atmosphériques en Région Wallonne pour 1996, Université de Liège, juillet 1998.

Van Hyfte, A. & Van Langenhove, H. (2000). Emissies van vluchtige organische stoffen in Vlaanderen: verfijning van de inventarisatie en van het relationeel verband met troposferische ozonvorming. Programma Beleidsgericht Onderzoek 1997, project PB097/29/160, 190 p.

Van Ransbeeck, N. (2013). Particulate matter, ammonia and greenhouse gases in pig fattening facilities: measuring strategies, indoor concentrations and emissions. Universiteit Gent, 165 p.

Van Rompaey, H., De Fré, R., De Spiegeleer, E., Polders, C., Vanderstraeten, P. & Wevers, M. (2001). Emissies van dioxins en PAK's door gebouwenverwarming met vaste brandstoffen. Study performed by VITO under the authority of AMINAL. 2001/IMS/R/059, 208 p.

Van Rompaey, H. & Wuyts, H. (1999). Collectieve registratie van industriële emissies. Studie uitgevoerd in opdracht van de Vlaamse Milieumaatschappij, project 1999/PPE/R/153, 55 p.

Vanherle, K, Van Zeebroeck, B. & Hulskotte, J. (2007). Emissiemodel voor spoorverkeer en scheepvaart in Vlaanderen: EMMOSS. Study performed by Transport and Mobility Leuven (TML) under the authority of VMM. 100 p. URL: <http://www.tmlleuven.be/project/emmos/index.htm>

Vanherle, K, Vanhove, F., Spitaels, K, Carlier, K. (2010). Actualisering en verfijning van het emissiemodel voor spoorverkeer en scheepvaart in Vlaanderen (EMMOSS). 85 p.

Wevers, M. (2002). Meetcampagnes dioxines en PAK's: emissies door open vuren, open tonnetjes, kachels en allesbranders en analyses van roetstalen uit schouwen. Study performed by VITO under the authority of AMINAL/Aminabel.

## Annex 1: Key category analysis

### 1.A Level assessment

The following tables show the level assessment for 2013 for all reported pollutants.

NOx Source Code	Source Category	Gg NO <sub>2</sub>	Level ass.	2013 Cum.Total
1A3bi	Road transport: Passenger cars	46.275	22.5%	22.5%
1A3biii	Road transport: Heavy duty vehicles and buses	40.621	19.8%	42.3%
1A4bi	Residential: Stationary	11.868	5.8%	48.1%
1A3bii	Road transport: Light duty vehicles	10.692	5.2%	53.3%
1A1a	Public electricity and heat production	10.34	5.0%	58.4%
2A1	Cement production	8.829	4.3%	62.7%
3Da2a	Animal manure applied to soils	5.654	2.8%	65.4%
2B10a	Chemical industry: Other (please specify in the IIR)	5.516	2.7%	68.1%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	5.027	2.4%	70.5%
3Da1	Inorganic N-fertilizers (includes also urea application)	4.73	2.3%	72.8%
2C1	Iron and steel production	4.264	2.1%	74.9%
1A4ai	Commercial/institutional: Stationary	4.027	2.0%	76.9%
1A3dii	National navigation (shipping)	4.022	2.0%	78.8%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	3.881	1.9%	80.7%

NM VOC Source Code	Source Category	Gg NM VOC	Level ass.	2013 Cum.Total
3B1b	Manure management - Non-dairy cattle	17.903	13.0%	13.0%
1A4bi	Residential: Stationary	16.971	12.3%	25.4%
2D3a	Domestic solvent use including fungicides	14.888	10.8%	36.2%
2D3d	Coating applications	12.308	9.0%	45.2%
2B10a	Chemical industry: Other (please specify in the IIR)	9.687	7.0%	52.2%
3B1a	Manure management - Dairy cattle	8.304	6.0%	58.3%
2D3g	Chemical products	4.533	3.3%	61.6%
3B3	Manure management - Swine	4.132	3.0%	64.6%
1A3bi	Road transport: Passenger cars	3.448	2.5%	67.1%
1B2aiv	Fugitive emissions oil: Refining / storage	3.11	2.3%	69.3%
1B2b	Fugitive emissions from natural gas (exploration, production, processing, transmission, storage, distribution and other)	3.075	2.2%	71.6%
2H2	Food and beverages industry	2.971	2.2%	73.7%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	2.66	1.9%	75.7%
3B4gii	Manure management - Broilers	2.514	1.8%	77.5%
2D3h	Printing	2.438	1.8%	79.3%
1A3bv	Road transport: Gasoline evaporation	2.359	1.7%	81.0%

<b>SOx</b>				<b>2013</b>
<b>Source Code</b>	<b>Source Category</b>	<b>Gg SO<sub>2</sub></b>	<b>Level ass.</b>	<b>Cum.Total</b>
1A4bi	Residential: Stationary	8.468	18.6%	18.6%
1A1b	Petroleum refining	6.831	15.0%	33.6%
2C1	Iron and steel production	6.024	13.2%	46.8%
2B10a	Chemical industry: Other (please specify in the IIR)	3.71	8.1%	54.9%
2A1	Cement production	3.199	7.0%	62.0%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	2.585	5.7%	67.6%
1A1a	Public electricity and heat production	1.966	4.3%	72.0%
1B2c	Venting and flaring (oil, gas, combined oil and gas)	1.791	3.9%	75.9%
2A6	Other mineral products (please specify in the IIR)	1.756	3.9%	79.7%
1A4ai	Commercial/institutional: Stationary	1.256	2.8%	82.5%

<b>NH3</b>				<b>2013</b>
<b>Source Code</b>	<b>Source Category</b>	<b>Gg NH3</b>	<b>Level ass.</b>	<b>Cum.Total</b>
3B3	Manure management - Swine	19.962	32.1%	32.1%
3B1b	Manure management - Non-dairy cattle	7.411	11.9%	44.0%
3Da1	Inorganic N-fertilizers (includes also urea application)	7.234	11.6%	55.6%
3Da2a	Animal manure applied to soils	6.632	10.7%	66.3%
3Da3	Urine and dung deposited by grazing animals	5.38	8.6%	74.9%
3B1a	Manure management - Dairy cattle	4.834	7.8%	82.7%

<b>PM2.5</b>				<b>2013</b>
<b>Source Code</b>	<b>Source Category</b>	<b>Gg PM10</b>	<b>Level ass.</b>	<b>Cum.Total</b>
1A4bi	Residential: Stationary	20.725	61.9%	61.9%
1A3bi	Road transport: Passenger cars	1.399	4.2%	66.1%
1A3bvi	Road transport: Automobile tyre and brake wear	1.074	3.2%	69.3%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.85	2.5%	71.9%
2A5a	Quarrying and mining of minerals other than coal	0.8	2.4%	74.3%
1A3biii	Road transport: Heavy duty vehicles and buses	0.726	2.2%	76.4%
1A3bii	Road transport: Light duty vehicles	0.708	2.1%	78.6%
1A2gviii	Stationary combustion in manufacturing industries and construction: Other (please specify in the IIR)	0.637	1.9%	80.5%

<b>PM10</b>				<b>2013</b>
-------------	--	--	--	-------------

Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
1A4bi	Residential: Stationary	21.255	49.4%	49.4%
2A5a	Quarrying and mining of minerals other than coal	2.376	5.5%	54.9%
1A3bvi	Road transport: Automobile tyre and brake wear	2.002	4.7%	59.6%
1A3bi	Road transport: Passenger cars	1.399	3.3%	62.8%
3B4gii	Manure mangement - Broilers	1.395	3.2%	66.1%
3B4gi	Manure mangement - Laying hens	1.333	3.1%	69.2%
3Da1	Inorganic N-fertilizers (includes also urea application)	1.113	2.6%	71.8%
1A3bvii	Road transport: Automobile road abrasion	1.096	2.5%	74.3%
3B3	Manure management - Swine	0.974	2.3%	76.6%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.874	2.0%	78.6%
2C1	Iron and steel production	0.791	1.8%	80.5%

TSP				2013
Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
1A4bi	Residential: Stationary	22.425	39.8%	39.8%
2A5a	Quarrying and mining of minerals other than coal	6.95	12.3%	52.1%
1A3bvi	Road transport: Automobile tyre and brake wear	2.64	4.7%	56.8%
1A3bvii	Road transport: Automobile road abrasion	2.191	3.9%	60.7%
3B3	Manure management - Swine	2.145	3.8%	64.5%
1A3bi	Road transport: Passenger cars	1.399	2.5%	67.0%
3B4gii	Manure mangement - Broilers	1.395	2.5%	69.4%
3B4gi	Manure mangement - Laying hens	1.333	2.4%	71.8%
1A2gvii	Mobile Combustion in manufacturing industries and construction: (please specify in the IIR)	1.152	2.0%	73.9%
3Da1	Inorganic N-fertilizers (includes also urea application)	1.113	2.0%	75.8%
2C1	Iron and steel production	0.95	1.7%	77.5%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.928	1.6%	79.2%
2B10a	Chemical industry: Other (please specify in the IIR)	0.826	1.5%	80.6%

BC				2013
Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
1A4bi	Residential: Stationary	2.306	41.6%	41.6%
1A3bi	Road transport: Passenger cars	1.106	20.0%	61.6%
1A3bii	Road transport: Light duty vehicles	0.551	10.0%	71.5%
1A3biii	Road transport: Heavy duty vehicles and buses	0.499	9.0%	80.5%

CO				2013
----	--	--	--	------

Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
2A2	Lime production	170.365	30.8%	30.8%
1A4bi	Residential: Stationary	130.234	23.5%	54.4%
2C1	Iron and steel production	112.79	20.4%	74.7%
1A3bi	Road transport: Passenger cars	39.623	7.2%	81.9%

<b>Pb</b>				
Source Code	Source Category	Gg PM10	Level ass.	2013 Cum.Total
2C1	Iron and steel production	14.383	54.5%	54.5%
1A3bvi	Road transport: Automobile tyre and brake wear	2.277	8.6%	63.1%
2C7c	Other metal production (please specify in the IIR)	2.154	8.2%	71.3%
1A4bi	Residential: Stationary	1.193	4.5%	75.8%
1B2aiv	Fugitive emissions oil: Refining / storage	1.087	4.1%	79.9%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	1.035	3.9%	83.9%

<b>Cd</b>				
Source Code	Source Category	Gg PM10	Level ass.	2013 Cum.Total
2C7c	Other metal production (please specify in the IIR)	0.809	34.9%	34.9%
1A4bi	Residential: Stationary	0.336	14.5%	49.4%
1B2aiv	Fugitive emissions oil: Refining / storage	0.221	9.6%	59.0%
2C1	Iron and steel production	0.221	9.6%	68.5%
1A1a	Public electricity and heat production	0.148	6.4%	74.9%
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	0.08	3.4%	78.3%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.078	3.4%	81.7%

<b>Hg</b>				
Source Code	Source Category	Gg PM10	Level ass.	2013 Cum.Total
2B10a	Chemical industry: Other (please specify in the IIR)	0.305	18.4%	18.4%
2A1	Cement production	0.238	14.4%	32.8%
1B2aiv	Fugitive emissions oil: Refining / storage	0.232	14.0%	46.7%
1A1a	Public electricity and heat production	0.172	10.4%	57.1%
2C1	Iron and steel production	0.161	9.7%	66.8%
1A4bi	Residential: Stationary	0.124	7.5%	74.3%
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.067	4.0%	78.3%
2C7c	Other metal production (please specify in the IIR)	0.055	3.3%	81.6%

<b>As</b>				
Source Code	Source Category	Gg PM10	Level ass.	2013 Cum.Total

Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
2C7c	Other metal production (please specify in the IIR)	0.619	38.2%	38.2%
1A1a	Public electricity and heat production	0.208	12.8%	51.0%
2C1	Iron and steel production	0.166	10.3%	61.3%
2A1	Cement production	0.112	6.9%	68.1%
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.093	5.7%	73.9%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.071	4.4%	78.3%
1B2aiv	Fugitive emissions oil: Refining / storage	0.053	3.3%	81.5%

Cr				2013
Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
2C1	Iron and steel production	1.337	22.5%	22.5%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	1.024	17.3%	39.8%
1A3bvi	Road transport: Automobile tyre and brake wear	0.835	14.1%	53.9%
1A4bi	Residential: Stationary	0.65	11.0%	64.9%
1A1a	Public electricity and heat production	0.516	8.7%	73.6%
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	0.199	3.4%	76.9%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.193	3.3%	80.2%

Cu				2013
Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
1A3bvi	Road transport: Automobile tyre and brake wear	18.221	58.3%	58.3%
1A3c	Railways	5.001	16.0%	74.3%
2G	Other product use (please specify in the IIR)	2.91	9.3%	83.6%

Ni				2013
Source Code	Source Category	Gg PM10	Level ass.	Cum.Total
1A3bvi	Road transport: Automobile tyre and brake wear	2.277	33.6%	33.6%
2C1	Iron and steel production	0.935	13.8%	47.4%
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.448	6.6%	54.0%
1B2aiv	Fugitive emissions oil: Refining / storage	0.372	5.5%	59.5%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.337	5.0%	64.5%
1A1a	Public electricity and heat production	0.313	4.6%	69.1%
1A1b	Petroleum refining	0.217	3.2%	72.3%
2B10a	Chemical industry: Other (please specify in the IIR)	0.179	2.6%	74.9%
2A1	Cement production	0.17	2.5%	77.4%
1A2b	Stationary combustion in manufacturing industries and construction: Non-ferrous metals	0.16	2.4%	79.8%
2C7c	Other metal production (please specify in the IIR)	0.158	2.3%	82.1%

Se		Gg PM10	Level ass.	2013
Source Code	Source Category			Cum.Total
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	1.629	46.2%	46.2%
1A1a	Public electricity and heat production	0.775	22.0%	68.2%
2A1	Cement production	0.496	14.1%	82.3%

Zn		Gg PM10	Level ass.	2013
Source Code	Source Category			Cum.Total
2C7c	Other metal production (please specify in the IIR)	18.133	22.4%	22.4%
1A4bi	Residential: Stationary	13.885	17.2%	39.6%
1A3bi	Road transport: Passenger cars	8.666	10.7%	50.3%
1A3bvi	Road transport: Automobile tyre and brake wear	8.468	10.5%	60.8%
2C1	Iron and steel production	7.532	9.3%	70.1%
1A3biii	Road transport: Heavy duty vehicles and buses	3.86	4.8%	74.9%
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	3.384	4.2%	79.1%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	3.383	4.2%	83.2%

PCDD/F		Gg PM10	Level ass.	2013
Source Code	Source Category			Cum.Total
1A4bi	Residential: Stationary	22.875	53.1%	53.1%
5C2	Open burning of waste	8.366	19.4%	72.5%
2C1	Iron and steel production	3.874	9.0%	81.5%

PAHs Total1-4		Gg PM10	Level ass.	2013
Source Code	Source Category			Cum.Total
2D3a	Domestic solvent use including fungicides	12.114	43.4%	43.4%
1A4bi	Residential: Stationary	10.133	36.3%	79.6%
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	4.043	14.5%	94.1%

HCB		Gg PM10	Level ass.	2013
Source Code	Source Category			Cum.Total
2L	Other production, consumption, storage, transportation or handling of bulk products (please specify in the IIR)	5.484	38.9%	38.9%
1A1a	Public electricity and heat production	3.839	27.2%	66.1%
2A1	Cement production	2.502	17.7%	83.8%

PCB		Gg PM10	Level ass.	2013
Source Code	Source Category			Cum.Total
2A1	Cement production	2.674	56.8%	56.8%
2C1	Iron and steel production	1.44	30.6%	87.3%

The following tables show the results of the key category analysis (level assessment) per year.

1990	Key source categories (sorted from high to low, from left to right)												Total
NO <sub>x</sub>	1A3bi	1A3bii	1A1a	1A4bi	2A1	1A2a	3Da2a	1A2f	1A1b	1A3bii	3Da1		81.3%
	27.1%	17.9%	14.8%	4.0%	3.6%	3.3%	2.3%	2.3%	2.2%	2.2%	1.7%		
NMVOC	1A3bi	2D3d	2B10a	3B1b	1A3bv	1B2aiv	3B1a	2D3h	2D3g	1A4bi	1B2av	2D3a	81.2%
	19.0%	14.6%	8.7%	6.3%	5.2%	5.0%	4.4%	4.1%	4.0%	3.8%	3.3%	2.8%	
SO <sub>x</sub>	1A1a	1A1b	1A4bi	1A4ci	2B10a	1A2f	1A2c	1A2e	1A2a	2C1			80.1%
	26.0%	11.2%	8.5%	7.8%	6.1%	4.8%	4.6%	4.4%	3.8%	2.9%			
NH <sub>3</sub>	3B3	3B1b	3B1a	3Da3	3Da2a								81.3%
	42.5%	13.2%	10.1%	7.8%	7.7%								
CO	1A3bi	2C1	1A2a	1A4bi									88.2%
	40.9%	19.0%	18.5%	9.7%									
Pb	1A3bi	2C1	5C1bi	2C7c									88.4%
	37.7%	19.9%	19.1%	11.6%									
Cd	2C7c	5C1bi	2C1	1A1a									82.7%
	35.2%	26.3%	15.0%	6.2%									
Hg	5C1bi	1A1a	2B10a	2A1									84.0%
	37.6%	23.3%	17.3%	5.7%									
As	2C1	1A1a	2C7c										80.5%
	32.8%	31.2%	16.5%										
Cr	2C1	5C1bi	1A1a										81.3%
	43.4%	29.2%	8.6%										
Cu	1A3bvi	2C1	1A1a	5C1bi	2C7c	1A2b							82.9%
	35.0%	17.2%	9.3%	8.6%	6.7%	6.2%							
Ni	1A2c	1A2e	5C1bi	2C1	1A2gvi	1A1a							87.0%
	22.1%	19.8%	12.4%	11.5%	10.9%	10.5%							
Se	1A1a	1A2f	2A2										86.5%
	53.9%	25.5%	7.1%										
Zn	2C1	5C1bi	2C7c	2B10a	1A2f	1A1a							81.7%
	19.8%	18.8%	17.3%	14.5%	6.6%	4.8%							
PCDD/ PCDF	2C1	1A1a	5C1a	5C1bii	2C7c								84.3%
	25.8%	25.7%	16.1%	11.7%	5.0%								
PAH	1B1b	2C1	1A4bi										82.8%
	38.7%	28.7%	15.4%										
HCB	1A2gvii	2L	1A1a	2C1									81.9%
	30.1%	28.2%	14.1%	9.6%									
PCB	2C1												93.7%
	93.7%												

1995	Key source categories (sorted from high to low, from left to right)													Total
NO <sub>x</sub>	1A3bi	1A3biii	1A1a	1A4bi	2A1	1A2f	1A2a	3Da2a	1A3bii	1A2c	1A1b	1A4cii		80.9%
	21.2%	19.1%	14.7%	4.8%	4.4%	3.3%	3.0%	2.5%	2.3%	2.1%	1.8%	1.8%		
NMVOC	1A3bi	2D3d	3B1b	2B10a	1B2aiv	2D3g	1A4bi	3B1a	2D3h	2D3a	1A3bv	1B2av	3B3	80.2%
	15.1%	14.4%	8.5%	6.9%	5.3%	4.3%	4.2%	4.2%	4.1%	4.1%	4.0%	3.3%	1.7%	
SO <sub>x</sub>	1A1a	1A1b	1A4bi	2B10a	1A2e	1A4ci	2C1	1A2f	1A2c	1A2a	1A3bi			80.2%
	30.3%	13.5%	9.3%	4.4%	4.2%	3.4%	3.3%	3.2%	3.2%	3.0%	2.3%			
NH <sub>3</sub>	3B3	3B1b	3Da3	3Da2a	3B1a	3Da1								86.7%
	43.4%	13.8%	8.2%	7.5%	7.1%	6.8%								
CO	1A3bi	2C1	1A2a	1A4bi										82.4%
	34.5%	22.3%	15.4%	10.3%										
Pb	1A3bi	2C1	2C7c	1A2f	5C1bi									86.3%
	29.8%	22.3%	17.7%	9.9%	6.5%									
Cd	2C7c	2C1	1A2f	1A1a	5C1bi									83.2%
	46.7%	15.7%	8.9%	6.0%	5.9%									
Hg	1A1a	2B10a	2A1	5C1a	2C1	5C1bi								85.4%
	29.4%	19.2%	11.4%	9.0%	8.8%	7.5%								
As	2C7c	2C1	1A1a											83.5%
	39.1%	34.4%	10.0%											
Cr	2C1	1A2f	2C7c	5C1bi	1A1a	1B1b								82.4%
	42.7%	20.1%	8.8%	4.1%	3.6%	3.2%								
Cu	1A3bvi	2C7c	2C1	1A3c										83.4%
	38.5%	24.8%	15.8%	4.4%										
Ni	1A2e	1A2c	1A1b	2C1	1A2gviii	1A2f								83.9%
	23.9%	20.8%	16.6%	10.4%	6.6%	5.5%								
Se	1A1a	2C7c	1A2f											85.5%
	37.3%	27.8%	20.4%											
Zn	2C7c	2C1	1A2f	1A4bi	1A3bi									84.3%
	35.1%	31.3%	8.5%	4.7%	4.7%									
PCDD/ PCDF	2C1	1A1a	5C1a	5C1bii	2C7c									81.0%
	32.4%	23.7%	11.4%	7.2%	6.3%									
PAH	2C1	1B1b	2D3a											81.4%
	32.6%	31.7%	17.1%											
HCB	1A1a	5C1bii	1A2gviii											83.5%
	55.3%	16.1%	12.1%											
PCB	2C1													93.5%
	93.5%													

2000	Key source categories (sorted from high to low, from left to right)																Total
NO <sub>x</sub>	1A3biii 22.9%	1A3bi 16.5%	1A1a 12.2%	2A1 5.1%	1A4bi 4.4%	1A2a 3.8%	1A3bii 3.1%	1A2f 2.8%	3Da2a 2.4%	1A4cii 2.1%	1A1b 2.1%	2B10a 2.1%	1A3dii 1.9%				81.3%
NM VOC	2D3d 14.8%	3B1b 9.4%	1A3bi 8.9%	2B10a 6.5%	2D3a 5.6%	1B2aiv 5.1%	3B1a 4.6%	2D3h 4.5%	1A4bi 4.4%	2D3g 4.1%	1A3bv 2.8%	1B2av 2.7%	2D3e 2.2%	3B3 2.0%	1A3biii 1.7%	2D3i 1.7%	81.0%
SO <sub>x</sub>	1A1a 20.1%	1A1b 13.4%	1A4bi 11.5%	1A2f 6.3%	2B10a 5.6%	2C1 4.6%	1A2a 4.4%	2A6 4.3%	2A2 3.4%	1A4ci 3.4%	1A2e 3.2%						80.2%
NH <sub>3</sub>	3B3 33.2%	3B1b 13.1%	3Da2a 10.7%	3Da3 9.8%	3B1a 8.9%	3Da1 8.4%											84.0%
PM <sub>2.5</sub>	1A4bi 28.1%	2C1 18.4%	1A3bi 10.7%	1A3biii 5.9%	1A1a 3.0%	1A3bii 2.9%	1A3bvi 2.4%	2A6 2.3%	1B1b 2.0%	2A5a 1.8%	1A2f 1.5%	1A1b 1.4%					80.4%
PM <sub>10</sub>	1A4bi 21.2%	2C1 17.2%	1A3bi 7.9%	1A1a 4.5%	1A3biii 4.3%	2A5a 4.0%	1A3bvi 3.3%	3B4gi 3.0%	3B4gii 2.5%	1B1b 2.4%	2A1 2.3%	1A3bii 2.2%	3Da1 2.1%	3B3 2.0%	1A3bvii 1.8%		80.6%
TSP	2C1 19.9%	1A4bi 16.0%	2A5a 7.6%	1A3bi 5.6%	1A1a 5.1%	1B1b 3.8%	1A3bii 3.1%	1A3bvi 3.1%	3B3 3.1%	2A2 2.8%	1A3bvi 2.6%	3B4gi 2.2%	2A1 2.0%	3B4gii 1.8%	1A2gvii 1.6%		80.2%
BC	1A3bi 32.7%	1A3bii 16.1%	1A4bi 15.7%	1A3bii 9.7%	1B1b 5.8%	2C1 5.1%											85.0%
CO	2C1 28.5%	1A2a 22.8%	1A3bi 20.2%	1A4bi 9.6%													81.0%
Pb	2C1 71.6%	2C7c 10.2%															81.8%
Cd	2C1 42.0%	2C7c 14.0%	1B2aiv 7.9%	1B1b 7.0%	1A1a 6.3%	1A4bi 5.7%											82.9%
Hg	2B10a 22.6%	2C1 21.1%	1A1a 18.5%	2A1 10.2%	1B2aiv 7.3%	2C7c 4.7%											84.4%
As	2C1 42.1%	2C7c 28.1%	1A1a 7.7%	1A2c 2.7%													80.6%
Cr	2C1 63.8%	1A2f 8.6%	1A3bvi 4.3%	1A1a 3.9%													80.6%
Cu	1A3bvi 43.9%	2C1 14.2%	1A3c 13.6%	2C7c 8.8%													80.4%
Ni	1A1b 29.4%	2C1 19.3%	1B2aiv 10.6%	1A2e 7.4%	1A2c 6.5%	1A3bvi 6.2%	1A2f 3.7%										82.9%
Se	1A2f 49.1%	2C7c 13.4%	1A1a 12.4%	2C1 5.8%													80.7%
Zn	2C1 43.9%	2C7c 20.3%	1A2f 8.0%	1A3bi 4.9%	1A3bvi 4.2%												81.3%
PCDD/ PCDF	2C1 47.4%	5C2 19.8%	1A4bi 12.1%	1A1a 9.0%													88.3%
PAH	1B1b 36.4%	2D3a 26.2%	1A4bi 17.3%	2C1 14.7%													94.6%
HCB	1A2gvii 39.6%	2L 30.7%	2A1 12.8%														83.1%
PCB	2C1 93.9%																93.9%

2005	Key source categories (sorted from high to low, from left to right)																	Total
NO <sub>x</sub>	1A3biii 25.1%	1A3bi 15.4%	1A1a 11.1%	1A4bi 4.5%	2A1 4.5%	1A3bii 3.9%	1A2f 3.2%	1A2a 2.6%	1A3dii 2.3%	3Da2a 2.2%	1A1b 2.1%	2B10a 2.0%	1A4cii 2.0%					80.9%
NM VOC	2D3d 15.6%	3B1b 10.4%	2D3a 7.4%	2B10a 7.2%	1A4bi 6.3%	1A3bi 5.0%	3B1a 4.8%	2D3h 4.0%	1B2aiv 3.9%	2D3g 3.9%	1A3bv 2.1%	3B3 2.1%	1A3bii 1.9%	1B2av 1.8%	1B2b 1.8%	1A4cii 1.6%	2H2 1.5%	81.2%
SO <sub>x</sub>	1A1a 20.4%	1A1b 13.3%	1A4bi 12.7%	1A2f 7.9%	2B10a 5.6%	2C1 4.5%	2A6 4.2%	1A4ci 4.0%	1A2a 3.7%	2A1 3.7%								80.1%
NH <sub>3</sub>	3B3 31.1%	3B1b 12.7%	3Da2a 11.6%	3Da3 10.6%	3Da1 9.0%	3B1a 8.6%												83.6%
PM <sub>2.5</sub>	1A4bi 38.6%	2C1 13.0%	1A3bi 8.0%	1A3biii 5.4%	1A3bii 3.1%	1A3bvi 2.9%	1A2f 2.1%	2A5a 2.1%	1A2gvii 1.8%	1A3bvii 1.6%	1A1a 1.6%							80.2%
PM <sub>10</sub>	1A4bi 29.7%	2C1 13.3%	1A3bi 6.0%	2A5a 4.7%	1A3biii 4.1%	1A3bvi 4.1%	3B4gi 3.0%	3Da1 2.4%	1A1a 2.4%	3B4gii 2.4%	1A3bii 2.3%	1A3bvi 2.2%	1B1b 2.1%	3B3 2.0%				80.6%
TSP	1A4bi 22.0%	2C1 15.6%	2A5a 8.7%	1A3bi 4.2%	1A3bvi 3.8%	1B1b 3.6%	1A1a 3.6%	1A3bvii 3.2%	3B3 3.1%	1A3biii 2.9%	2A2 2.7%	3B4gi 2.1%	1A2gvii 1.7%	3Da1 1.7%	3B4gii 1.7%			80.5%
BC	1A3bi 28.2%	1A4bi 20.9%	1A3biii 16.1%	1A3bii 11.3%	2C1 4.7%													81.2%
CO	2C1 32.2%	1A2a 20.9%	1A4bi 12.7%	1A3bi 12.4%	2A2 4.8%													83.0%
Pb	2C1 69.5%	2C7c 13.1%																82.6%
Cd	2C1 33.2%	2C7c 15.8%	1B2aiv 9.2%	1A4bi 8.8%	1B1b 7.0%	1A1a 6.0%	1A2f 5.9%											85.8%
Hg	2C1 33.7%	2B10a 12.4%	2A1 10.8%	1B2aiv 10.6%	1A1a 8.7%	2C7c 5.1%												81.2%
As	2C1 31.6%	2C7c 29.8%	1A1a 12.8%	1A2f 4.5%	1A2c 3.5%													82.2%
Cr	2C1 63.3%	1A1a 6.7%	1A3bvi 5.0%	1B1b 3.4%	1A2a 3.3%													81.6%
Cu	1A3bvi 49.7%	1A3c 14.1%	2C1 10.0%	2G 7.5%														81.4%
Ni	1A1b 24.2%	2C1 13.5%	1A1a 13.5%	1A2c 9.1%	1A3bvi 8.2%	1A2e 7.0%	1B2aiv 6.4%											81.9%
Se	1A2f 92.9%																	92.9%
Zn	2C1 45.4%	2C7c 16.9%	1A4bi 7.0%	1A3bi 6.6%	1A3bvi 6.4%													82.3%
PCDD/ PCDF	5C2 28.9%	1A4bi 26.5%	2C1 25.6%															81.0%
PAH	1B1b 37.3%	2D3a 31.0%	1A4bi 19.6%															87.9%
HCB	2A1 29.6%	2L 29.4%	2C1 22.5%															81.5%
PCB	2C1 94.6%																	94.6%

2010	Key source categories (sorted from high to low, from left to right)																Total
NO <sub>x</sub>	1A3biii 24.0%	1A3bi 20.8%	1A1a 6.2%	1A4bi 5.3%	1A3bii 4.9%	2A1 4.2%	3Da2a 2.5%	1A2f 2.4%	1A4cii 2.4%	2B10a 2.3%	2C1 2.1%	3Da1 2.0%	1A3dii 1.9%				80.9%
NM VOC	3B1b 12.4%	1A4bi 11.3%	2D3d 10.5%	2D3a 9.4%	2B10a 7.5%	3B1a 5.4%	2D3g 3.6%	1A3bi 3.1%	3B3 2.7%	1B2aiv 2.6%	1B2b 2.3%	2D3h 2.2%	2H2 1.9%	1A3bv 1.8%	2D3e 1.8%	1A4cii 1.8%	80.1%
SO <sub>x</sub>	1A4bi 14.5%	1A1b 14.3%	1A2f 10.5%	2C1 8.3%	2A1 7.8%	2B10a 7.0%	2A6 4.9%	1A1a 4.5%	1B2c 4.5%	1A2a 3.1%	1A4ci 2.7%						81.8%
NH <sub>3</sub>	3B3 31.8%	3B1b 12.1%	3Da2a 11.4%	3Da1 11.2%	3Da3 8.8%	3B1a 7.7%											83.1%
PM <sub>2,5</sub>	1A4bi 57.8%	1A3bi 6.3%	2C1 3.7%	1A3biii 3.3%	1A3bvi 3.2%	1A3bii 2.7%	2A5a 2.0%	1A3bvii 1.8%									80.7%
PM <sub>10</sub>	1A4bi 47.0%	1A3bi 5.0%	1A3bvi 4.7%	2A5a 4.6%	2C1 3.5%	3B4gi 2.9%	3B4gii 2.7%	1A3biii 2.6%	1A3bvi 2.6%	3Da1 2.5%	1A3bii 2.2%						80.2%
TSP	1A4bi 38.8%	2A5a 9.0%	1A3bvi 4.8%	1A3bvii 4.0%	1A3bi 3.9%	2C1 3.9%	3B3 3.7%	1B1b 2.5%	3B4gi 2.2%	3B4gii 2.1%	1A3bii 2.0%	1A2gvii 2.0%	3Da1 1.9%				80.9%
BC	1A4bi 32.5%	1A3bi 26.5%	1A3bii 11.8%	1A3biii 11.4%													82.2%
CO	2C1 31.0%	1A4bi 25.2%	1A3bi 9.9%	2A2 7.0%	1A2a 5.9%	1A3biii 3.1%											82.1%
Pb	2C1 56.4%	2C7c 6.6%	1A3bvi 6.1%	2A2 5.8%	2L 4.7%	1A2a 3.1%											82.7%
Cd	2C7c 23.8%	2C1 21.7%	1A4bi 13.2%	1B2aiv 8.5%	1A1a 7.5%	1B1b 4.4%	1A2f 4.2%										83.3%
Hg	2C1 22.6%	1A1a 14.6%	1B2aiv 12.1%	2A1 11.9%	2B10a 10.4%	1A2c 3.5%	1A4bi 3.4%	2A2 3.1%									81.7%
As	2C7c 26.7%	2C1 25.2%	1A1a 13.1%	1A2a 4.5%	1A2c 4.5%	1A2f 3.8%	1A2gviii 3.8%										81.7%
Cr	2C1 50.2%	1A2a 13.3%	1A3bvi 6.8%	2C7c 5.1%	1A4bi 5.1%												80.5%
Cu	1A3bvi 56.3%	1A3c 14.1%	2G 8.2%	2C1 6.5%													85.0%
Ni	2C1 21.4%	1A3bvi 21.3%	1B2aiv 12.6%	1A2c 6.5%	1A2f 6.1%	1A2a 5.8%	1A2e 4.2%	2B10a 4.0%									81.9%
Se	1A2f 80.8%																80.8%
Zn	2C1 35.4%	1A4bi 13.0%	2C7c 11.5%	1A3bi 8.4%	1A3bvi 8.3%	1A3biii 4.0%											80.7%
PCDD/ PCDF	1A4bi 45.2%	5C2 22.4%	2C1 12.0%	1A3bi 7.9%													87.4%
PAH	2D3a 36.6%	1A4bi 32.2%	1B1b 22.6%														91.5%
HCB	2L 45.5%	2A1 26.8%	1A1a 14.8%														87.1%
PCB	2A1 74.5%	2C1 24.5%															99.0%

2011	Key source categories (sorted from high to low, from left to right)																Total
NOx	1A3biii 23.1%	1A3bi 22.0%	1A1a 5.3%	1A3bii 5.1%	2A1 5.0%	1A4bi 4.5%	3Da2a 2.5%	1A2f 2.5%	1A4cii 2.4%	2B10a 2.4%	2C1 2.2%	3Da1 2.1%	1A3dii 2.0%				81.1%
NM VOC	3B1b 13.1%	2D3d 10.8%	2D3a 10.3%	1A4bi 8.9%	2B10a 7.2%	3B1a 5.7%	2D3g 4.1%	3B3 2.9%	1A3bi 2.8%	1B2aiv 2.3%	1B2b 2.3%	1A3bv 2.2%	2H2 2.2%	2D3h 1.9%	1A4cii 1.9%	3B4gii 1.7%	80.3%
SOx	1A1b 15.6%	1A4bi 14.1%	1A2f 11.3%	2C1 10.1%	2A1 8.8%	2B10a 8.4%	2A6 4.9%	1A1a 3.9%	1B2c 3.9%								81.2%
NH3	3B3 31.7%	3B1b 12.0%	3Da1 11.2%	3Da2a 11.1%	3Da3 8.7%	3B1a 7.8%											82.5%
PM2.5	1A4bi 51.2%	1A3bi 6.2%	1A3bvi 3.9%	2C1 3.7%	1A3biii 3.5%	1A3bii 2.9%	2A5a 2.5%	1A2d 2.4%	1A3bvii 2.1%	1A2gviii 1.8%							80.2%
PM10	1A4bi 40.1%	2A5a 5.7%	1A3bvi 5.5%	1A3bi 4.8%	2C1 3.4%	3B4gi 3.3%	3B4gii 3.3%	1A3bvii 3.0%	3Da1 2.9%	1A3biii 2.6%	3B3 2.5%	1A3bii 2.2%	1A2d 1.9%				81.1%
TSP	1A4bi 32.1%	2A5a 11.3%	1A3bvi 5.5%	1A3bvii 4.6%	3B3 4.2%	1A3bi 3.6%	2C1 3.1%	1B1b 2.5%	3B4gi 2.5%	3B4gii 2.5%	1A2gvii 2.3%	3Da1 2.2%	1A3biii 2.0%	1A3bii 1.7%			80.1%
BC	1A4bi 28.9%	1A3bi 25.7%	1A3bii 12.4%	1A3biii 12.0%	2C1 2.6%												81.5%
CO	2C1 32.9%	1A4bi 23.3%	1A3bi 10.7%	2A2 6.3%	1A3biii 3.7%	1A3biv 3.1%	1A4bii 3.0%										83.0%
Pb	2C1 53.5%	2C7c 8.9%	1A3bvi 8.1%	1A2a 4.0%	1B2aiv 3.6%	1A1a 3.6%											81.7%
Cd	2C7c 28.5%	2C1 18.6%	1A4bi 10.1%	1B2aiv 9.0%	1A1a 8.6%	1B1b 4.7%	1A2d 2.8%										82.4%
Hg	1A1a 20.4%	2C1 16.3%	2A1 13.3%	1B2aiv 12.2%	2B10a 11.3%	1A2c 3.5%	2C7c 3.1%										80.2%
As	2C7c 30.7%	2C1 24.3%	1A1a 13.5%	1A2c 5.4%	1A2f 3.4%	1B1b 3.3%											80.5%
Cr	2C1 41.6%	1A2a 18.0%	1A3bvi 7.8%	1A1a 7.5%	2C7c 4.5%	1A4bi 4.2%											83.8%
Cu	1A3bvi 57.3%	1A3c 14.2%	2G 8.4%	2C1 6.1%													86.0%
Ni	1A3bvi 22.3%	2C1 18.6%	1A2c 14.2%	1B2aiv 9.3%	1A2a 7.3%	2B10a 4.9%	1A2e 3.7%										80.2%
Se	1A2f 44.3%	1A1a 26.1%	2A1 9.0%	1A2c 7.5%													86.9%
Zn	2C1 31.9%	2C7c 15.5%	1A4bi 9.9%	1A3bi 9.0%	1A3bvi 8.8%	1A3biii 4.1%	1A2a 3.5%										82.7%
PCDD/ PCDF	1A4bi 39.2%	5C2 24.1%	2C1 16.9%														80.2%
PAH	2D3a 38.9%	1A4bi 24.8%	1B1b 22.5%														86.2%
HCB	2C1 45.3%	2L 26.2%	2A1 15.0%														86.5%
PCB	2A1 79.1%	2C1 19.7%															98.8%

2012	Key source categories (sorted from high to low, from left to right)																Total
NOx	1A3bi	1A3biii	1A1a	1A3bii	1A4bi	2A1	3Da2a	1A4cii	2B10a	1A2f	3Da1	2C1	1A3dii	1A4ai			81.2%
	22.1%	20.7%	5.2%	5.1%	5.1%	4.9%	2.7%	2.5%	2.5%	2.4%	2.2%	2.0%	2.0%	1.7%			
NMVOC	3B1b	1A4bi	2D3a	2D3d	2B10a	3B1a	2D3g	3B3	1A3bi	2H2	1B2b	1B2aiv	2D3h	1A4cii	1A3bv	3B4gii	80.5%
	13.0%	10.9%	10.5%	9.6%	7.4%	5.8%	3.8%	3.0%	2.6%	2.2%	2.2%	2.1%	1.9%	1.9%	1.9%	1.8%	
SOx	1A1b	1A4bi	2C1	2A1	2B10a	1A2f	1A1a	2A6	1B2c	1A4ai							81.8%
	16.1%	15.8%	12.2%	8.3%	7.9%	7.9%	4.3%	3.7%	3.0%	2.5%							
NH3	3B3	3B1b	3Da1	3Da2a	3Da3	3B1a											82.4%
	31.8%	11.8%	11.2%	11.2%	8.7%	7.7%											
PM2,5	1A4bi	1A3bi	1A3bvi	2C1	1A2d	1A3biii	2A5a	1A3bii	1A3bvii								81.0%
	57.8%	4.9%	3.4%	3.0%	2.6%	2.6%	2.5%	2.4%	1.9%								
PM10	1A4bi	2A5a	1A3bvi	1A3bi	3B4gi	3B4gii	2C1	3Da1	1A3bvii	3B3	1A2d	1A3biii					81.7%
	46.1%	5.7%	4.9%	3.8%	3.2%	3.2%	2.9%	2.7%	2.7%	2.4%	2.1%	2.0%					
TSP	1A4bi	2A5a	1A3bvi	1A3bvii	3B3	1A3bi	2C1	3B4gi	3B4gii	1A2gvii	1B1b	3Da1	1A2d	2C1			80.9%
	36.9%	12.8%	4.9%	4.0%	4.0%	2.9%	2.5%	2.5%	2.4%	2.2%	2.2%	2.0%	1.7%				
BC	1A4bi	1A3bi	1A3bii	1A3biii	1A2d												81.6%
	36.1%	22.3%	11.1%	10.0%	2.2%												
CO	1A4bi	2C1	1A3bi	1A3biii	1A4bii	1A3biv											82.9%
	31.2%	30.6%	11.0%	3.6%	3.4%	3.1%											
Pb	2C1	1A3bvi	2C7c	1A2a	1B2aiv												80.4%
	57.9%	7.7%	7.4%	3.8%	3.6%												
Cd	2C7c	1A4bi	2C1	1B2aiv	1A1a	1B1b	1A2d										82.5%
	31.5%	13.8%	10.6%	10.2%	8.2%	4.6%	3.6%										
Hg	2A1	1B2aiv	2C1	2B10a	1A1a	1A2c	1A2f	1A4bi	2C7c								83.9%
	16.7%	15.6%	13.9%	12.7%	8.4%	4.6%	4.2%	3.9%	3.9%								
As	2C7c	2C1	1A1a	1A2c	1A2f	1B1b											81.2%
	38.1%	16.3%	14.4%	5.1%	3.9%	3.4%											
Cr	2C1	1A2a	1A3bvi	1A1a	1A4bi												82.7%
	44.5%	18.3%	7.8%	6.7%	5.4%												
Cu	1A3bvi	1A3c	2G														82.3%
	56.0%	17.5%	8.8%														
Ni	1A3bvi	1A2c	2C1	1A2a	1A2f	1B2aiv	1A1a	1A2d	2C7c								82.5%
	27.5%	14.8%	10.7%	8.9%	5.4%	4.6%	4.2%	3.3%	3.1%								
Se	1A2f	1A1a	2A1														81.1%
	43.3%	28.3%	9.4%														
Zn	2C7c	1A4bi	1A3bi	1A3bvi	2C1	1A3biii	1A2a										81.0%
	31.7%	14.1%	9.8%	9.6%	7.4%	4.4%	3.9%										
PCDD/ PCDF	1A4bi	2C1	5C2														82.6%
	41.3%	22.3%	19.0%														
PAH	2D3a	1A4bi	1B1b														93.2%
	41.7%	31.6%	19.9%														
HCB	2L	2C1	2A1														80.1%
	33.2%	26.4%	20.5%														
PCB	2C1	2A1															91.6%
	50.4%	41.2%															

2013	Key source categories (sorted from high to low, from left to right)																	Total
NOx	1A3bi	1A3biii	1A4bi	1A3bii	1A1a	2A1	3Da2a	2B10a	1A4cii	3Da1	2C1	1A4ai	1A3dii	1A2f				80.7%
	22.5%	19.8%	5.8%	5.2%	5.0%	4.3%	2.8%	2.7%	2.4%	2.3%	2.1%	2.0%	2.0%	1.9%				
NMVOC	3B1b	1A4bi	2D3a	2D3d	2B10a	3B1a	2D3g	3B3	1A3bi	1B2aiv	1B2b	2H2	1A4cii	3B4gii	2D3h	1A3bv		81.0%
	13.0%	12.3%	10.8%	9.0%	7.0%	6.0%	3.3%	3.0%	2.5%	2.3%	2.2%	2.2%	1.9%	1.8%	1.8%	1.7%		
SOx	1A4bi	1A1b	2C1	2B10a	2A1	1A2f	1A1a	1B2c	2A6	1A4ai								82.5%
	18.6%	15.0%	13.2%	8.1%	7.0%	5.7%	4.3%	3.9%	3.9%	2.8%								
NH3	3B3	3B1b	3Da1	3Da2a	3Da3	3B1a												82.7%
	32.1%	11.9%	11.6%	10.7%	8.6%	7.8%												
PM2.5	1A4bi	1A3bi	1A3bvi	1A2d	2A5a	1A3biii	1A3bii	1A2gviii										80.5%
	61.9%	4.2%	3.2%	2.5%	2.4%	2.2%	2.1%	1.9%										
PM10	1A4bi	2A5a	1A3bvi	1A3bi	3B4gii	3B4gi	3Da1	1A3bvii	3B3	1A2d	2C1							80.5%
	49.4%	5.5%	4.7%	3.3%	3.2%	3.1%	2.6%	2.5%	2.3%	2.0%	1.8%							
TSP	1A4bi	2A5a	1A3bvi	1A3bvii	3B3	1A3bi	3B4gii	3B4gi	1A2gvii	3Da1	2C1	1A2d	2B10a					80.6%
	39.8%	12.3%	4.7%	3.9%	3.8%	2.5%	2.5%	2.4%	2.0%	2.0%	1.7%	1.6%	1.5%					
BC	1A4bi	1A3bi	1A3bii	1A3biii														80.5%
	41.6%	20.0%	10.0%	9.0%														
CO	2A2	1A4bi	2C1	1A3bi														81.9%
	30.8%	23.5%	20.4%	7.2%														
Pb	2C1	1A3bvi	2C7c	1A4bi	1B2aiv	1A2a												83.9%
	54.5%	8.6%	8.2%	4.5%	4.1%	3.9%												
Cd	2C7c	1A4bi	1B2aiv	2C1	1A1a	1B1b	1A2d											81.7%
	34.9%	14.5%	9.6%	9.6%	6.4%	3.4%	3.4%											
Hg	2B10a	2A1	1B2aiv	1A1a	2C1	1A4bi	1A2c	2C7c										81.6%
	18.4%	14.4%	14.0%	10.4%	9.7%	7.5%	4.0%	3.3%										
As	2C7c	1A1a	2C1	2A1	1A2c	1A2f	1B2aiv											81.5%
	38.2%	12.8%	10.3%	6.9%	5.7%	4.4%	3.3%											
Cr	2C1	1A2a	1A3bvi	1A4bi	1A1a	1B1b	1A2f											80.2%
	22.5%	17.3%	14.1%	11.0%	8.7%	3.4%	3.3%											
Cu	1A3bvi	1A3c	2G															83.6%
	58.3%	16.0%	9.3%															
Ni	1A3bvi	2C1	1A2c	1B2aiv	1A2f	1A1a	1A1b	2B10a	2A1	1A2b	2C7c							82.1%
	33.6%	13.8%	6.6%	5.5%	5.0%	4.6%	3.2%	2.6%	2.5%	2.4%	2.3%							
Se	1A2f	1A1a	2A1															82.3%
	46.2%	22.0%	14.1%															
Zn	2C7c	1A4bi	1A3bi	1A3bvi	2C1	1A3biii	1A2d	1A2a										83.2%
	22.4%	17.2%	10.7%	10.5%	9.3%	4.8%	4.2%	4.2%										
PCDD/ PCDF	1A4bi	5C2	2C1															81.5%
	53.1%	19.4%	9.0%															
PAH	2D3a	1A4bi	1B1b															94.1%
	43.4%	36.3%	14.5%															
HCB	2L	1A1a	2A1															83.8%
	38.9%	27.2%	17.7%															
PCB	2A1	2C1																87.3%
	56.8%	30.6%																

## 1.B Trend assessment

The following tables show the trend assessment for 2013 for all reported pollutants.

NOx		1990	2013	trend	Ass. %	contrib	cum.total
1A1a	Public electricity and heat production	60.73	10.34	0.049	25.20%	25.20%	
1A3bi	Road transport: Passenger cars	111.27	46.28	0.023	11.90%	37.10%	
1A3bii	Road transport: Light duty vehicles	8.86	10.69	0.015	7.90%	45.00%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	13.56	1.5	0.013	6.70%	51.60%	
1A3biii	Road transport: Heavy duty vehicles and buses	73.2	40.62	0.01	5.00%	56.60%	
1A4bi	Residential: Stationary	16.53	11.87	0.009	4.50%	61.10%	
2B10a	Chemical industry: Other (please specify in the IIR)	5.12	5.52	0.007	3.70%	64.80%	
1A4ai	Commercial/institutional: Stationary	3.49	4.03	0.006	2.90%	67.70%	
1A3di(ii)	International inland waterways	2.67	3.6	0.006	2.80%	70.50%	
2A2	Lime production	2.71	3.55	0.005	2.80%	73.30%	
1A4ci	Agriculture/Forestry/Fishing: Stationary	3.2	3.37	0.004	2.20%	75.50%	
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	6.79	5.03	0.004	2.00%	77.60%	
1A3dii	National navigation (shipping)	4.84	4.02	0.004	2.00%	79.60%	
2C1	Iron and steel production	5.38	4.26	0.004	2.00%	81.50%	

NMVOC		1990	2013	trend	Ass. %	contrib	cum.total
1A3bi	Road transport: Passenger cars	65.03	3.45	0.067	21.70%	21.70%	
1A4bi	Residential: Stationary	13.01	16.97	0.034	11.20%	32.80%	
2D3a	Domestic solvent use including fungicides	9.41	14.89	0.033	10.60%	43.40%	
3B1b	Manure management - Non-dairy cattle	21.45	17.90	0.027	8.80%	52.20%	
2D3d	Coating applications	49.72	12.31	0.023	7.30%	59.60%	
1A3bv	Road transport: Gasoline evaporation	17.88	2.36	0.014	4.60%	64.20%	
1B2aiv	Fugitive emissions oil: Refining / storage	16.93	3.11	0.011	3.50%	67.70%	
2D3h	Printing	13.96	2.44	0.009	3.00%	70.80%	
1B2av	Distribution of oil products	11.37	1.80	0.008	2.60%	73.40%	
3B3	Manure management - Swine	4.54	4.13	0.007	2.20%	75.60%	
3B1a	Manure management - Dairy cattle	15.04	8.30	0.007	2.10%	77.70%	
2B10a	Chemical industry: Other (please specify in the IIR)	29.56	9.69	0.006	2.10%	79.90%	
2H2	Food and beverages industry	2.10	2.97	0.006	2.00%	81.90%	

SOx		1990	2013	trend	Ass. %	contrib	cum.total
1A1a	Public electricity and heat production	95.2	1.97	0.027	24.90%	24.90%	
2C1	Iron and steel production	10.51	6.02	0.013	11.90%	36.80%	
1A4bi	Residential: Stationary	31.24	8.47	0.013	11.50%	48.30%	
1A4ci	Agriculture/Forestry/Fishing: Stationary	28.41	0.71	0.008	7.10%	55.40%	
2A1	Cement production	4.33	3.2	0.007	6.70%	62.10%	
1B2c	Venting and flaring (oil, gas, combined oil and gas)	0.25	1.79	0.005	4.40%	66.60%	
1A1b	Petroleum refining	40.85	6.83	0.005	4.40%	70.90%	
2A6	Other mineral products (please specify in the IIR)	0.46	1.76	0.005	4.30%	75.20%	
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	16.96	0.83	0.004	3.20%	78.40%	
1A2e	Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	16.03	1.03	0.003	2.40%	80.90%	

NH3		1990	2013	trend	Ass. %	contrib	cum.total
3B3	Manure management - Swine	49.85	19.96	0.055	29.70%	29.70%	
3Da1	Inorganic N-fertilizers (includes also urea application)	7.27	7.23	0.029	15.50%	45.20%	
3Da2a	Animal manure applied to soils	9.04	6.63	0.016	8.40%	53.70%	
3B1a	Manure management - Dairy cattle	11.85	4.83	0.012	6.70%	60.30%	
1A4bi	Residential: Stationary	0.71	1.82	0.012	6.60%	67.00%	
3B4gii	Manure management - Broilers	1.39	1.92	0.01	5.40%	72.40%	
3B4gi	Manure management - Laying hens	4.52	1.26	0.01	5.20%	77.60%	
5D1	Domestic wastewater handling	3.36	0.96	0.007	3.80%	81.40%	

PM2.5		2000	2013	trend	Ass. %	contrib	cum.total
1A4bi	Residential: Stationary	11.63	20.72	0.274	42.90%	42.90%	
2C1	Iron and steel production	7.59	0.62	0.134	20.90%	63.80%	
1A3bi	Road transport: Passenger cars	4.41	1.4	0.053	8.20%	72.10%	
1A3biii	Road transport: Heavy duty vehicles and buses	2.43	0.73	0.03	4.70%	76.80%	
1A1a	Public electricity and heat production	1.25	0.27	0.018	2.80%	79.60%	
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.18	0.85	0.017	2.70%	82.30%	

PM10		2000	2013	trend	Ass. %	contrib	cum.total
1A4bi	Residential: Stationary	11.9	21.26	0.216	37.80%	37.80%	
2C1	Iron and steel production	9.67	0.79	0.118	20.60%	58.40%	
1A3bi	Road transport: Passenger cars	4.41	1.4	0.035	6.20%	64.60%	
1A1a	Public electricity and heat production	2.52	0.3	0.029	5.10%	69.70%	
1A3biii	Road transport: Heavy duty vehicles and buses	2.43	0.73	0.02	3.50%	73.20%	
2A1	Cement production	1.32	0.03	0.017	3.10%	76.20%	
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	1.33	0.26	0.014	2.40%	78.60%	
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.2	0.87	0.013	2.20%	80.90%	

TSP		2000	2013	trend	Ass. %	contrib	cum.total
1A4bi	Residential: Stationary	12.62	22.42	0.17	30.90%	30.90%	
2C1	Iron and steel production	15.69	0.95	0.131	23.70%	54.60%	
2A5a	Quarrying and mining of minerals other than coal	5.96	6.95	0.034	6.20%	60.80%	
1A1a	Public electricity and heat production	4.02	0.34	0.032	5.90%	66.70%	
1A3bi	Road transport: Passenger cars	4.41	1.4	0.022	4.10%	70.80%	
2A2	Lime production	2.17	0.04	0.019	3.50%	74.20%	
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	2.99	0.79	0.017	3.10%	77.40%	
1A3biii	Road transport: Heavy duty vehicles and buses	2.43	0.73	0.013	2.30%	79.70%	
2A1	Cement production	1.55	0.13	0.012	2.30%	82.00%	

BC		2000	2013	trend	Ass. %	contrib	cum.total
1A4bi	Residential: Stationary	1.4	2.31	0.162	43.20%	43.20%	
1A3bi	Road transport: Passenger cars	2.9	1.11	0.08	21.30%	64.50%	
1A3biii	Road transport: Heavy duty vehicles and buses	1.43	0.5	0.044	11.80%	76.30%	
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	0.51	0.07	0.028	7.50%	83.80%	

CO		1990	2013	trend	Ass. %	contrib	cum.total
1A3bi	Road transport: Passenger cars	578.73	39.62	0.132	31.60%	31.60%	
2A2	Lime production	19.68	170.36	0.115	27.60%	59.20%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	261.36	1.39	0.071	17.10%	76.30%	
1A4bi	Residential: Stationary	137.59	130.23	0.054	13.00%	89.30%	

Pb		1990	2013	trend	Ass. %	contrib	cum.total
1A3bi	Road transport: Passenger cars	95.82	0.22	0.038	34.70%	34.70%	
2C1	Iron and steel production	50.63	14.38	0.036	32.50%	67.30%	
1A3bvi	Road transport: Automobile tyre and brake wear	1.75	2.28	0.008	7.50%	74.80%	
1B2aiv	Fugitive emissions oil: Refining / storage	0	1.09	0.004	3.90%	78.60%	
1A4bi	Residential: Stationary	2.72	1.19	0.004	3.20%	81.90%	

Cd		1990	2013	trend	Ass. %	contrib	cum.total
1A4bi	Residential: Stationary	0.15	0.34	0.045	31.60%	31.60%	
2C1	Iron and steel production	0.92	0.22	0.02	14.20%	45.90%	
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.01	0.08	0.012	8.50%	54.40%	
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	0.38	0.08	0.01	7.00%	61.40%	
1A2gviii	Stationary combustion in manufacturing industries and construction: Other (please specify in the IIR)	0.02	0.07	0.01	6.90%	68.30%	
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.04	0.07	0.008	5.90%	74.20%	
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.03	0.06	0.007	4.90%	79.10%	
2B10a	Chemical industry: Other (please specify in the IIR)	0.13	0	0.007	4.80%	83.90%	

Hg		1990	2013	trend	Ass. %	contrib	cum.total
1A1a	Public electricity and heat production	1.32	0.17	0.038	24.50%	24.50%	
2A1	Cement production	0.33	0.24	0.025	16.30%	40.90%	
1A4bi	Residential: Stationary	0.13	0.12	0.015	9.80%	50.60%	
2C1	Iron and steel production	0.28	0.16	0.014	9.00%	59.60%	
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.03	0.07	0.01	6.70%	66.30%	
2C7c	Other metal production (please specify in the IIR)	0.03	0.05	0.008	5.30%	71.60%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.01	0.04	0.006	3.90%	75.50%	
2A2	Lime production	0.04	0.05	0.006	3.90%	79.40%	
5C1bv	Cremation	0.02	0.04	0.006	3.90%	83.30%	

As		1990	2013	trend	Ass. %	contrib	cum.total
2C1	Iron and steel production	2.04	0.17	0.059	26.50%	26.50%	
2C7c	Other metal production (please specify in the IIR)	1.02	0.62	0.057	25.50%	52.00%	
1A1a	Public electricity and heat production	1.94	0.21	0.048	21.60%	73.60%	
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.05	0.09	0.013	5.80%	79.40%	
2A1	Cement production	0.12	0.11	0.013	5.80%	85.30%	

Cr		1990	2013	trend	Ass. %	contrib	cum.total
2C1	Iron and steel production	14.27	1.34	0.038	28.50%	28.50%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.25	1.02	0.03	22.50%	51.00%	
1A3bvi	Road transport: Automobile tyre and brake wear	0.64	0.84	0.022	16.50%	67.50%	
1A4bi	Residential: Stationary	0.46	0.65	0.017	13.00%	80.60%	

Cu		1990	2013	trend	Ass. %	contrib	cum.total
1A3bvi	Road transport: Automobile tyre and brake wear	14.05	18.22	0.181	30.90%	30.90%	
2C1	Iron and steel production	6.9	0.83	0.113	19.30%	50.10%	
1A3c	Railways	1.68	5	0.092	15.70%	65.80%	
1A1a	Public electricity and heat production	3.72	0.52	0.059	10.10%	75.90%	
1A2b	Stationary combustion in manufacturing industries and construction: Non-ferrous metals	2.47	0.02	0.047	8.10%	83.90%	

Ni		1990	2013	trend	Ass. %	contrib	cum.total
1A3bvi	Road transport: Automobile tyre and brake wear	1.75	2.28	0.029	28.20%	28.20%	
1A2e	Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	14.66	0.15	0.016	15.90%	44.10%	
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	16.39	0.45	0.014	14.00%	58.10%	
5C1bi	Industrial waste incineration	9.16	0.01	0.011	11.10%	69.20%	
1A2gviii	Stationary combustion in manufacturing industries and construction: Other (please specify in the IIR)	8.04	0.1	0.009	8.40%	77.60%	
1A1a	Public electricity and heat production	7.75	0.31	0.005	5.30%	82.80%	

Se		1990	2013	trend	Ass. %	contrib	cum.total
1A1a	Public electricity and heat production	2.68	0.77	0.226	38.80%	38.80%	
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	1.27	1.63	0.147	25.20%	64.00%	
2A1	Cement production	0.16	0.5	0.077	13.30%	77.30%	
1A2c	Stationary combustion in manufacturing industries and construction: Chemicals	0.02	0.27	0.051	8.80%	86.00%	

Zn		1990	2013	trend	Ass. %	contrib	cum.total
2B10a	Chemical industry: Other (please specify in the IIR)	33.88	0.45	0.048	16.30%	16.30%	
1A4bi	Residential: Stationary	9.1	13.88	0.046	15.60%	31.90%	
2C1	Iron and steel production	46.29	7.53	0.036	12.20%	44.10%	
1A3bvi	Road transport: Automobile tyre and brake wear	6.15	8.47	0.027	9.20%	53.30%	
1A3bi	Road transport: Passenger cars	8.08	8.67	0.025	8.50%	61.80%	
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	15.43	0.44	0.021	7.10%	68.90%	
2C7c	Other metal production (please specify in the IIR)	40.59	18.13	0.018	6.00%	74.90%	
1A2d	Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	0.61	3.38	0.014	4.60%	79.50%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	1.48	3.38	0.012	4.20%	83.60%	

PCDD/PCDF		1990	2013	trend	Ass. %	contrib	cum.total
1A4bi	Residential: Stationary	18.09	22.88	0.037	32.80%	32.80%	
1A1a	Public electricity and heat production	148.49	0.97	0.018	15.40%	48.20%	
2C1	Iron and steel production	148.76	3.87	0.013	11.00%	59.30%	
5C1a	Municipal waste incineration	92.57	0.01	0.012	10.50%	69.80%	
5C2	Open burning of waste	21.05	8.37	0.012	10.40%	80.20%	

PAHsTotal1-4		1990	2013	trend	Ass. %	contrib	cum.total
2D3a	Domestic solvent use including fungicides	11.03	12.11	0.104	27.30%	27.30%	
2C1	Iron and steel production	22.72	0.05	0.101	26.50%	53.70%	
1B1b	Fugitive emission from solid fuels: Solid fuel transformation	30.64	4.04	0.086	22.50%	76.20%	
1A4bi	Residential: Stationary	12.15	10.13	0.074	19.40%	95.60%	

HCB		1990	2013	trend	Ass. %	contrib	cum.total
1A2gviii	Stationary combustion in manufacturing industries and construction: Other (please specify in the IIR)	29.46	0	0.043	39.80%	39.80%	
1A1a	Public electricity and heat production	13.75	3.84	0.019	17.40%	57.30%	
2L	Other production, consumption, storage, transportation or handling of bulk products (please specify in the IIR)	27.55	5.48	0.015	14.20%	71.40%	
2A1	Cement production	8.67	2.5	0.013	11.70%	83.20%	

PCBs		1990	2013	trend	Ass. %	contrib	cum.total
2C1	Iron and steel production	98.54	1.44	0.028	51.30%	51.30%	
2A1	Cement production	3.26	2.67	0.024	43.60%	94.90%	

## Annex 2: NMVOC emissions and TIER methods used in the Solvents sector.

Summary of the NMVOC emissions and the TIER methodologies used per region are in the tables below.

### FLANDERS

NMVOC emissions (tonnes/year) - codes NFR 2014		2005	2010	2013	Tier method for 2013	Comment
<b>2D</b>		<b>41436</b>	<b>27300</b>	<b>22612</b>		
<b>2D3a</b>	<b>Domestic solvent use including fungicides</b>	<b>7792</b>	<b>8278</b>	<b>8450</b>		
	Domestic solvent use (other than paint application)	7792	8278	8450	Tier 2	Activity data is the population. The emission factors for the different product groups have been determined for the year 2008. Same global emission factor has been assumed for the recent years.
<b>2D3d</b>	<b>Coating applications</b>	<b>19851</b>	<b>10249</b>	<b>7280</b>		
	Manufacture of automobile	4953	2433	1700	Tier 3	Use of facility data
	Paint application: car refinishing	1500	785	642	Tier 2	Sales of the different paints and their solvent content are used.
	Construction and buildings	4001	1950	1741	Tier 2	Sales of decorative paint (WB & SB) and their solvent content are used.
	Domestic use	2468	863	851	Tier 2	Sales of decorative paint (WB & SB) and their solvent content are used.
	Paint application: metal (excl. Manufacture of automobile)	3146	1442	881	Tier 3	Use of facility data
	Paint application: wood	1217	609	278	Tier 3	Use of facility data
	Paint application: textile (solvent coat)	825	629	109	Tier 3	Use of facility data
	Paint application: textile (carpet latexing)	53	38	33	Tier 2	Quantities of carpet with region specific information are used.
	Paint application: paper	42	59	1	Tier 3	Use of facility data
	Paint application: synthetics	778	570	174	Tier 3	Use of facility data
	Road markings	870	870	870	Tier 2	Quantities of paints with region specific information are used.
<b>2D3e</b>	<b>Degreasing</b>	<b>1520</b>	<b>1177</b>	<b>874</b>		
	Metal degreasing	1261	940	543	Tier 2	Consumption of chlorinated /non-chlorinated solvents with region specific factor for abatement technologies
	Other industrial cleaning: tank cleaning	259	237	331	Tier 2	Flemish BAT study
<b>2D3f</b>	<b>Dry cleaning</b>	<b>424</b>	<b>591</b>	<b>254</b>		
	Dry cleaning	424	591	254	Tier 2	Consumption of chlorinated /non-chlorinated solvents with region specific information
<b>2D3g</b>	<b>Chemical products</b>	<b>3947</b>	<b>2705</b>	<b>2688</b>		
	Polyester processing, Polyurethane processing, Polystyrene foam processing	2812	2072	2031	Tier 3	Use of facility data
	Rubber processing	101	57	23	Tier 3/Tier 1	Tier 3: Use of facility data; Tier 1: number of tires produced
	Pharmaceutical products manufacturing	388	202	208	Tier 3/Tier 2	Tier 3: Use of facility data; Tier 2: survey with region specific information

	Paints manufacturing, Inks manufacturing	445	268	348	Tier 3	Use of facility data
	Glues manufacturing	201	106	78	Tier 2	Flemish glues production figures with region specific information
2D3h	Printing	5331	1949	1434		
	Printing industry	5331	1949	1434	Tier 3/Tier 2	Tier 3: Use of facility data; Tier 2: Fetra/Febelgra survey with country specific information
2D3i	Other solvent use (please specify in the IIR)	2573	2352	1632		
	Fat, edible and non-edible oil extraction	541	745	566	Tier 3	Use of facility data
	Application of glues and adhesives	1839	1482	996	Tier 3	Use of facility data
	Preservation of wood	138	55	10	Tier 2	Activity data (Flemish BAT study) and their solvent content are used.
	Recovery of waste solvents	55	70	61	Tier 1	Quantities of recycled waste are used

## WALLONIA

NMVOC emissions (tonnes/year) - codes NFR 2013		2005	2010	2013	Tier method for 2013	Comment
2D & 2G	Other solvent and product use	19190	17146	13896		
2D3a	Domestic solvent use including fungicides	4680	4940	5031		
	060408 Domestic solvent use (other than paint application)	4680	4940	5031	Tier 2	Activity data is the population. The emission factors for the different product groups have been determined for the year 2008. Same global emission factor has been assumed for the recent years.
2D3b	Road paving with asphalt	NE	NE	NE		
2D3c	Asphalt roofing	NE	NE	NE		
2D3d	Coating applications	7430	5429	4262		
	060101 Manufacture of automobile	0	0	0	-	In Wallonia, there is no activity for this sector any more.
	060102 Car Repairing	775	406	332	Tier 2	Sales of the different paints and their solvent content are used.
	060103 Construction and buildings	2207	1076	895	Tier 2	Sales of decorative paint (WB & SB) and their solvent content are used.
	060104 Domestic use	1008	655	508	Tier 2	Sales of decorative paint (WB & SB) and their solvent content are used.
	060105 Coil Coating	30	10	8	Tier 3	Use of facility data
	060106 Boat building	7	7	5	Tier 3	Use of facility data
	060107 Wood	276	288	281	Tier 1	Sales of paint, proportion of WB and SB and solvent content are assumed.
	060108 Other industrial paint application	2520	2391	1653	Tier 1	Sales of paint and solvent content are assumed.
	060109 Other non industrial paint application	607	596	580	Tier 2	Emissions of road marking: Paint consumption (WB & SB) and solvent content are used.
2D3e	Degreasing	1107	1557	1015		
	060201 Metal degreasing	748	1152	660	Tier 1	Consumption of chlorinated solvent is known but no information is available for non-chlorinated solvent.
	060203 Electronic components manufacturing	IE	IE	IE		Included in 060204
	060204 Other industrial cleaning	359	404	355	Tier 1	Consumption of chlorinated solvent is known but no information is available for non-chlorinated

						solvent.
<b>2D3f</b>	<b>Dry cleaning</b>	<b>343</b>	<b>280</b>	<b>151</b>		
	060202 Dry cleaning	343	280	151	Tier 1	Consumption of chlorinated solvent is known but no information is available for non-chlorinated solvent.
<b>2D3g</b>	<b>Chemical products</b>	<b>3278</b>	<b>2830</b>	<b>1845</b>		
	060301 Polyester processing	283	464	464	Tier 1	National statistics are not available any more. Data cannot be actualized.
	060302 Polyvinylchloride processing	125	125	125	Tier 1	National statistics are not available any more. Data cannot be actualized.
	060303 Polyurethane processing	79	79	79	Tier 1	National statistics are not available any more. Data cannot be actualized.
	060304 Polystyrene foam processing	266	266	266	Tier 1	National statistics are not available any more. Data cannot be actualized.
	060305 Rubber processing	0	0	0	-	In Wallonia, there is no activity for this sector any more.
	060306 Pharmaceutical products manufacturing	460	341	172	Tier 3	Use of facility data
	060307 Paints manufacturing	529	438	369	Tier 1	Global solvent consumption and emission factor are assumed.
	060308 Inks manufacturing	22	10	14	Tier 3	Use of facility data
	060309 Glues manufacturing	19	0	0	-	In Wallonia, this activity is not significant. The NMVOC emissions of the few producers are reported under SNAP 040527 - NFR 2B10a.
	060310 Asphalt blowing	NO	NO	NO	-	Not occurring in Wallonia.
	060311 Adhesive, magnetic tapes, films and photographs	1481	1013	331	Tier 3	Use of facility data
	060312 Textile finishing	NE	NE	NE	-	In Wallonia, this activity is not significant. The NMVOC emissions are not estimated.
	060313 Leather tanning	15	19	24	Tier 3	Use of facility data
	060314 Other		75	2	Tier 3	Use of facility data
<b>2D3h</b>	<b>Printing</b>	<b>1339</b>	<b>1223</b>	<b>787</b>		
	060403 Printing industry	1339	1223	787	Tier 2	Part of the emissions comes from facility data. Global ink sales, proportion of each printing technique and average solvent content of the ink for each printing technique are used to calculate the rest of the emissions.
<b>2D3i</b>	<b>Other solvent use (please specify in the IIR)</b>	<b>NO</b>	<b>NO</b>	<b>NO</b>		
<b>2G</b>	<b>Other product use (please specify in the IIR)</b>	<b>1013</b>	<b>888</b>	<b>805</b>		
	060401 Glass wool enduction	NE	NE	NE	-	In Wallonia, this activity is not significant. The NMVOC emissions are not estimated.
	060402 Mineral wool enduction	NE	NE	NE	-	In Wallonia, this activity is not significant. The NMVOC emissions are not estimated.
	060404 Fat, edible and non-edible oil extraction	IE	IE	IE	-	Emissions of one producer are reported SNAP 060314 - NFR 2D3g.
	060405 Application of glues and adhesives	339	336	335	Tier 2	Detailed information are available for the year 2002. Same emissions have been assumed for the recent years.
	060406 Preservation of wood	672	550	469	Tier 3	Use of facility data
	060407 Underseal treatment and conservation of vehicles	NO	NO	NO	-	Not occurring in Wallonia (this activity is not present any more).
	060409 Vehicles dewaxing	0	0	0	-	In Wallonia, this activity is not present any more.
	060411 Domestic use of pharmaceutical products	NE	NE	NE	-	These emissions are not estimated in Wallonia.
	060412 Other (preservation of seeds,...)	2	2	2	Tier 3	Use of facility data

## BRUSSELS CAPITAL REGION

NMVOC emissions (tonnes/year) - codes NFR 2014		2005	2010	2013	Tier method for 2013	Comment
<b>2D</b>		<b>7794</b>	<b>8279</b>	<b>8451</b>		
<b>2D3a</b>	<b>Domestic solvent use including fungicides</b>	<b>7792</b>	<b>8278</b>	<b>8450</b>		
	Domestic solvent use (other than paint application)	1.191	1.328	1.407	Tier 2	Emissions are calculated considering population, expenses on solvent products and emission factors.
<b>2D3d</b>	<b>Coating applications</b>	<b>2</b>	<b>1</b>	<b>1</b>		
	Manufacture of automobile	0.757	0.192	0.352	Tier 3	Data is coming directly from the companies (NMVOC balance)
	Paint application: car refinishing	0.035	0.017	0.017	Tier 2	Based on average emissions from car repairing establishments. AD is the number of enterprises.
	Domestic use	0.993	0.451	0.397	Tier 2	Emissions are calculated considering paint sales, solvent content and emission factors.
<b>2D3e</b>	<b>Degreasing</b>	<b>0</b>	<b>0</b>	<b>0</b>		
	Metal degreasing	0.013	0.014	0.026	Tier 3	Data is coming directly from the companies (NMVOC balance)
<b>2D3f</b>	<b>Dry cleaning</b>	<b>0</b>	<b>0</b>	<b>0</b>		
	Dry cleaning	0.005	0.006	0.006	Tier 2	Based on average emissions from car repairing establishments. AD is the number of enterprises.
<b>2D3h</b>	<b>Printing</b>	<b>1</b>	<b>0</b>	<b>0</b>		
	Printing industry	0.725	0.233	0.218	Tier 3/Tier 2	Tier 3: Use of facility data; Tier 2:Based on average emissions from car repairing establishments. AD is the number of enterprises.