



2023

INFORMATIVE INVENTORY REPORT OF GEORGIA 1990-2021

Ministry of Environmental Protection and Agriculture of Georgia
Ambient Air Division

Submitted under the Convention on Long-Range Transboundary Air Pollution

LIST OF ABBREVIATIONS

MEPA	– Ministry of Environmental Protection and Agriculture of Georgia
EMEP	– The European Monitoring and Evaluation Programme
EEA	– European Economic Area
GEOSTAT	– National Statistics Office of Georgia
IPCC	– Intergovernmental Panel on Climate Change
CLRTAP	– Convention on Long-Range Transboundary Air Pollution
COPERT 4	– Road transport database
CNG	– Compressed natural gas
IIR	– Informative Inventory Report (UNECE)
LPS	– Large point sources, equals to the definition of E-PRTR installations
NFR	– Nomenclature for reporting (IPCC code of categories)
QA/QC	– Quality assurance/quality control:
UNECE	– United Nations Economic Commission for Europe

Pollutants

As	– Arsenic
Cd	– Cadmium
Cr	– Chromium
Cu	– Copper
CO	– Carbon monoxide
HCB	– Hexachlorobenzene
Hg	– Mercury
HM	– Heavy metals
NH ₃	– Ammonia
Ni	– Nickel
NMVOC	– Non-methane volatile organic compounds
NO ₂	– Nitrogen dioxide
NO _x	– Nitrogen oxides, nitric oxide and nitrogen dioxide, expressed as nitrogen dioxide
PAH	– Polyaromatic hydrocarbons expressed as the sum of benzo(a)pyrene, benzo(b)fluoranthene, benzo(k)fluoranthene and indeno(1,2,3,-cd)pyrene
Pb	– Lead
PCDD/PCDF	– Dioxins and furans: 1,2,3,7,8-PeCDD; 2,3,4,7,8-PeCDF; 1,2,3,4,7,8-HxCDF; 1,2,3,6,7,8-HxCDF

PCB	– Polychlorinated biphenyls
PCP	– Pentachlorophenol
PFCs	– Perfluorocarbons
PM2.5	– Particulate matter; particles on the order of ~ 2.5 micrometers or less
PM10	– Particulate matter; particles on the order of ~10 micrometers or less
POP	– Persistent organic pollutants
Se	– Selenium
SO ₂	– Sulphur dioxide
SO _x	– Sulphur oxides, all sulphur compounds expressed as sulphur dioxide
TSP	– Total suspended particulates
Zn	– Zinc

Contents

Executive Summary	4
1. Introduction	7
1.1. National Inventory Background	7
1.2. Institutional Arrangements	7
1.3. Inventory preparation process	8
1.4. Methods and data sources	9
1.5. Key categories	9
1.6. QA/QC and verification methods	24
1.7. General assessment of completeness	24
List of notation keys	24
Sources not estimated	25
Sources included elsewhere	27
2. Explanation of key trends	28
Nitrogen Oxides	29
Non-methane volatile compounds	29
Sulphur Dioxide	30
Ammonia	30
Particulates	31
Black Carbon	32
Carbon Monoxide	33
Priority Heavy Metals	34
Additional Heavy Metals	35
Dioxins/ Furans	36
PAHs / benzo(a)pyrene	36
POPs / HCB and PCBs	37
3. Energy (NFR sector 1)	37
Energy industries (1A1)	38
Manufacturing industries and construction (1A2)	39
Transport (1A3)	40
Small combustion (1A4)	41
Fugitive emissions from fuels (1B)	45
4. Industrial processes and product use (NFR sector 2)	45
Mineral Products (2A)	46
Chemical Industry (2B)	47
Metal Production (2C)	47
Solvents (2D)	49
Other (2H, 2I and 2K)	49
5. Agriculture (NFR sector 3)	51
Manure Management (3B)	52
Agricultural Soils (3D)	52
6. Waste (NFR sector 5)	55
Solid waste disposal on land (5A)	55
Waste incineration (5C)	56
Wastewater handling (5D)	56
7. Recalculations and improvements	58
Recalculations	58
Planned improvements	58
8. IIR References	58

Executive Summary

Georgia is a party to the 1979 Geneva Convention on Long-range Transboundary Air Pollution since 1999. The present report is the sixth Informative Inventory Report (IIR) submitted by Georgia under the Convention on Long-Range Transboundary Air Pollution. The first IIR was submitted in 2015. The report provides background information on Georgia's emission inventory data.

Georgia reports emissions of NO_x, NMVOC, SO₂, NH₃, PM_{2.5}, PM₁₀, TSP, BC, CO, Pb, Cd, Hg, As, Cr, Cu, Ni, Se, Zn, PCDD/ PCDF, benzo(a) pyrene, benzo(b) fluoranthene, benzo(k) fluoranthene, Indeno (1,2,3-cd) pyrene, HCB, PCBs, in the following sectors: Energy, Industrial Processes and Product Use, Agriculture and Waste. Georgia also reports emission data from large point sources. The report covers period from 1990 to 2021.

The pollutants reported by Georgia show the following trends:




Figure 1.1 Trends of main pollutants, 1990-2021




Figure 1.2 Trends of particulate matters, 1990-2021




Figure 1.3 Trend of Carbon Monoxide (CO), 1990-2021




Figure 1.4 Trends of priority heavy metals, 1990-2021




Figure 1.5 Trends of additional heavy metals, 1990-2021




Figure 1.6 Trend of PCDD/ PCDF, 1990-2021




Figure 1.7 Trends of PAHs, 1990-2021




Figure 1.8 Trends of HCB and PCBs, 1990-2021

1. Introduction

1.1. National Inventory Background

Georgia joined the Convention on Long-Range Transboundary Pollution in 1999. Georgia annually provides a national inventory of air pollutants. The following pollutants are covered:

Table 1.1 List of pollutants by sector

Sector	Pollutant / 1990-2021					
Energy	Main Pollutants	PM	CO	Priority Heavy Metals	Additional Heavy Metals	POPs
Industrial Processes and Product Use	Main Pollutants	PM	CO	Priority Heavy Metals	Additional Heavy Metals	POPs
Agriculture	Main Pollutants ¹	PM ²				
Waste	Main Pollutants	PM	CO	Priority Heavy Metals	Additional Heavy Metals ³	

1.2. Institutional Arrangements

In Georgia, the Ministry of Environmental Protection and Agriculture (MEPA) is responsible for preparation of the inventory. This task is located within the Ambient Air Division, which collects activity data from the National Statistics Office of Georgia (GEOSTAT), National Environmental Agency (NEA) and various companies. MEPA carries out the emission calculation based on the collected data. Quality checking/control is also carried out by MEPA. MEPA is responsible for reporting emission data to the UNECE as well. The responsibilities for preparing the inventory are described in the following figure.




Figure 1.9 Responsibilities for preparing of emission inventory

¹ Except for SO_x

² Except for BC

³ Except for Se and Zn

1.3. Inventory preparation process

In the first step of inventory preparation, MEPA obtains data from GEOSTAT and State Registry of Emissions from Stationary Pollution Sources that is managed by the Department of Environmental Assessment of NEA. The same Department provides data on wastewater handling. GEOSTAT on its own collects data from various state institution and companies.

Emissions are calculated based on the standard methods and procedures, such as EMEP/EEA Guidebook, national methodology, country-specific emission factors.

Activity data and emission factors are stored in Excel files. Data is backed-up and archived at MEPA's Ambient Air Division in different computers and virtual server.




Figure 1.10 Emission inventory structure

1.4. Methods and data sources

Emissions from the Transport and Agriculture sectors are calculated based on Tier 1 EMEP/EEA methodology, along with the recommended Tier 1 emission factors from GB2019. To estimate emissions from Energy and Industrial sectors Tier 1 and Tier 2 EMEP/EEA methodology were used. For the subcategory 2A6 (concrete and brick production) a national methodology⁴ is applied. Emissions from Waste sector Tier 1 EMEP/EEA methodology and plant specific emissions from state reporting system.

1. ENERGY

1A1a Tier 1/2 method, EMEP/EEA Guidebook – 2019⁵.

1A2b, 1A2d, 1A2e, 1A2f: Tier 2 method, EMEP/EEA Guidebook - 2019.

1A3b (i-vi), 1A3c and 1A3dii: Tier 1 method, EMEP/EEA Guidebook – 2019.

1A2a, 1A4ai⁶, 1A4bi, 1A4ci, 1A4cii, 1B1a, 1B1b, 1B2ai, 1B2aiv, 1B2av, 1B2b, 1B2c: Tier 1 method, EMEP/EEA Guidebook – 2019.

2. INDUSTRIAL PROCESSES AND PRODUCT USE

2A1, 2A2, 2A3, 2A5a, 2B1, 2B2, 2C1, 2C2, 2D3a, 2D3b, 2D3d⁷, 2H1, 2I, 2K: Tier 1 method, EMEP/EEA Guidebook - 2019.

2A6: National Methodology for emission calculation from concrete and brick production.

2B10a, 2C3, 2C5, 2H2: Tier 2 method, EMEP/EEA Guidebook - 2019.

3. AGRICULTURE

3B1a, 3B1b, 3B2, 3B3, 3B4a, 3B4d, 3B4e, 3B4gi, 3B4gii, 3B4giii, 3B4giv, 3Da1, 3Da2a⁸, 3Da3, 3Dc, 3De: Tier 1 method, EMEP/EEA Guidebook-2019.

4. WASTE

5A, 5D1, 5D2: Tier 1 method, EMEP/EEA Guidebook - 2019.

5C1bi, 5C1biii: Tier 1 and 3 (plant specific emissions from state reporting system) method, EMEP/EEA Guidebook - 2019.

Data sources for the inventory mainly comes from GEOSTAT. Data on fuel consumption in 1990-2018 was obtained from International Energy Agency. State Registry of Emissions from Stationary Pollution Sources, which is managed and verified by the Department of Environmental Assessment of NEA, provides information on industrial activity in certain economic sectors. NEA also provides data on wastewater handling. Data on CH₄ emissions from solid waste disposal on land were obtained from Georgia's Biennial Update Reports to the UNFCCC.

1.5. Key categories

This chapter presents the results of key sources analyses.

It is good practice for each country to identify its national key categories in a systematic and objective way. This can be achieved by a quantitative analysis of the relationship between the magnitude of

⁴ N435 Order of the Government on instrumental method for determination of actual amounts of emissions into ambient air from stationary pollution source, standard list of emission measuring equipment, and methodology for calculation of actual amounts of emissions into ambient air from stationary pollution source according to technological processes (31/12/13)

⁵ Only Tier 2 methods are applied for emissions since 2012

⁶ Tier 2 methods are applied for Ni

⁷ 1990-2008 have not been estimated due to lack of statistic data

⁸ 1990-2013 have not been estimated due to lack of statistic data

emissions in any one year (level) and the change in the emissions year to year (trend) of each category's emissions compared to the total national emissions.

Key sources analysis is prepared based on methodology described in Chapter 2 of the EMEP/EEA air pollutant emission inventory Guidebook 2019. The methodology covers Approaches 1 and 2 for both level and trend assessment. Both approaches identify key categories in terms of their contribution to the absolute level of the national emissions.

In Approach 1: the key categories are identified using a predetermined cumulative emissions threshold. Key categories are those which, when summed together in descending order of magnitude, cumulatively add up to 80% of the total level.

In Approach 2: the key categories can be derived by inventory compilers if category uncertainties or parameter uncertainties are available. Under Approach 2 the categories are sorted according to their contribution to uncertainty.

For identification of the key categories for level and trend assessment, approach 1 has been selected.

Level assessment

The contribution of each source category to the total national inventory level is calculated according to equation (1) (level assessment (Approach 1)):

Key category level assessment = source category estimate / total contribution

$$L_{x,t} = E_{x,t} / \sum E_t$$

Where:

$L_{x,t}$ = level assessment for source x in latest inventory year (year Gg)

$E_{x,t}$ = value of emission estimate of source category x in year Gg

$\sum E_t$ = total contribution, which is the sum of the emissions in year Gg, calculated using the aggregation level chosen by the country for key category analysis.

Key categories according to equation (1) are those that, when summed together in descending order of magnitude, add up to 80 % of the sum of all $L_{x,t}$. Tables 1.2 - 1.26 present the source category, sorted by largest contribution to national total.

Table 1.2 Key categories for NO_x emissions for the year 2021

Level Assessment					
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A3biii	Road transport: Heavy duty vehicles and buses	12.079	30.9%	30.9%	
1A3bi	Road transport: Passenger cars	6.666	17.1%	48.0%	
2B2	Nitric acid production	3.804	9.7%	57.8%	
3Da2a	Animal manure applied to soils	3.643	9.3%	67.1%	
1A4bi	Residential: Stationary	2.732	7.0%	74.1%	
3Da1	Inorganic N-fertilizers (includes also urea application)	2.680	6.9%	81.0%	
Trend Assessment					
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	27.68	0.843	0.667	42.9% 42.9%
1A3biii	Road transport: Heavy duty vehicles and buses	13.02	12.079	0.270	17.4% 60.3%
1A3bi	Road transport: Passenger cars	5.68	6.666	0.188	12.1% 72.4%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	1.96	2.167	0.058	3.7% 76.2%
1A4ai	Commercial/institutional: Stationary	3.46	0.683	0.054	3.5% 79.7%

Table 1.3 Key categories for NMVOC emissions for the year 2021

Level Assessment					
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A3bi	Road transport: Passenger cars	7.300	18.1%	18.1%	
1A4bi	Residential: Stationary	5.961	14.8%	32.9%	
2D3d	Coating applications	5.487	13.6%	46.5%	
2D3a	Domestic solvent use including fungicides	4.474	11.1%	57.6%	
3B1a	Manure management - Dairy cattle	3.635	9.0%	66.6%	
2H2	Food and beverages industry	3.061	7.6%	74.2%	
1A3bv	Road transport: Gasoline evaporation	2.634	6.5%	80.7%	
Trend Assessment					
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend Cumulative Total of L _{x,t}
2H2	Food and beverages industry	10.03	3.061	0.146	25.5% 25.5%
1A3bi	Road transport: Passenger cars	6.45	7.300	0.085	14.8% 40.3%

Informative Inventory Report of Georgia 2023

1A3bv	Road transport: Gasoline evaporation	1.00	2.634	0.063	11.1%	51.3%
1B1a	Fugitive emission from solid fuels: Coal mining and handling	2.09	0.184	0.046	8.0%	59.3%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	1.70	0.184	0.036	6.3%	65.5%
3B4gii	Manure magement - Broilers	2.97	1.1	0.032	5.6%	71.1%
1A4bi	Residential: Stationary	8.92	15.1	0.021	3.7%	74.9%
3B3	Manure management - Swine	0.86	0.4	0.019	3.3%	78.2%
5A	Biological treatment of waste - Solid waste disposal on land	0.49	0.7	0.016	2.8%	81.0%

Table 1.4 Key categories for SO_x emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	4.780		80.3%	80.3%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	82.05	0.005	13.765	44.3%	44.3%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	7.55	4.780	12.992	41.8%	86.1%

Table 1.5 Key categories for NH₃ emissions for the year 2021

Level Assessment				
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}
3B1a	Manure management - Dairy cattle	8.605	25.4%	25.4%
3Da2a	Animal manure applied to soils	8.086	23.9%	49.3%
3Da3	Urine and dung deposited by grazing animals	3.980	11.8%	61.1%
3B1b	Manure management - Non-dairy cattle	3.243	9.6%	70.7%
5D1	Domestic wastewater handling	2.893	8.6%	79.3%
3Da1	Inorganic N-fertilizers (includes also urea application)	2.185	6.5%	85.7%
Trend Assessment				

Informative Inventory Report of Georgia 2023

NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
3B3	Manure management - Swine	4.87	0.734	0.112	34.6%	34.6%
3B1a	Manure management - Dairy cattle	10.51	8.605	0.066	20.6%	55.3%
3B4gii	Manure management - Broilers	4.90	2.046	0.055	17.1%	72.3%
3Da2a	Animal manure applied to soils	13.15	8.086	0.034	10.6%	82.9%

Table 1.6 Key categories for PM_{2.5} emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
1A4bi	Residential: Stationary	7.303	72.1%	72.1%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.574	5.7%	77.7%		
1A3bvi	Road transport: Automobile tyre and brake wear	0.407	4.0%	81.7%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	5.33	0.004	0.525	43.3%	43.3%
1A4bi	Residential: Stationary	10.86	7.303	0.342	28.2%	71.5%
1A3bvi	Road transport: Automobile tyre and brake wear	0.08	0.407	0.071	5.9%	77.4%
1A3bvii	Road transport: Automobile road abrasion	0.05	0.249	0.044	3.6%	81.0%

Table 1.7 Key categories for PM₁₀ emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
1A4bi	Residential: Stationary	7.499	59.4%	59.4%		
2A5a	Quarrying and mining of minerals other than coal	0.999	7.9%	67.3%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.622	4.9%	72.2%		
2A2	Lime production	0.444	3.5%	75.7%		
2A1	Cement production	0.340	2.7%	78.4%		
1A3biii	Road transport: Heavy duty vehicles and buses	0.340	2.7%	81.1%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}

Informative Inventory Report of Georgia 2023

1A1a	Public electricity and heat production	3.85	0.004	0.304	35.0%	35.0%
1A4bi	Residential: Stationary	11.15	7.499	0.197	22.6%	57.6%
2A5a	Quarrying and mining of minerals other than coal	1.32	0.999	0.039	4.5%	62.1%
3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products	1.09	0.330	0.039	4.5%	66.6%
2C2	Ferroalloys production	0.06	0.303	0.039	4.5%	71.0%
1A4ai	Commercial/institutional: Stationary	0.44	0.1	0.034	3.9%	74.9%
1A3bvi	Road transport: Automobile tyre and brake wear	0.15	0.1	0.033	3.8%	78.7%
2A2	Lime production	0.44	0.0	0.029	3.3%	82.0%

Table 1.8 Key categories for TSP emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
1A4bi	Residential: Stationary	7.890	38.7%	38.7%		
2A6	Other mineral products (please specify in the IIR)	3.429	16.8%	55.5%		
2A5a	Quarrying and mining of minerals other than coal	2.038	10.0%	65.5%		
2A2	Lime production	1.086	5.3%	70.8%		
3B4gii	Manure mangement - Broilers	1.046	5.1%	76.0%		
2D3b	Road paving with asphalt	0.659	3.2%	79.2%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.659	3.2%	82.4%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
2A6	Other mineral products (please specify in the IIR)	1.94	3.429	0.169	24.7%	24.7%
1A1a	Public electricity and heat production	3.01	0.004	0.147	21.5%	46.2%
2I	Wood processing	1.32	0.181	0.051	7.4%	53.6%
3B4gii	Manure mangement - Broilers	2.50	1.046	0.042	6.1%	59.7%
1A4bi	Residential: Stationary	11.74	7.890	0.033	4.8%	64.5%
2A2	Lime production	1.09	0.1	0.030	4.4%	68.9%
3B3	Manure management - Swine	0.78	0.4	0.029	4.2%	73.1%
3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products	1.09	1.0	0.028	4.1%	77.2%
2A5a	Quarrying and mining of minerals other than coal	2.69	0.0	0.025	3.7%	80.9%

Informative Inventory Report of Georgia 2023

Table 1.9 Key categories for BC emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A4bi	Residential: Stationary	0.728		66.4%	66.4%	
1A3biii	Road transport: Heavy duty vehicles and buses	0.180		16.4%	82.8%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A4bi	Residential: Stationary	1.08	0.728	0.161	22.3%	22.3%
1A1a	Public electricity and heat production	0.16	0.000	0.148	20.5%	42.8%
1A4ai	Commercial/institutional: Stationary	0.15	0.000	0.136	18.8%	61.6%
1A3biii	Road transport: Heavy duty vehicles and buses	0.19	0.180	0.106	14.7%	76.3%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	0.07	0.008	0.053	7.4%	83.7%

Table 1.10 Key categories for CO emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A3bi	Road transport: Passenger cars	58.597		52.7%	52.7%	
1A4bi	Residential: Stationary	40.343		36.3%	88.9%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A3bi	Road transport: Passenger cars	53.59	58.597	0.179	45.3%	45.3%
1A4bi	Residential: Stationary	61.16	40.343	0.095	24.0%	69.4%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	4.75	0.516	0.037	9.4%	78.7%
1A4ai	Commercial/institutional: Stationary	2.11	0.268	0.016	4.0%	82.8%

Table 1.11 Key categories for Pb emissions for the year 2021

Level Assessment					
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] $E_{x,t}$		Level Assessment $L_{x,t}$	Cumulative Total of $L_{x,t}$
2C1	Iron and steel production	1.357		49.9%	49.9%

Informative Inventory Report of Georgia 2023

1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.712	26.2%	76.1%		
1A4bi	Residential: Stationary	0.265	9.7%	85.8%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	1.12	0.712	0.676	29.0%	29.0%
2C1	Iron and steel production	6.75	1.357	0.409	17.5%	46.5%
1A4cii	Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	0.97	0.009	0.345	14.8%	61.3%
1A1a	Public electricity and heat production	0.79	0.000	0.290	12.5%	73.8%
1A4bi	Residential: Stationary	0.44	0.265	0.241	10.3%	84.1%

Table 1.12 Key categories for Cd emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
1A4bi	Residential: Stationary	0.127	71.4%	71.4%		
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.012	6.5%	77.9%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.010	5.4%	83.3%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	0.20	0.000	1.111	41.2%	41.2%
1A4bi	Residential: Stationary	0.19	0.127	0.996	36.9%	78.1%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.01	0.012	0.147	5.4%	83.5%

Table 1.13 Key categories for Hg emissions for the year 2021

Level Assessment				
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.071	29.1%	29.1%
5C1biii	Clinical waste incineration	0.050	20.5%	49.7%

Informative Inventory Report of Georgia 2023

1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.042	17.3%	67.0%		
2K	Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)	0.037	15.2%	82.2%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
2C1	Iron and steel production	0.15	0.030	0.380	31.3%	31.3%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.04	0.071	0.350	28.8%	60.1%
1A1a	Public electricity and heat production	0.08	0.002	0.306	25.1%	85.2%

Table 1.14 Key categories for As emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
2C1	Iron and steel production	0.118	58.3%	58.3%		
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.039	19.0%	77.4%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.021	10.5%	87.9%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	0.71	0.002	3.420	49.0%	49.0%
2C1	Iron and steel production	0.59	0.118	1.241	17.8%	66.8%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.02	0.039	1.235	17.7%	84.5%

Table 1.15 Key categories for Cr emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
2C1	Iron and steel production	1.328		76.8%	76.8%	
1A4bi	Residential: Stationary	0.225		13.0%	89.9%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}

Informative Inventory Report of Georgia 2023

1A4bi	Residential: Stationary	0.34	0.225	0.403	29.4%	29.4%
2C1	Iron and steel production	6.60	1.328	0.301	22.0%	51.3%
1A1a	Public electricity and heat production	0.44	0.000	0.257	18.8%	70.1%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.04	0.060	0.137	10.0%	80.1%

Table 1.16 Key categories for Cu emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.094	29.4%	29.4%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.093	29.1%	58.5%		
1A4bi	Residential: Stationary	0.059	18.4%	76.9%		
1A3bi	Road transport: Passenger cars	0.028	8.9%	85.8%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	0.91	0.000	2.850	44.4%	44.4%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.06	0.094	1.229	19.2%	63.6%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.15	0.093	0.935	14.6%	78.1%
1A4bi	Residential: Stationary	0.10	0.059	0.579	9.0%	87.2%

Table 1.17 Key categories for Ni emissions for the year 2021

Level Assessment				
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.071	28.0%	28.0%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.069	27.1%	55.2%
2C1	Iron and steel production	0.041	16.2%	71.4%
2A3	Glass production	0.031	12.2%	83.6%
Trend Assessment				

Informative Inventory Report of Georgia 2023

NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] $E_{x,0}$	Latest Year (2021) Estimate [Gg] $E_{x,t}$	Trend Assessment $L_{x,t}$	% Contribution to the trend	Cumulative Total of $L_{x,t}$
1A1a	Public electricity and heat production	39.90	0.000	156.863	47.0%	47.0%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.04	0.071	47.285	14.2%	61.2%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.11	0.069	45.559	13.7%	74.9%
2C1	Iron and steel production	0.21	0.041	26.709	8.0%	82.9%

Table 1.18 Key categories for Se emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate		Level Assessment	Cumulative Total of	
		[Gg] E _{x,t}		L _{x,t}	L _{x,t}	
2A3	Glass production	0.051		46.2%	46.2%	
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.037		33.5%	79.7%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.010		8.7%	88.4%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A1a	Public electricity and heat production	0.56	0.000	5.083	47.1%	47.1%
1A2f	Stationary combustion in manufacturing industries and construction: Non-metallic minerals	0.02	0.037	2.468	22.9%	70.0%
2A3	Glass production	0.16	0.051	2.239	20.8%	90.7%

Table 1.19 Key categories for Zn emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A4bi	Residential: Stationary	5.016		48.8%	48.8%	
1A3bi	Road transport: Passenger cars	1.494		14.5%	63.4%	
2C1	Iron and steel production	1.180		11.5%	74.9%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	1.063		10.3%	85.2%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}

Informative Inventory Report of Georgia 2023

1A1a	Public electricity and heat production	13.92	0.000	1.355	40.9%	40.9%
1A4bi	Residential: Stationary	7.44	5.016	0.838	25.3%	66.1%
1A3bi	Road transport: Passenger cars	1.37	1.494	0.332	10.0%	76.2%
2C1	Iron and steel production	5.87	1.180	0.204	6.1%	82.3%

Table 1.20 Key categories for PCDD/ PCDF emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A4bi	Residential: Stationary	7.905		77.8%	77.8%	
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	1.078		10.6%	88.5%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A4bi	Residential: Stationary	11.94	7.905	0.280	45.1%	45.1%
2C1	Iron and steel production	4.40	0.885	0.270	43.6%	88.7%

Table 1.21 Key categories for benzo(a)pyrene emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A4bi	Residential: Stationary	1.185		82.1%	82.1%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A4bi	Residential: Stationary	1.83	1.185	0.122	64.5%	64.5%
1A4ai	Commercial/institutional: Stationary	0.03	0.000	0.023	12.3%	76.9%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.38	0.242	0.021	11.0%	87.8%

Table 1.22 Key categories for benzo(b)fluoranthene emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A4bi	Residential: Stationary	1.087		76.0%	76.0%	

Informative Inventory Report of Georgia 2023

1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.316	22.0%	98.0%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A4bi	Residential: Stationary	1.73	1.087	0.142	57.2%	57.2%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.51	0.316	0.040	15.9%	73.1%
1A4ai	Commercial/institutional: Stationary	0.05	0.000	0.033	13.2%	86.3%

Table 1.23 Key categories for benzo(k)fluoranthene emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}	Level Assessment L _{x,t}	Cumulative Total of L _{x,t}		
1A4bi	Residential: Stationary	0.412	73.6%	73.6%		
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.127	22.7%	96.3%		
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A4bi	Residential: Stationary	0.66	0.412	0.149	53.3%	53.3%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.20	0.127	0.053	19.0%	72.3%
1A4ai	Commercial/institutional: Stationary	0.02	0.000	0.031	11.1%	83.4%

Table 1.24 Key categories for Indeno(1,2,3-cd)pyrene emissions for the year 2021

Level Assessment						
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] E _{x,t}		Level Assessment L _{x,t}	Cumulative Total of L _{x,t}	
1A4bi	Residential: Stationary	0.696		86.0%	86.0%	
Trend Assessment						
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] E _{x,0}	Latest Year (2021) Estimate [Gg] E _{x,t}	Trend Assessment L _{x,t}	% Contribution to the trend	Cumulative Total of L _{x,t}
1A4bi	Residential: Stationary	1.07	0.696	0.097	66.5%	66.5%
1A4ai	Commercial/institutional: Stationary	0.01	0.000	0.017	11.5%	78.0%
1A2a	Stationary combustion in manufacturing industries and construction: Iron and steel	0.16	0.099	0.009	6.0%	84.0%

Table 1.25 Key categories for HCB emissions for the year 2021

Level Assessment					
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] $E_{x,t}$		Level Assessment $L_{x,t}$	Cumulative Total of $L_{x,t}$
2C3	Aluminum production	28.938		99.8%	99.8%
Trend Assessment					
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] $E_{x,0}$	Latest Year (2021) Estimate [Gg] $E_{x,t}$	Trend Assessment $L_{x,t}$	% Contribution to the trend
1A4bi	Residential: Stationary	0.07	0.049	0.002	53.8%
2C1	Iron and steel production	0.04	0.009	0.002	33.0%

Table 1.26 Key categories for PCB emissions for the year 2021

Level Assessment					
NFR Category Code	NFR Category	Latest Year (2021) Estimate [Gg] $E_{x,t}$		Level Assessment $L_{x,t}$	Cumulative Total of $L_{x,t}$
2K	"Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)"	372.860		99.5%	99.5%
Trend Assessment					
NFR Category Code	NFR Category	Base Year (1990) Estimate [Gg] $E_{x,0}$	Latest Year (2021) Estimate [Gg] $E_{x,t}$	Trend Assessment $L_{x,t}$	% Contribution to the trend
2K	Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)	541.00	372.860	0.007	47.8%
2C1	Iron and steel production	3.67	0.738	0.007	44.2%

Trend assessment

The purpose of the trend assessment is to identify categories that may not be large enough to be identified by the level assessment, but whose trend contributes significantly to the trend of the overall inventory, and should therefore receive particular attention. The trend assessment is calculated according to equation (2) (level assessment (Approach 1)):

$$T_x = (E_x - E_{x,0}) / (\sum_i E_{i,t} - \sum_i E_{i,0})$$

Where:

T_x = trend assessment of source category x in year t as compared to the base year (year 0) or starting year of the inventory

$E_x - E_{x,0}$ = values of estimates of source category x in year t and 0 respectively

$\sum_i E_{i,t} - \sum_i E_{i,0}$ = sum of emissions across all n source categories (i = 1, ...x, n) (total inventory estimates) in years t and 0, respectively

Informative Inventory Report of Georgia 2023

The trend of a category refers to the change in the source category emissions over time. It is computed as an absolute value for the source category x by subtracting value of the base year, or the starting year (year 0), estimate for source category x from the latest inventory year (year t) estimate and dividing by the overall difference between the target year (year t) and the base year (year 0) total inventories (the inventory trend). The percentage contribution from category x for year t to the trend is then calculated by dividing T_x , by the sum of the trend assessment of all categories of the inventory.

The trend assessment then sorts categories by magnitude (highest to lowest) of their contribution to the trend, regardless of whether category trend is increasing or decreasing. Categories whose cumulative percentage contribution is greater than 80% should be identified as key. The results of trend assessment are given in second sub-tables in Tables 1.2 - 1.26.

1.6. QA/QC and verification methods

The following quality control measures are carried out:

Check for transcription errors and data comparison

For point sources, the first check is made during the approval of the submitted annual reports, and then in process of the data analysis. Statistical data is compared to data available from previous years. In case of discrepancies, data from other sources (e.g. from companies) are used. If the data available to the Ministry shows higher levels than the statistical data, the levels available to the Ministry are used.

Check of calculated emissions

A staff member who did not make a specific calculation checks the colleague's approach and results. All results are compared to the values of previous years.

In addition, the following measure is carried out:

Review of methods and emission factors

Emission factors are updated when new EMEP/EEA-Guidebooks are published. Other guidebooks are monitored. The national methodology is also updated continuously.

1.7. General assessment of completeness

List of notation keys

In the following table, notation keys are listed (as defined in the UNFCCC reporting guidelines (ECE/EB.AIR/125)):

- (a) "NE" (not estimated), for activity data and/or emissions by sources of pollutants which have not been estimated but for which a corresponding activity may occur within a Party. Where NE is used in an inventory to report emissions of pollutants, the Party should indicate why such emissions have not been estimated;
- (b) "IE" (included elsewhere), for emissions by sources of pollutants estimated but included elsewhere in the inventory instead of under the expected source category. Where IE is used in an inventory, the Party should indicate where in the inventory the emissions for the displaced source category have been included, and the Party should explain such a deviation from the inclusion under the expected category;
- (c) "C" (confidential information), for emissions by sources of pollutants of which the reporting could lead to the disclosure of confidential information. The source category where these emissions are included should be indicated;
- (d) "NA" (not applicable), for activities under a given source category that do occur within the Party but do not result in emissions of a specific pollutant;
- (e) "NO" (not occurring), for categories or processes within a particular source category that do not occur within a Party;
- (f) "NR" (not relevant). According to paragraph 37 in the Guidelines, emission inventory reporting for the main pollutants should cover all years from 1990 onwards if data are available. However, NR is

introduced to ease the reporting where reporting of emissions is not strictly required by the different protocols, e.g., emissions for some Parties prior to agreed base years.

Sources not estimated

The following categories have not been estimated:

List of important sectors with “NE” and short justification why these sectors have not been estimated.

Table 1.14 Sources not estimated (NE)

NFR14 code	Substance(s)	Reason for not estimated
1A1a	NH ₃ , PCBs, PCDD/F, HCB	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
1A2a	NH ₃	
1A2b	All except for NO _x and SO _x	
1A2c	All	Emission occur, but have not been estimated due to lack of statistic data and emission factors in national methodology
1A2d	NH ₃	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
1A2f	NH ₃	
1A3ai(i)	NO _x , NMVOC, SO _x , PMs CO	Emission occur, but have not been estimated due to lack of statistic data
1A3aii(i)	NO _x , NMVOC, SO _x , PMs CO	
1A3bi	Hg, As, PCDD/F, HCB, PCBs	Emissions occur, but have not been estimated due to lack of emission factors in methodology (COPERT 4 version v11.2 fills NFR table (including notation keys NE)).
1A3bii	Hg, As, PCDD/F, HCB, PCBs	
1A3biii	Hg, As, PCDD/F, HCB, PCBs	
1A3biv	All	Emission occur, but have not been estimated due to lack of statistic data and emission factors in methodology (COPERT 4 version v11.2 fills NFR table (including notation keys NE)).
1A3bv	POPs except for total PAHs	Emissions occur, but have not been estimated due to lack of emission factors in methodology (COPERT 4 version v11.2 fills NFR table (including notation keys NE)).
1A3bvi	HMs, PAHs, HCB, PCBs	
1A3bvii	HMs, PAHs, HCB, PCBs	
1A3c	BC, Pb, Hg, As, PCDD/PCDF, Benzo(k)fluoranthene, Indeno(1,2,3-cd)pyrene	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
1A3dii	NH ₃	
1A3ei	All	Emission occur, but have not been estimated due to lack of statistic data
1A4aii	All	
1A4bii	All	
1A4cii	Hg, As, PCDD/PCDF, Benzo(k)fluoranthene, Indeno(1,2,3-cd)pyrene, HCB, PCBs	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)

1A4ciii	All	
1B1a	BC, HMs	
1B2ai	SOx, PCDD/PCDF	
1B2av	SOx, PCDD/PCDF	
1B2b	SOx, PCDD/PCDF	
1B2c	NH ₃ , PMs, HMs, PCDD/F, PAHs	
2A1	All except for PMs and PCBs	
2A2	NOx, NMVOC, SOx, CO, Pb, Cd, Hg	
2A3	Main Pollutants, CO, POPs except for PCBs	Emission occur, but have not been estimated due to lack of statistic data
2A5b	TSP, PM _{2.5} , PM ₁₀	
2A5c	TSP, PM _{2.5} , PM ₁₀	
2A6	PM _{2.5} , PM ₁₀ , BC, benzo(a) pyrene	
2B1	NMVOC, SOx, PM _{2.5}	
2B2	NH ₃ , PM _{2.5}	
2B10a	All except for NH ₃ and TSP	
2C1	NH ₃ , PAHs except for total	Emission occur, but have not been estimated due to lack of emission factors in national methodology
2C2	Main Pollutants, CO, HMs, PCDD/F, PAHs	
2C3	Main Pollutants, CO, HMs, PAHs	
2C5	NMVOC, NH ₃ , BC, CO, Hg, Cr, Cu, Ni, Se, PAHs, HCB	
2C7d	TSP, PM _{2.5} , PM ₁₀	
2D3a	TSP, PM _{2.5} , PM ₁₀	
2D3b	NOx, SOx, CO, POPs except for PCBs	
2D3c	NOx, NMVOC, PMs, CO, Pb, Cd, Hg, POPs except for PCBs	
2D3e	NMVOC, PM _{2.5}	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
2D3f	NMVOC, PM _{2.5}	
2D3g	All	
2D3h	NMVOC, PM _{2.5} , BC	
2D3i	All	
2G	All	
2H1	NH ₃ , PAHs, HCB	
2H2	PMs	
2I	Main Pollutants, CO, PM ₁₀ , PM _{2.5} , BC, As, Cu	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
2K	HMs except for Hg, HCB	
3B4f	NOx, NMVOC, NH ₃ , PM _{2.5} , PM ₁₀ , TSP	Emission occur, but have not been estimated due to lack of statistic data

3Da1	PM _{2.5} , PM ₁₀ , TSP	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
3Da2b	NH ₃ , NOx	Emission occur, but have not been estimated due to lack of statistic data
3Da2c	NH ₃ , NOx	
3Da3	NOx	
3Da4	NH ₃ , NOx	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
3Db	NOx	
3De	NH ₃	
3Df	HCB	Emission occur, but have not been estimated due to lack of statistic data
3F	All	Emission occur, but have not been estimated due to lack of statistic data and emission factors in methodology (EMEP-EEA guidebook – 2019)
3I	NH ₃	Emission occur, but have not been estimated due to lack of statistic data
5A	NH ₃ , PM _{2.5} , PM ₁₀ , TSP, CO, Hg	Emission occur, but have not been estimated due to lack of statistic data and emission factors in methodology (EMEP-EEA guidebook – 2019)
5B1	Main Pollutants, PMs, CO	
5C1a	All	Emission occur, but have not been estimated due to lack of statistic data
5C1bi	NH ₃ , Cd, Cu, Ni, Se, Zn, POPs except for PCBs	Emission occur, but have not been estimated due to lack of emission factors in national or international methodology (EMEP-EEA guidebook – 2019)
5C1bii	All except for PCBs	Emission occur, but have not been estimated due to lack of statistic data and emission factors in methodology (EMEP-EEA guidebook – 2019)
5C1biii	NMVOC, NH ₃ , PM _{2.5} , PM ₁₀ , Se, Zn, POPs	Emission occur, but have not been estimated due to lack of statistic data and emission factors in national or international methodology (EMEP-EEA guidebook – 2019)
5C1biv	All	Emission occur, but have not been estimated due to lack of statistic data and emission factors in national or international methodology (EMEP-EEA guidebook – 2019)
5C2	All except for PCBs	
5D1	PMs, HMs	Emission occur, but have not been estimated due to lack of emission factors in methodology (EMEP-EEA guidebook – 2019)
5D2	NH ₃ , TSP, PM ₁₀ , PM _{2.5} , HMs	
5E	All	Emission occur, but have not been estimated due to lack of statistic data and emission factors in methodology (EMEP-EEA guidebook – 2019)

Sources included elsewhere

List of important categories with “IE” and short explanation in which category they are included.

Table 1.15 Sources included elsewhere (IE)

NFR14 code	Substance(s)	Included in NFR code
1A2f	PM _{2.5} , PM ₁₀ , TSP, BC	2A1, 2A2, 2A3 and 2A6
2C1	NOx, SOx, CO	1A2a

2C5	NO _x , SO _x	1A2a
3B4a	NO _x , NMVOC, NH ₃ , PM _{2.5} , PM ₁₀ , TSP	3B1b
3B4gi	NO _x , NMVOC, NH ₃ , PM _{2.5} , PM ₁₀ , TSP	3B4gii

2. Explanation of key trends

In Georgia, ambient air pollution is mainly caused by emissions from motor vehicles, the energy, industrial and agriculture sectors. Emission trends for main pollutants are presented in figure 2.1.




Figure 2.1 Main pollutants, trends over time, 1990-2021.

- The general economic activity decreased in 1990s due to economic crisis caused by dissolution of Soviet Union. Subsequently, emissions of main pollutants declined sharply. Increased economic activity from the middle of 2000s led to increased emissions of most pollutants, but these trends were disproportionate as the latter lagged behind economic trend. The main reason of this was an application of cleaner technologies that abated pollution from various sectors.
- The coronavirus (COVID19) pandemic caused negative economic effects in 2020 that resulted in reduced production and consumption as well as decreased emissions of almost all pollutants. In 2021 economic recovery triggered rising emissions compared to 2020 especially in energy and industrial sectors.
- Intense rise in emissions of NO_x from millennium was caused by rapid increase of car fleet in Georgia that remains one of the main sources of nitrogen oxides.
- Rise in emissions of NMVOC in 2009 is related to calculation of subcategory on paint (coating), application (2D3d) for which activity data before 2009 is not available. Sharp rise of emissions since 2013 was caused by introduction of national energy balance that provided slightly different and more precise activity data for residential stationary combustion as well as increased emissions from transport due to rapid expansion of car fleet in Georgia.

- Dramatic rise of SO_x emissions in 2013 was caused by increased consumption of coal for iron and steel production. Due to same reason SO_x emissions again increased in 2021.
- Trend of NH₃ emissions is mostly dependent on the production in the agricultural sector.
- A sudden growth of particulates emissions in 2013 was caused by launching of national energy balance that provided slightly different and more precise activity data for residential stationary combustion compare to the data of the International Energy Agency, which was used for all previous years. While the reduction trend since 2016 is caused by decreased consumption of firewood in households.

Nitrogen Oxides




Figure 2.2 Trend of NO_x emissions 2014-2021

Energy sector has the biggest share in total NO_x emissions (about 72%). Approximately 52% of total NO_x emissions and 72% of energy sector emissions comes from transport.

Emissions of NO_x from transport had steady decreasing trend in 2016-2020, which was caused by increasing share of new and clean transport types in import of vehicles and in the car fleet as well. The process was supported by increased taxes for the importation of fuel (petrol and diesel) and old cars and by reduction of excise duties for import of cleaner vehicles (electric/hybrid) in 2016-2017. However, in 2021, due to sharp increase of vehicles import in post pandemic period and increased share of old and petrol/diesel engine cars in the import, emissions of NO_x were raised by 5.8%. Rising emissions of NO_x from transport was counterbalanced by 9% reduction in emissions from industry due to decreased nitric acid production in the country. As a result, total emissions of NO_x increased slightly (by 3%) in 2021 compared to previous year.

Non-methane volatile compounds




Figure 2.3 Trend of NMVOC emissions 2014-2021

Increased consumption of petrol by passenger cars caused rise in emissions of NMVOC in 2016. In 2016-2020 emissions have decreased stably by 11% due to reduced emissions in energy sector, in particular reduced consumption of biomass by households. However, in 2021, emissions have raised by 4% compared to previous year. The reasons of this growth mainly are increased emissions from transport due to expanded car fleet and fuel consumption, and increased emissions from coating application due to increased amount of consumed paints.

Energy sector is the main source of pollution with NMVOC (about 47%), while road transport within energy sector accounts for 27% of total emissions.

Sulphur Dioxide




Figure 2.4 Trend of SO₂ emissions 2014-2021

Stationary combustion in manufacturing industries and construction accounted for nearly 94% of SO_x emissions in 2021.

Significant decrease in 2020 compared to 2017 was caused by diminished and zero consumption of coal for public electricity and heat production (2018-2020), iron and steel (2020), and food production (2020). In 2021 emissions were increased by 32% due to increased consumption of coal in iron and steel production.

Ammonia




Figure 2.5 Trend of NH₃ emissions 2014-2021

Substantial increase of emissions in 2015 was caused by calculation of NH_3 emissions from domestic wastewater handling which previously was not able due to absence of relevant activity data for past years. Since 2015 NH_3 emissions started decreasing from all sectors and have decreased by 15% in 4-year term. However, emissions from agricultural sector, which accounts for 87% of total NH_3 emissions, increased in 2020 by 5% due to enlarged livestock population size in the sector and slightly decreased in 2021 due to shrinking in population.

Particulates




Figure 2.6 Trend of PM2.5, PM10 and TSP emissions 2014-2021

Total emissions of PM2.5 decreased by 38% from 2014 to 2021, PM10 – by 33% and TSP – by 21%. The sharp reduction of particulate matter's emissions since 2014 was mostly achieved by decreasing biomass consumption in households.




Figure 2.7 Sources of PM2.5, PM10 and TSP emissions in 2014 and 2021

In contrast, PM emissions from industry have rising trend. TSP emissions increased by 37% in 2014-2021 mainly due to increased lime, concrete and asphalt production and quarrying of bigger amount of minerals other than coal.

Energy sector remains as a main source of PM emissions as 90% of PM_{2.5} emissions, 73% of PM₁₀ emissions and 46% of TSP emissions came from energy sector in 2021. Industry is second biggest polluter responsible for about 7% of PM_{2.5} emissions, 18% of PM₁₀ emissions and 41% of TSP emissions as of 2021. The share of energy sector in PM emissions decreases year by year due to alteration of solid fuel consumption in residential sector by natural gas, while industry's share increases as a result of increased production and lack of application of modern clean technologies in the sector.

Black Carbon




Figure 2.8 Trend of BC emissions 2014-2021

Emissions of BC have decreased since 2014 by 39% due to reduced emissions from residential combustion that is a result of wider application of cleaner energy sources and less consumption of solid fuels by households.

Energy sector is a source for about 97% of BC emissions, while only residential stationary combustion within energy sector is responsible for 66% of total emissions.

Carbon Monoxide




Figure 2.9 Trend of CO emissions 2014-2021

CO emissions were reduced by 19% in 2021 compared to 2014. The reduction of CO emissions was a result of its decreased release from residential sector as emissions from residential combustion have decreased by 48% in 2014-2021. Emissions from residential sector increased slightly in 2021.

CO emissions from transport have variable trend: it increases from 2014 to 2016 and decreases from 2017. The reduction was caused by decrease of vehicles import and growing share of new and fuel-efficient vehicles in import that was supported through economic and financial instruments since 2017. However, CO emissions from transport again increases in 2021 by 8% compared to 2020 due to rising import of old and petrol engine cars.

All in all, CO emissions were increased by 7% in 2021 compared to previous year.

Energy sector is a main source of CO emissions. In 2014 the shares of emissions from transport and residential combustion sources were 38% and 56%, since that emissions from transport outnumbered emissions from residential sources and by 2021 the shares are following: 56% to 36%.

Priority Heavy Metals




Figure 2.10 Trend of Pb emissions 2014-2021




Figure 2.11 Trend of Cd and Hg emissions 2014-2021

There are different trends among priority heavy metals emissions over 7-year period. Pb emissions increased by 33%, Hg emissions increased by 45%, while Cd emissions were reduced by 39%. The main reason for growth of Pb emissions is increased iron and steel production since 2017. Increased Hg emissions in 2017-2019 was caused by rising production of iron and steel in Georgia, while sharp rise in emission in 2020-2021 was further triggered by increased amount of medical waste incinerated. In case of Cd emissions, reduction was caused by decreased wood consumption in households that triggered shrank of emissions by 39% in 2014-2021.

The main source of Pb emissions is iron and steel production (both combustion and industry sectors) that accounted for 76% of total emissions in 2021. Emissions from energy sector decreased by 9%, while emissions from industry increased by 99% in 2021 compared to 2014.

71% of Cd emissions came from residential stationary combustion in 2021, though the share was even more in 2014 – 85%. Energy sector, mainly combustion in cement and iron and steel production, is a source for 52% of Hg emissions.

Additional Heavy Metals




Figure 2.12 Trend of additional heavy metals emissions 2014-2021

In 2021 compared to 2014 emissions of As and Cr increased by 55% and 39% respectively, while emissions of Zn and Cu decreased by 27% and 8%. Emissions of Se have variable trend as it rises in 2015-2017, reduces in 2018-2020 and again increases in 2021 due to increased glass production. Ni emissions are stable until 2021, when it increases by 9% due to increased coal consumption in iron and steel production.

Rising emissions of As and Cr is also connected with the increased production of iron and steel since 2014. Reductions of Cu and Zn emissions are the result of sharply diminished biomass consumption in residential sector. Variability of Se emissions was mainly conditioned by changeable amount of produced glass and consuming coal for public electricity and heat production in 2016-2018.




Figure 2.13 Sources of additional heavy metals emissions in 2014 and 2021

As outlined in figure 2.13 the share of energy sector in emissions of additional heavy metals is decreasing year by year, while industry gradually is becoming dominant.

Dioxins/ Furans




Figure 2.14 Trend of PCDD/ PCDF emissions 2014-2021

PCDD/ PCDF emissions were increased in 2021 compared to 2020 by 6% due to increased coal consumption in iron and steel production and slightly increased consumption of wood in residential sector in 2021.

In general, PCDD/PCDF emissions were reduced by 40% in 2021 compared to 2014. The reduction of PCDD/ PCDF emissions was a result of its decreased discharge from residential sector as emissions from residential combustion, which comprise 78% of total emissions as of 2021, have reduced by 49% in 2014-2021. Unlike energy sector, emissions of PCDD/ PCDF from industry have steady increasing trend as it increased by 113% in the mentioned period due to rise in iron and steel production, which constitutes major industrial source of PCDD/PCDF emissions in Georgia.

PAHs / benzo(a)pyrene




Figure 2.15 Trend of BaP emissions 2014-2021

Benzo(a)pyrene (BaP) is basically emitted from energy sector. BaP emissions were reduced by 43% in 2021 compared to 2014. The reduction of BaP emissions was a result of its decreased discharge from residential sector. Emissions from residential combustion, which comprise 82% of total emissions as of

2021, have decreased by 49% in 2014-2021 due to diminished firewood consumption in households. However, emissions were increased in 2021 compared to 2020 by 8% that was a result of increased coal consumption for iron and steel production and slightly increased consumption of wood in residential sector.

HCB and PCBs




Figure 2.15 Trend of HCB and PCBs emissions 2014-2021

PCBs emissions slightly increased in 2014-2021 due to rising demand by population for POPs and heavy metals. Variability of HCB emissions trend is caused by changeable amount of aluminum production in the mentioned period, which is responsible for more than 99% of emissions.

3. Energy (NFR sector 1)

Present submission includes emission data from activities 1A1a, 1A2a, 1A2b, 1A2d, 1A2e, 1A2f, 1A3b (i-iii, v-vii), 1A3c, 1A3dii, 1A4ai, 1A4bi, 1A4ci, 1A4cii, 1B1a, 1B1b, 1B2ai, 1B2aiv, 1B2av, 1B2b, 1B2c in 1990-2021⁹. Since 2013, GEOSTAT prepares yearly National Energy Balance, which gives more precise activity data for this sector.

Emissions in energy sector mainly come from fuel combustion. Minor fugitive emissions from fuel exploration generated as well. This sector covers five key activities: public electricity and heat production, combustion in manufacturing industries and construction, transport, small combustion and fugitive emissions. The energy sector is the main source of NO_x, SO₂, NMVOC, CO and PMs emissions in Georgia. In 2021, this sector contributed 72% of total NO_x emissions, 98% of total SO₂ emissions, 47% of total NMVOC emissions, 46% of total TSP emissions and 99.9% of total CO emissions.

⁹ However, some activities did not occur in some years of the reporting period including 1B1b, which did not occur since 1998.




Figure 3.1 Emissions from energy sector in 2021

Transport is the major contributor of NO_x (72.4%) and NMVOC (56.4%) emissions in the energy sector. Share of industrial combustion in total SO₂ emissions in energy sector is 94.4%. Small combustion is responsible for the 84.1% of TSP and 50.3% of NH₃ emissions in this sector.

Energy industries (1A1)

Source category description

Emissions in this category mostly come from natural gas consumption.




Figure 3.2 Emissions from public electricity and heat production 2007-2021

Temporary Increase of TSP emissions in 2016-2018 is related to consumption of coal by thermal power plant at that time.

Methodology

Emissions are estimated for 1990-2021 using EMEP/EEA Guidebook – 2019, Tier 1/2 approach based on data from International Energy Agency (1990-2018) and National Energy Balance (2019-2021).

Manufacturing industries and construction (1A2)

Source category description

This category covers emissions occurred by combustion processes in industrial sector. The main emission sources in this category are metallurgy and production of mineral materials. The category is responsible for 93.8% of total SO_x emissions in the country.




Figure 3.3 Emissions from combustion in manufacturing industries 2007-2021

Increasing trend of SO₂ emissions from 2011 is resulted from increased coal fuel consumption in industry sector (mostly in iron and steel production) and partially by rising cement production, while increase in NO_x emissions is caused mainly by expansion in cement production that more than tripled in 2010-2021, and partially by increased coal consumption in iron and steel production. Decreasing emissions of SO₂ in 2014-2016 were related to the reduced consumption of coal and heavy oil within those years, mainly caused by shifting back from coal to natural gas in iron and steel industry. However, coal consumption again increased from 2017, with an exclusion due to pandemic in 2020, and peaked in 2021 that resulted in rising emissions of SO_x. TSP and NMVOC emissions also increased in 2021 due to same reason.

Significant increase of TSP and NMVOC emissions from 2011 is also a result of increased coal fuel consumption in iron and steel production. Dramatic rise in SO₂, TSP and NMVOC emissions in 2013 is also related to introduction of national energy balance, which provides detailed information on fuel consumption in categories that was not available before.

Methodology

Emissions are estimated for 1990-2021 using EMEP/EEA Guidebook – 2019, Tier 1/2 approach based on data from International Energy Agency (1A2a, 1A2d, 1A2e - 1990-2018) and from National Energy Balance (1A2a, 1A2d, 1A2e – 2019-2021), national inventory of stationary emission sources

(1A2b, 1A2f – 2013-2021¹⁰), and production data provided by GEOSTAT and GHG emissions reports (1A2f – before 2013).

Transport (1A3)

Source category description

This category includes railways, national navigation (shipping) and all types of vehicles (passenger cars, light duty vehicles, heavy-duty trucks, buses, motorcycles¹¹) except off-road transport (agricultural and industrial machinery, etc.). Road transport is the main source of air pollution in Georgia. The number of transport vehicles has doubled within the last decade.




Figure 3.4 Emissions of NOx and NMVOC from transport 2007-2021




Figure 3.5 Emissions of TSP, SOx and NH3 from transport 2007-2021

Emission trends of NOx, CO, TSP, NMVOC and NH₃ from this sector is gradually increasing (peaked in 2016) alongside growing number of vehicle in the country. From 2007 to 2021, emissions of NOx were increased by 102.7%, TSP by 51.8%, NMVOC by 88.5%, CO by 64.4% and emissions of NH₃ by 47.4 %.

¹⁰ In 2020-2021, emissions from gypsum production and in 2019 emissions from bricks production under 1A2f were calculated based on production data provided by GEOSTAT.

¹¹ 1Abiv is not estimated.

Reducing trend in emissions in 2016-2020 was a result of environmental policy in the transport sector. In particular, promotion of cleaner vehicles (hybrid and electric vehicles) and increased environmental taxes for the import of fuel engine and old vehicles. However, in 2021, due to sharp increase of vehicles import in post pandemic period and increased share of old and petrol/diesel engine cars in the import, emissions were increased.

Reduction of CO and low-level increase in NMVOC emissions in 2013 compared to 2012 were caused by reduction of petrol consumption due to switching of passenger cars from petrol fuel to compressed natural gas (CNG). Increasing emissions since 2014 are related to growing petrol consumption in these years caused by increasing car fleet.

Emissions of SO_x are gradually decreasing in parallel with reduction of sulphur content limits in national standards for petrol and diesel (for petrol: from 500 ppm to 10 ppm and for diesel: from 500 ppm to 50 ppm). In 2021, these are 9 times less compared to 2007.




Figure 3.6 Share of emissions from transport in total emissions in 2021

Transport was a source of 52% of NO_x emissions, 27% of NMVOC emissions and 56% of CO emissions in 2021.

Methodology

Road transport emissions are calculated by Tier 1 method of the EMEP/EEA Guidebook – 2019. Emissions from railways and national navigation (shipping) are estimated using EMEP/EEA Guidebook – 2019, Tier 1 approaches based on data from International Energy Agency (1990-2012) and from National Energy Balance (2013-2021).

Small combustion (1A4)

Source category description

Emissions in this category come from stationary combustion in commercial/institutional, residential and agriculture/forestry/fishing, plus from off-road vehicles and other machinery of agriculture/forestry/fishing.




Figure 3.7 Emissions of main pollutants from small combustion in 2013-2021




Figure 3.8 Emissions of particulate matters from small combustion in 2013-2021

Due to reduction in biomass and coal consumption in residential sector, emissions of NMVOC, SOx, NH₃, PMs and CO have decreased, however emissions increased in 2021 due to increased consumption of wood in the residential sector. Emissions of NOx remains stable for years since increased consumption of alternative fuels (mainly gas) still constitutes source for significant amount of NOx.




Figure 3.9 Share of emissions from small combustion in total emissions in 2021

PMs are mainly emitted from small combustion sector. In addition, the sector is responsible for 37% of CO, 16% of NMVOC and 8% of NOx emissions.




Figure 3.10 NOx emissions by sources of pollution in small combustion sector in 2021




Figure 3.11 NMVOC emissions by sources of pollution in small combustion sector in 2021




Figure 3.12 TSP emissions by sources of pollution in small combustion sector in 2021




Figure 3.13 CO emissions by sources of pollution in small combustion sector in 2021

Residential stationary combustion within small combustion sector is a main emission source of NO_x, NMVOC, SO_x, NH₃, CO and TSP. Also, 71% of Cd, 18% of Cu, 49% of Zn, 78% of dioxins/furans and 63% of PAHs are emitted from this subcategory.




Figure 3.14 Share of HMs and POPs emissions from residential combustion in total emissions in 2021

Methodology

Emissions are estimated according to EMEP/EEA Guidebook – 2019, Tier 1 approach based on the activity data from International Energy Agency (1990-2012) and National Energy Balance (2013-2021)¹².

Fugitive emissions from fuels (1B)

This category covers fugitive emissions from coal mining and handling, solid fuel transformation, oil and natural gas exploration.




Figure 3.15 Fugitive emissions of NMVOC, NOx, SOx, TSP and CO in 2009-2021

1B emits meaningful amount of NMVOC only, which is dependent on oil consumption and coal mining. Hence, emissions of NMVOC peaked in 2016 and again in 2021 because of increased oil consumption and coal mining. TSP emissions mainly comes from coal mining and handling. Emissions of NOx, SOx and CO from 2015 is related to resumed oil refining and gas flaring in oil refineries in Georgia. While significant reduction in TSP emissions is caused by decreased amount of produced coal.

Methodology

Emissions are estimated by EMEP/EEA Guidebook – 2019, Tier 1 approach based on the activity data from International Energy Agency (1B2av in 1990-2012), National Energy Balance (1B2aiv, 1B2av and 1B2c in 2013-2021) and from GEOSTAT (1B1a, 1B1b, 1B2ai, 1B2aiv in 1990-2012, 1B2b, 1B2c in 1990-2012).

4. Industrial processes and product use (NFR sector 2)

Dissolution of the Soviet Union accompanied with the collapse of the economy in the 1990s resulted in a significant decrease of industrial activities in Georgia. There has been some growth in this sector

¹² In case of 1A4bi liquid fuel data was taken from International Energy Agency for the years of 2013-2018.

in more recent years. The main activities in this sector are manufacturing of mineral products, chemical industry, metal production as well as paper, wood and food industries.




Figure 4.1 Emissions from industry sector in 2021

Share of chemical industry in total NO_x emissions in industry sector is 99.1%. Solvent subsector is responsible for 76% of NMVOC emissions. Manufacturing of mineral products is the major contributor of SO_x (69.2%) and TSP (82.9%) emissions from this sector.

Mineral Products (2A)

Source category description

In this category, cement, lime, bricks, concrete, gypsum, gravel and glass production are reported.




Figure 4.2 Emissions from mineral products 2007-2021

The most important pollutant emitted from this category is particulate matters. TSP emissions from 2A increased by 3-times in 2007-2021 and accounted for 34% of total TSP emissions in 2021. Increasing trend of this pollutant's emissions since 2008 is caused by rising amount gravel, concrete and gypsum

production. However, emissions of TSP decreased in 2020-2021 mainly due to reducing amount of gypsum and lime production.

Methodology

Emissions are estimated based on EMEP/EEA Guidebook – 2019, Tier 1 approach (2A1, 2A2, 2A3, 2A5) and national methodology (2A6). In case of cement production abatement coefficients were defined considering country and plant specifics. Activity data came from GEOSTAT, state reporting system for stationary sources and GHG emissions reports.

Chemical Industry (2B)

Source category description

This category covers emissions from ammonia, nitric acid and fertilizer production.




Figure 4.3 Emissions from chemical industry 2007-2021

Chemical industry emitted 10.3% of total NOx emissions in 2021. The main source from the category in this regard is nitric acid production (2B2) that decreased in 2020-2021 resulting reducing emissions of NOx. Significant decrease of emissions in 2016 was caused by switching to production data that is retrieved from state reporting system for stationary sources. The main source of TSP emissions from the sector is fertilizer production that is reported under 2B10a.

Methodology

Emissions are estimated based on EMEP/EEA Guidebook – 2019, Tier 1/2 approach. Activity data came from GEOSTAT (1990-2005), state reporting system for stationary sources (2016-2021), GHG emissions reports (2006-2013) and through extrapolation (2014-2015).

Metal Production (2C)

Source category description

In Georgia, there is ferroalloys and secondary iron/steel, lead and aluminum production.




Figure 4.4 Emissions from metal production 2007-2021

Ferroalloys production (2C2) is main source of TSP emissions, while iron and steel production is an only source for NMVOC in this category. Therefore, increased TSP emissions in 2016-2018 was caused by rising ferroalloys production in Georgia. Ferroalloys production sharply decreased in 2020 due to the effects of the coronavirus pandemic and reduced prices for the relevant materials. Production again increased in 2021 as a result of rising prices and demand for the product that conditioned growth in emissions of TSP.




Figure 4.5 Share of heavy metals emissions from iron and steel production in total emissions in 2021

Iron and steel production (2C1) is a key source of heavy metals emissions especially Pb (50%), As (58%) and Cr (77%). Aluminum production (2C3) is responsible for 99.8% of HCB emissions in Georgia.

Methodology

Emissions from are estimated using EMEP/EEA Guidebook – 2019, Tier 1/2 approaches. In case of aluminum and lead abatement coefficients were defined considering country and plant specifics. Activity data came from GEOSTAT, state reporting system for stationary sources and GHG emissions reports.

Solvents (2D)

Source category description

This category covers only three activities - road paving with asphalt, coating application and domestic solvent use including fungicides.




Figure 4.6 Emissions from solvents 2007-2021

NMVOC is the most important pollutant from this category as 24.7% of total emissions comes from 2D as of 2021. It increases due to rising quantity of applied paints year-by-year. Sharp increase in 2009 was caused by obtaining data on paints application that was unavailable before 2009.

Source of emissions of another important pollutant TSP is asphalt production. TSP emissions increases from 2014 as a result of increased production of asphalt for road paving.

Sharp reduction in emissions of both pollutant in 2020 is related to economic effects of the coronavirus pandemic and decreased consumption of the relevant products.

Methodology

Emissions from are estimated using EMEP/EEA Guidebook – 2019, Tier 1 approach. In case of road paving with asphalt abatement coefficients are defined considering country and plant specifics. Activity data came from GEOSTAT and state reporting system for stationary sources.

Other (2H, 2I and 2K)

Source category description

This category covers pulp and paper, food and drink industry, wood processing and consumption of POPs and heavy metals.

In Georgia, there is secondary paper processing only. Food comprises bread production, sugar production, margarine production, flour production, animal feed production, coffee processing, fish processing, meat processing. Under drink production, beer, wine, spirits and brandy are included.

In the past large wood processing companies existed in Georgia. Nowadays small plants remain which process logs and produce wooden boards etc.




Figure 4.7 Emissions from other industrial processes 2007-2021

Reductions of NMVOC and TSP emissions from this sector in 2008 related to the global economic crisis. Further reduction of NMVOC emissions since 2015 is caused by a sharp decline in sugar production, which was reduced to zero in 2017. Sharp increase in 2018 resulted from the reintroduction of sugar production. Rising NOx and SOx emissions from the category is caused by increased secondary paper processing.

Food and beverages industry is a responsible for 7.7% of NMVOC emissions as of 2021. Additionally, consumption of POPs and heavy metals is a source for 99.5% of total PCBs emissions.

Methodology

Emissions from are estimated using EMEP/EEA Guidebook – 2019, Tier 1 (2H1, 2I, 2K) and Tier 2 (2H2) approaches. In case of wood processing abatement coefficients are defined considering national methodology. Activity data came from GEOSTAT and state reporting system for stationary sources. Activity data to estimate emissions from consumption of POPs and heavy metals is a number of population in Georgia.

5. Agriculture (NFR sector 3)

Emission inventory from agriculture sector includes animal husbandry, the application of inorganic and organic fertilizers, farm-level agricultural operations and crops cultivation.




Figure 5.1 Emissions from agriculture sector in 2021

Agriculture sector is the main emitter of ammonia (87%) in the country. It also accounts for 17% of total NOx emissions and 18% of total NMVOC emissions as of 2021.




Figure 5.2 Emissions from agriculture 2007-2021

Sharp drop in emissions in 2014 is related to recalculations of activity data in agriculture sector by GEOSTAT in 2017 for the years of 2014-2015 based on the results of the census. Recalculations of previous years (2002-2013) were not implemented because a lack of human and financial resources in GEOSTAT. Emissions of ammonia have decreasing trend since 2014 mainly due to reducing production in animal husbandry and reduced levels of fertilizers consumption.

Manure Management (3B)

Source category description




Figure 5.3 Emissions from livestock manure management 2007-2021

Manure management is the most significant source of ammonia emissions as it accounts for 45% of total NH₃ emissions. It is also responsible for 97% of NMVOC and 87% of TSP emissions from agriculture.

Drop of trend in 2014 is related to recalculations of activity data in agriculture sector by GEOSTAT. Emissions increases in 2020 due to increased output of agricultural sector.

Methodology

Emissions are calculated using the EMEP/EEA Guidebook – 2019, tier 1 approach.

Agricultural Soils (3D)

Source category description

Under this category, NO_x and NH₃ emissions from application of inorganic and organic fertilizers and particulate matters emissions from farm-level agricultural operations (storage, handling and transport of agricultural product) are provided. Additionally, emissions of NMVOC have occurred from grain fields.




Figure 5.4 Emissions from use of inorganic N-fertilizers 2007-2021




Figure 5.5 Emissions from application of animal manure to soils 2014-2021




Figure 5.6 Emissions from urine and dung deposited by grazing animals 2007-2021




Figure 5.7 Emissions from farm-level agricultural operations including storage, handling and transport of agricultural products 2007-2021




Figure 5.8 Emissions of NMVOC from cultivated crops 2007-2021

Crop production and agricultural soils is a source for 42% of total ammonia emissions. Almost all (94%) NO_x emissions from agriculture comes from this subsector, which accounts for 16% of total NO_x emissions as of 2021. Subcategory - manure application to soils (3Da2a) - is only responsible for 9% of total NO_x and 24% of total NH₃ emissions. In addition, 54% of NO_x emissions and 27% of NH₃ emissions from agriculture comes from 3Da2a.

Drop of emissions in 2014 is related to recalculations of activity data in agriculture sector by GEOSTAT. Further decrease of NO_x emissions from use of inorganic N-fertilizers in 2017 (figure 5.3) are resulted by sharp reduction of use of fertilizers.

Methodology

Emissions are calculated using the EMEP/EEA Guidebook – 2019, tier 1 approach.

6. Waste (NFR sector 5)

This sector covers solid waste disposal on land, waste incineration and wastewater handling categories. The biggest polluting category in this sector is solid waste disposal on land from where comes about 99.7% of NMVOC emissions.




Figure 6.1 Emissions of NMVOC from waste sector in 2021 (kt)




Figure 6.2 Emissions of NOx, SOx and TSP from waste sector in 2021

Solid waste disposal on land (5A)

Source category description




Figure 6.3 Emissions from solid waste disposal on land 2014-2021

In general, emissions of NMVOC from solid waste disposal is rising due to increased accumulation of wastes on landfills.

Methodology

Emissions are calculated using EMEP/EEA Guidebook – 2019, Tier 1 approach. Data on CH₄ emissions from solid waste disposal on land were obtained from Georgia's Biennial Update Reports (BUR) to the UNFCCC. Emissions for 2018-2020 were extrapolated, since data on CH₄ emissions in BUR is available until 2018.

Waste incineration (5C)

Source category description

This category includes industrial waste incineration and clinical waste incineration. Due to lack of activity data emissions from this category were estimated only from 2013.




Figure 6.4 Emissions from waste incineration 2014-2021

Increased emissions in this category is related to installing of new incinerators and consequently, increased amount of waste incinerated.

Methodology

Emissions are estimated based on plant specific emissions (from state reporting system for stationary sources).

Wastewater handling (5D)

Source category description

This category covers industrial domestic wastewater handling and industrial wastewater handling.




Figure 6.5 Emissions from wastewater handling 2015-2021

Until 2012 wastewater treatment infrastructure in Georgia had been gradually demolished. Since 2012 new WWTPs were built and in parallel, amount of treated wastewater were increased. As a result, emissions of MNVOC from this category started increasing and doubled in 2011-2021.

Due to lack of activity data on a number of population not connected to centralized wastewater collection system NH_3 emissions from this category were estimated only from 2015.

Methodology

Emissions are calculated using EMEP/EEA Guidebook – 2019, Tier 1 approach. Activity data were gained from the Integrated Management Division of NEA.

7. Recalculations and improvements

Recalculations

Activity data of 1A2b was corrected for 2020 as it was indicated in wrong unit.

Emissions from 1A3bv-1A3bvii in 2004-2006 and in 2017-2021 were recalculated because relevant activity data were recalculated by GEOSTAT.

Emissions of TSP from 2A6 in 2001-2020 were recalculated due to estimation of emissions from gypsum production that had not been taken into account in previous submissions.

Emissions from 2H2 in 2020 were slightly changed once coffee production data had been reported incorrectly before.

Emissions of NO₂ from 3DA2a were recalculated for 2014-2020 and calculated for 1990-2013 according to the technical correction suggested by ERT in 2023.

Planned improvements

For the next year, it is planned to calculate emissions from aviation and construction and demolition, and recalculate emissions from road transport using more modern software tool COPERT 5.

8. IIR References

1. Mareckova, K.; Wankmüller, R.; Moosmann, L.; Pinterits, M.; Tista, M. Inventory Review 2014. Review of emission data reported under the LRTAP Convention and NEC Directive. Stage 1 and Review. Status of gridded and LPS data. Technical Report CEIP 2014/1.
http://www.ceip.at/review_proces_intro/review_reports/;
2. UNECE, 2014. Guidelines for Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution (ECE/EB.AIR/125);
3. Annexes to UNECE reporting guidelines (ECE/EB.AIR/125)
http://www.ceip.at/fileadmin/inhalte/emep/2014_Guidelines/Annexes_revised_140417.zip
4. UNECE, 2013. Decision 2013/4 reporting of emissions and projections data under the Convention and its protocols in force (ECE/EB.AIR/122/Add.1.);
5. # 435 Order of the Government on instrumental method for determination of actual amounts of emissions into ambient air from stationary pollution source, standard list of emission measuring equipment, and methodology for calculation of actual amounts of emissions into ambient air from stationary pollution source according to technological processes (31/12/13);
6. EMEP/EEA air pollutant emission inventory guidebook – 2019:
<http://www.eea.europa.eu/publications/emep-eea-guidebook-2019>;
7. Energy Balance of Georgia, 2013;

8. Energy Balance of Georgia, 2014;
9. Energy Balance of Georgia, 2015;
10. Energy Balance of Georgia, 2016;
11. Energy Balance of Georgia, 2017;
12. Energy Balance of Georgia, 2018;
13. Energy Balance of Georgia, 2019;
14. Energy Balance of Georgia, 2020;
15. Energy Balance of Georgia, 2021;
16. AGRICULTURE OF GEORGIA - Statistical Publications 2007;
17. AGRICULTURE OF GEORGIA - Statistical Publications 2008;
18. AGRICULTURE OF GEORGIA - Statistical Publications 2009;
19. AGRICULTURE OF GEORGIA - Statistical Publications 2010;
20. AGRICULTURE OF GEORGIA - Statistical Publications 2011;
21. AGRICULTURE OF GEORGIA - Statistical Publications 2012;
22. AGRICULTURE OF GEORGIA - Statistical Publications 2013;
23. AGRICULTURE OF GEORGIA - Statistical Publications 2014;
24. AGRICULTURE OF GEORGIA - Statistical Publications 2015;
25. AGRICULTURE OF GEORGIA - Statistical Publications 2016;
26. AGRICULTURE OF GEORGIA - Statistical Publications 2017;
27. AGRICULTURE OF GEORGIA - Statistical Publications 2018.
28. AGRICULTURE OF GEORGIA - Statistical Publications 2019.
29. AGRICULTURE OF GEORGIA - Statistical Publications 2020.
30. AGRICULTURE OF GEORGIA - Statistical Publications 2021.
31. GEOSTAT, H-3. Composition of Road Motor Vehicle Fleet by Fuel Type:
https://geostat.ge/media/48286/H-3.-Composition-of-road-motor-vehicle-fleet-by-fuel-type_ENG.XLS
32. International Energy Agency, Energy data and statistics: <https://www.iea.org/data-and-statistics/data-browser?country=WORLD&fuel=Energy%20supply&indicator=TPESbySource>
33. Georgia's 2nd Biennial Update Report:
https://unfccc.int/sites/default/files/resource/2019.06.13_BUR2_2019_Eng.pdf